

Qualitative Evaluation of Land Suitability for Olive, Potato and Cotton Cultivation in Tarom in Zanjan

Zahra Mirzae Shiri*, Manochehr Farbodi

Soil Science Department, Agriculture Faculty, Miyaneh Branch-Islamic Azad University, Miyaneh, Iran

*Corresponding author: Zahra Mirzae Shiri, Email: zahramirzaieshiri@yahoo.com

Submitted: July 28, 2020; Revised: December 15, 2020, February 1, 2021, February 17, 2021;

Accepted: March 13, 2021

ABSTRACT

The use of soil as one of the main factors in the manufacture of agricultural products for human consumption must be carried out using the right basis and scientific principles. One of the ways to achieve this important goal is to assess land suitability for common crops in the study area. It is also necessary to determine product compatibility based on the soil and climate conditions to reduce the risk of investment because there is no history of the studied plant's cultivation in the region. Therefore, this study aims to evaluate the suitability of land for cotton, potato, and olive crops at the bottom of the valley in Tadarom in Tarom region, Zanjan province to ensure proper use planning. After sampling and performing physical, chemical, and profile tests using the classification key of 2014, the soils were classified into different Aridi Sols categories, namely Typic calcigypsid, haplocambid, and torriortent. The climate, soil, and terrain data of the area were then collected from the relevant agencies. Land suitability assessment for olive, potato, and cotton was also performed using the simple constraint, parametric, FAO system methods. The evaluation results showed that the Tahdareh area with series 1 soils was in the not suitable (N) and subclass (Ns) categories using the history and second root methods, respectively with depth and pebbles as the limiting factors. Meanwhile, soil series 2, 3, 4, and 5 were placed in the critical fit (S_3) and proportionality (S_2) classes by the story and second root methods, respectively, where they were hindered by drainage. Evaluation of these lands for potato crop by story method for series 1, 2 and 3 shows that they were in Ns class, while 4 was in the proportionality (S_3) category with lime as the limiting factor. The non suitable class was obtained using the story method (N) for series 2, 3, 4 and 5 and they were hindered by soil depth, while class S_3 was obtained for all the soils with the texture and gravel as limit factors. The results showed that the studied lands were relatively suitable for olive groves and the conditions can be improved with corrective factors for drainage and the amount of gravel. The storytelling method has very strict output while presenting its results. The study area was considered a land with critical fit (S_3) class for olive cultivation, but it was not very consistent with field observations. Furthermore, the second root method had the greatest compliance with the existing realities in the region and it can be considered the most reliable parametric technique in the qualitative assessment of land suitability.

Keywords: Land evaluation; land use; parametric method; quality proportion

INTRODUCTION

Over the years, humans have made several efforts to achieve the optimal usage of critical resources around the world. One of the most important natural resources used is soil, which serves as the basis of all agricultural activities. Human and animal life also depends on its level of exploitation, where excessive usage causes degradation. Therefore, soil utilization must be carried out carefully to avoid damage to valuable resources while reaching maximum production (Kilic *et al.*, 2005).

The increase in population and food demand as well as the subsequent degradation of these materials have led to various crises in the modern era (ZaliVargahan *et al.*, 2001; Santana-Cordero *et al.*, 2016; Hanh *et al.*, 2017). The inappropriate use of land is considered one of the most influential destructive factors of natural resources (Lal, 2009; Brevik, 2013). At present, land suitability classification has been considered (Hudson and Birnie, 2000) as a basic solution to certain problems, such as population growth, fertility reduction (Panagos *et al.*, 2015), soil and water pollution (Chartzoulakis and Bertaki, 2015; Sam *et al.*, 2016). Their potential must be evaluated to ensure they are assigned to the most profitable and sustainable operating system.

Land evaluation studies the potential of an area for its intended uses before any application for production. Furthermore, it assesses two important aspects, namely the physical aspects covering soil, topography, climate as well as the socio-economic aspects, such as land size, management level, labor availability and market access. Over time, the physical characteristics of the earth are almost constant, while the socio-economic factors have high variations. The main aim of land suitability assessment is to study the physical, social and economic aspects of land for optimal and sustainable use (Norton, 2003).

The FAO assessment system can fulfill the aforementioned objectives as an applied approach. Furthermore, while providing an optimal cropping pattern, it can also predict the productivity of each crop and provide guidance on the type of management required. One of its advantages is the adherence to a single framework proposed by the World Food Organization in 1976 for global implementation. This system allows the possibility to exchange and utilize the results of studies around the world as well as to collaborate with other countries (Ashraf and Normohammad, 2011).

Assessing land based on its suitability as well as agricultural and natural resources evaluation helps to reveal its potential and productivity patterns. Evaluation is often carried out to study its characteristics and economic conditions, which helps to achieve the proper

use of resources without soil degradation (FAO, 1976, 2007; Niekerk, 2010). One of the ways to optimally use land and prevent destruction is to carry out activities based on its production capacity (Rossiter, 1996; Anaya-Romero *et al.*, 2015).

Land assessment is often performed to evaluate the relationship between the land and its use type. It also helps to determine how the area can be properly utilized as well as to provide an estimate of the required amount of inputs and outputs (Sys *et al.*, 1993). In most developing countries, qualitative suitability assessment is a central focus of land evaluation including suitability studies for potato (Caldiz *et al.*, 2001), tomato (Shepande, 2002), cotton (Kalogirou, 2002) and olive (Wahba *et al.*, 2007).

The sources and capacity of land must be assessed before planning its best usage purpose. Therefore, recognizing the production capacity and allocating them to the best and most profitable type of use as well as paying attention to soil fertility and protection are very important. In evaluation studies, the capacity is often studied and determined based on the type of use. The first step in usage planning is land suitability assessment, where the most important task is understanding the relationship between the natural environment and the types of possible uses.

The problems of land misuse, uncontrolled growth of urban areas and declining area due to cultivation have gained public attention around the world. For policymakers, appropriate and optimal land use methods are very important for planning and effective management of agriculture in the region. The study of soil quality indicators in different areas provides the possibility of better utilization. Therefore, this study aims to evaluate the suitability of lands based on the FAO model using simple and parametric constraints as well as to investigate some soil quality indicators in the valley lands of the Tarom region in Zanjan province for olive, cotton and potato cultivation.

MATERIALS AND METHODS

This study was carried out in Tarom city, Zanjan province, Iran, and the total land used was 570 hectares. Based on the meteorological information of the water synoptic station, the average maximum daily temperature is 36.2°C in August with an average minimum daily temperature of 2.1°C in February. The maximum relative humidity of 70% is often experienced in December and the minimum of 61% in July and February. The absolute maximum and minimum temperatures were 48.5°C in August and -9.5°C in January, respectively. Furthermore, the area had an

the properties. When the amount of lime decreases with depth, it is used up to a depth of 30 cm for evaluation, or it is calculated using weighted coefficients of the average amount to the root penetration depth. Furthermore, the irrigation of calcareous soils causes changes in their physical properties as the irrigated layers harden and soil permeability to the root reduces. The high amount of lime in the soil also causes a reduction in its water storage capacity. It is important to note that low amounts have positive effects on plant nutrition and soil structure improvement.

In the assessment of gypsum status, when its average amount decreases with depth, a value of 30 cm is considered or the evaluation is carried out based on the average value in the root depth and weighting factors. Furthermore, the mean soil pH was calculated for a depth of 25 cm. Using the weighted average coefficients, the average EC was calculated in Decisiemens per meter for olive and potato up to depths of 150 cm and 100 cm, respectively.

Method of Classifying Lands

Depending on the extent to which land specifications can meet the needs of the crop, its suitability is expected to vary. Consequently, the value of characteristics for a particular plant is very important in the evaluation process. In this study, two methods were used to determine land class: simple restriction method and parametric method.

Simple restriction method

In this method, the most limited land characteristic for crop growth determines the suitability class. To check the meteorological information and determine the climatic compatibility, the weather data for Abar city station, namely 10 km from the study area was used. The studied climatic parameters include rainfall, temperature, relative humidity and sunlight. The results obtained from the calculation of weather data were compared with the needs of the product, hence, the class depends on each climatic variable. The lowest category obtained was assumed to be the effect of weather. The variables related to geography and soil produced another class, subsequently the lowest was considered the total land class.

Parametric method

In the parametric method, a quantitative rating was assigned to each attribute, and when they are desirable for the target crop, the maximum rating of 100 is used. Meanwhile, when the same attribute is restricted, a lower value was assigned, after which the degrees were then used to calculate the land index.

Climate evaluation was also carried out based on variables of temperature, rainfall, radiation and relative humidity. The lowest grade assigned to each group was selected and used by the Storie and Root Secondary Climate Index methods (Equation 1).

$$25 < CI < 92.5 \quad CR = 16.67 + 0.9 \times CI \quad (1)$$

$$CI < 25 \quad CR = 1.6 \times CI \quad (2)$$

The land index was calculated using the degrees assigned to each land attribute with the Equation 3.

$$I \times \left(\frac{C}{100} \right) = A \times \left(\frac{B}{100} \right) \quad (3)$$

Where I = Storie index and A, B, C, etc. are the degrees.

This method uses the Khidir equation presented in 1986 and known as the second root, which minimizes the interactions between the factors to some extent (Equation 4).

$$I = R_{min} \sqrt{\left(\left(\frac{A}{100} \right) \times \left(\frac{B}{100} \right) \times \dots \right)} \quad (4)$$

Where I = Storie index, Rmin: the lowest degree between the different properties, and A, B and ... other degrees of attributes apart from the lowest.

Using the calculated index, the land suitability class was determined as very good grade (S_1), relatively good (S_2), critical fit (S_3) and not-suitable (N) with value ranges of 100-75, 75-50, 25-25 and 0-25, respectively.

RESULTS AND DISCUSSION

Evaluation of soil properties

The results of soil properties assessment using physical and salinity properties in different series and units were presented for olive plants (Tables 1 and 2), potatoes (Tables 3 and 4) and cotton (Tables 5 and 6). Furthermore, the average soil texture profile in series 1 sandy loam, series 2 sandy loam, series 3 sandy clay loam soil, series 4 clay loam sand and series 5 clay loam were determined based on the mean values of sand, silt and clay as well as the texture triangle. The amount of pebbles in series 1 were 15-35 %, >75 %, >75 %, 15-35 %, 35-75%. Meanwhile, value range of 15-35% was obtained in series 2, 3, 4 with 15-30% in series 5, as shown in Table 1. Based on the results, series 2 and 3 have soil depths of 120 cm, while values of 20 cm, 130 cm and 120 cm were obtained in 1, 4 and 5, as shown in Tables 1, 3 and 5. At these levels, series 1, 2, 3, 4 and 5 have lime content of 14.7, 17.90,

16.16, 7.5 and 27.5, respectively, as shown in Tables 1 and 6. For gypsum, a similar approach was also used, but its content was not reported in the studied profile. Electrical conductivity of soil saturated extract was used to evaluate its salinity. In the study area, a mean value of 0.9 dS m⁻¹ was obtained for different soil series under olive, potato and cotton cultivations, as shown in Tables 1, 3 and 5. Salinity values of 0.7, 0.85, 0.9 and 1 dS m⁻¹, were recorded in units 2, 3, 4 and 5 under olive cultivation, respectively, as shown in Table 2.

Final Land Evaluation

The land evaluation value was calculated by combining soil and climate assessment results. Calculation of land index was carried out the methods in Equation 5 and 6.

$$LI = \left(CR \times \frac{BC}{100} \times \frac{C}{100} \right) \quad (5)$$

$$LI = R_{min} \sqrt{\left(\left(\frac{A}{100} \right) \times \left(\frac{B}{100} \right) \times 100 \right)} \quad (6)$$

C and B are related to the soil, while CR is associated with climate. R_{min} is the product of the lowest degree assigned to any soil properties multiplied by the root of the product of the others. The results showed that the lowest climatic index and climatic degree were obtained from olive, while the highest amount of indices were recorded in potato. Table 7 shows the final evaluation of the qualitative suitability of the lands in the Tahdara area using the parametric method, which consists of storie and second root techniques for the cultivation of olive, potato and cotton plants.

Climate characteristics assessment

The climatic characteristic for the target crop were obtained based on the standard tables presented by conveyor Givi (1997) as well as the results of soil analysis and related evaluation experiments. In the parametric method, the lowest degree assigned to each group was selected, and then converted to climate index by using storie and second root methods. Subsequently, the appropriate equations for the climatic degree were obtained. The final result of evaluation for olive using the simple and parametric constraint method revealed that the land was in S_2 , while S_1 , S_2 and S_1 were obtained for potato with the simple constraint, Storie and second root methods, respectively. Furthermore, the class of S_2 was recorded for cotton using the simple constraint, storie and the second root methods, as shown in Figure 7. After determining the degrees in the climate needs,

the lowest in each group was then selected. They were then calculated along with the climate index using the storie and second root method, as shown in Table 7.

Qualitative land suitability assessment provides useful information about different crops and the most suitable place to produce them. The results showed that the lowest index and climatic grade were obtained from the olive plant. Furthermore, Mashayekhi (2015) qualitatively and quantitatively evaluated the suitability of land for local crops in the Khodafarin region. The results of climatic evaluation using the simple method and severity showed low suitability S_3 for cotton cultivation in this area. The main limiting factors were high gypsum, pH and unsuitable weather. Rahimi *et al.* (2009) assessed olives on a 2,200ha plot in Roodbar with 6 soil series and phases, where the results showed that soils can be classified into two categories, namely Aridisols and Entosol/Torriorthents. Qualitative evaluation was carried out in the study using the limitation and parametric methods. Olive production was based on regional cultivation management practices in the Roodbar region. The results showed that the climate conditions are very suitable (S_1), but there were important topographic limiting factors, such as large stones, shallow soils, salinity and alkalinity issues. Menjiver *et al.* (2003) conducted a study on land suitability for olive plants in Spain, where 35 research profiles were selected and evaluated using 6 methods. They revealed that according to the FAO method, the most limiting factors in this region were high soil moisture and extreme slope. In the system, all of the studied areas were in the suitable class N_1 and the reduction of any extreme conditions required a lot of expenditure. Similarly, Igue *et al.* (2005) used the same method to evaluate the potential of Azucan lands in West Africa for flax cultivation. Their results showed that soil index for different soils in the area ranged from critical to unsuitable, while the limiting factors were methane production, adverse climatic conditions, physical and chemical properties, soil texture and organic carbon. Al-Mashreki *et al.* (2011) investigated land suitability for rainfed sorghum using GIS and remote sensing technologies in IBI province in southwest Yemen. The results revealed that approximately 5% of the study area was very suitable for sorghum cultivation. Meanwhile, 25%, 31%, 24% and 15% were relatively suitable, critical, inappropriate and temporarily unsuitable, respectively.

Adaptation of climatic, soil and topography needs with parametric method was carried out. Furthermore, Table 7 shows the results of the process including climate index and class, land index as well as final qualitative class. The use of the storie and second root

Table 1. Soil properties evaluation results of different Unit 1 series for olives

Features	series 1/1			series 1/2			series 1/3			series 1/4			series 1/5		
	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value
Slope (%)	96	S ₁	5-2	98	S ₁	2-0	96	S ₁	5-2	96	S ₁	5-2	96	S ₁	5-2
Flooding	85	S ₂	None	100	S ₁	None	100	S ₁	None	85	S ₂	None	85	S ₂	None
Drainage	85	S ₂	Average	72.5	S ₂	Average	72.5	S ₂	Average	85	S ₂	Average	85	S ₂	Average
Texture and structure	100	S ₁	SL	100	S ₁	SL	100	S ₁	SL	100	S ₁	SL	100	S ₁	SL
Pebbles (vol %)	75	S ₁	15-35	0	N ₂	>%75	0	N ₂	75-35	75	S ₁	15-35	0	N ₂	35-75
Depth (cm)	0	N ₂	20	0	N ₂	20	0	N ₂	20	0	N ₂	20	0	N ₂	20
Lime (%)	100	S ₁	14.7	100	S ₁	14.7	100	S ₁	14.7	100	S ₁	14.7	100	S ₁	14.7
Gypeum (%)	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
Organic matter (%)	72.5	S ₁	1.86	72.5	S ₁	1.86	72.5	S ₁	1.86	72.5	S ₁	1.86	72.5	S ₁	1.86
EC (ds m ⁻¹)	95.96	S ₁	0.9	95.56	S ₁	0.9	95.56	S ₁	0.9	95.56	S ₁	0.9	95.56	S ₁	0.9

Table 2. Soil properties evaluation results for Units 2 to 5 for olives

Features	Unit 2			Unit 3			Unit 4			Unit 5		
	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value
Slope (%)	98	S ₁	2-0	96	S ₁	2-5	96	S ₁	5-2	96	S ₁	5-2
Flooding	100	S ₁	None	85	S ₂	None	100	S ₁	None	85	S ₂	None
Drainage	72.5	S ₂	Average	85	S ₂	Average	72.5	S ₂	Average	85	S ₂	Average
Texture and structure	100	S ₁	SL	98	S ₁	LCS	100	S ₁	SL	100	S ₁	SL
Pebbles (vol %)	75	S ₁	15-35	75	S ₁	15-35	0	N ₂	75-35	75	S ₁	15-35
Depth (cm)	85.33	S ₁	120	85	S ₁	120	0	N ₂	20	0	N ₂	20
Lime (%)	100	S ₁	17.90	100	S ₁	16.16	100	S ₁	14.7	100	S ₁	14.7
Gypeum (%)	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
Organic matter (%)	85.27	S ₁	0.82	100	S ₁	2.4	72.5	S ₁	1.86	72.5	S ₁	1.86
EC (dS m ⁻¹)	95.47	S ₁	0.7	100	S ₁	0.85	95.56	S ₁	0.9	95.56	S ₁	0.9

Table 3. Results of soil properties assessment of different unit 1 for potatoes

Features	series 1/1			series 1/2			series 1/3			series 1/4			series 1/5		
	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value
Flooding	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
Drainage	85	S ₁	Average	85	S ₁	Average	85	S ₁	Average	85	S ₁	Average	85	S ₁	Average
Texture and structure	95	S ₁	SL	95	S ₁	SL	95	S ₁	SL	95	S ₁	SL	95	S ₁	SL
Pebbles (vol%)	72.5	S ₂	15-35	50	S ₃	35-75	50	S ₃	35-75	72.5	S ₂	15-35	50	S ₃	35-75
Depth (cm)	60	S ₃	20	60	S ₃	20	60	S ₃	20	60	S ₃	20	60	S ₃	20
Lime (%)	98.4	S ₁	14.7	98.4	S ₁	14.7	98.4	S ₁	14.7	98.4	S ₁	14.7	98.4	S ₁	14.7
Gypeum (%)	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
pH	7.9	S ₁	7.76	7.9	S ₂	7.76	7.9	S ₂	7.76	100	S ₂	7.76	79	S ₂	7.76
EC (dS m ⁻¹)	99.5	S ₂	0.9	99.5	S ₁	0.9	99.5	S ₁	0.9	85	S ₁	0.9	99.5	S ₁	0.9

Table 4. Results of soil properties assessment of units 2 to 5 for potatoes

Features	Unit 2			Unit 3			Unit 4			Unit 5		
	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value
Flooding	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
Drainage	85	S ₁	Average	85	S ₁	Average	85	S ₁	Average	85	S ₁	Average
Texture and structure	95	S ₁	SL	100	S ₁	LCS	100	S ₁	LCS	95	S ₁	LCS
Pebbles (vol%)	72.5	S ₂	15-35	72.5	S ₂	15-35	72.5	S ₂	15-35	95	S ₁	15-35
Depth (cm)	100	S ₁	120	100	S ₁	120	100	S ₁	130	100	S ₁	120
Lime (%)	40	N ₁	25.25	40	N ₁	24.95	97.68	S ₁	12.12	72.4	S ₂	42.44
Gypeum (%)	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
pH	82	S ₂	7.85	80.5	S ₂	7.82	71.5	S ₂	7.46	75.75	S ₂	7.63
EC (dS m ⁻¹)	98	S ₁	0.94	87.02	S ₁	1.40	88.35	S ₁	1.67	85.7	S ₁	1.14

Table 5. Results of soil properties assessment of different unit series 1 for cotton

Features	series 1/1			series 1/2			series 1/3			series 1/4			series 1/5		
	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value
Slope (%)	98	S ₁	2-5	99	S ₁	0-2	98	S ₁	2-5	98	S ₁	2-5	98	S ₁	2-5
Flooding	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
Drainage	100	S ₁	Average	100	S ₁	Average	100	S ₁	Average	100	S ₁	Average	100	S ₁	Average
Texture and structure	60	S ₃	SL	60	S ₃	SL	60	S ₃	SL	60	S ₃	SL	60	S ₃	SL
Pebbles (vol %)	95	S ₁	15-35	100	S ₂	35-75	25	N ₂	35-75	72.5	S ₂	15-35	25	N ₂	35-75
Depth (cm)	0	N ₂	20	0	N ₂	20	0	N ₂	20	0	N ₂	20	0	N ₂	20
Lime (%)	89.7	S ₁	14.7	89.7	S ₁	14.7	89.7	S ₁	14.7	89.7	S ₁	14.7	89.7	S ₁	14.7
pH	93.6	S ₁	7.76	93.6	S ₁	7.76	93.6	S ₁	7.76	93.6	S ₁	7.76	93.6	S ₁	7.76
EC (dS m ⁻¹)	100	S ₁	0.9	100	S ₁	0.9	100	S ₁	0.9	100	S ₁	0.9	100	S ₁	0.9

Table 6. Soil evaluation results of units 2 to 5 for cotton

Features	Unit 2			Unit 3			Unit 4			Unit 5		
	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value	Numeric method	Limitation	Value
Slope %	99	S ₁	0-2	98	S ₁	2-5	99	S ₁	2-0	99	S ₁	2-0
Flooding	100	S ₁	None	100	S ₁	None	100	S ₁	None	100	S ₁	None
Drainage	100	S ₁	Average	100	S ₁	Average	100	S ₁	Average	100	S ₁	Average
Texture and structure	60	S ₃	SL	85	S ₂	SCL	85	S ₂	SCL	85	S ₂	SCL
Pebbles (vol %)	72.5	S ₂	15-35	72.5	S ₂	15-35	90	S ₁	3-15	72.5	S ₂	15-35
Depth (cm)	98	S ₁	120	98	S ₁	120	98	S ₁	120	91	S ₁	130
Lime (%)	94.72	S ₁	19.72	93.56	S ₁	18.56	99.69	S ₁	9.38	45.1	S ₃	32.55
pH	95	S ₂	7.88	87.5	S ₂	7.82	89.6	S ₁	7.46	89.6	S ₁	7.63
EC (dS m ⁻¹)	100	S ₁	0.7	100	S ₁	0.85	95.96	S ₁	0.9	95.6	S ₁	1.0

Table 7. Parametric method for qualitative suitability of cropland for olive, potato, cotton cultivation

Plant	Soil series	Climate index		Climate degree		Land index		Land class	
Olive	1.4	62.5	62.5	72.92	72.92	0	0	N _s	N _s
Olive	1.5	62.5	62.5	72.92	72.92	0	0	N _s	N _s
Olive	2	62.5	62.5	72.92	72.92	26.99	26.44	S _{3w}	S _{3w}
Olive	3	62.5	62.5	72.92	72.92	42.13	42.55	S _{3c}	S _{2c}
Olive	4	62.5	62.5	72.92	72.92	32.79	57.10	S _{3w}	S _{2w}
Olive	5	62.5	62.5	72.92	72.92	36.50	60.24	S _{3w}	S _{2w}
Potato	1.1	72.93	79.1	82.31	87.86	22.36	40.2	N _s	S _{3s}
Potato	1.2	72.93	79.1	82.31	87.86	15.42	30.5	N _s	S _{3s}
Potato	1.3	72.93	79.1	82.31	87.86	15.42	18.5	N _s	N _s
Potato	1.4	72.93	79.1	82.31	87.86	22.36	40.37	N _s	S _{3s}
Potato	1.5	72.93	79.1	82.31	87.86	15.42	30.60	N _s	S _{3s}
Potato	2	72.93	79.1	82.31	87.86	15.48	27.2	N _s	S _{3s}
Potato	3	72.93	79.1	82.31	87.86	14.21	26.28	N _s	S _{3s}
Potato	4	72.93	79.1	82.31	87.86	31.29	52.14	S _{3f}	S _{2f}
Potato	5	72.93	79.1	82.31	87.86	29.67	25.44	S _{3s}	S _{3s}
Cotton	1.1	72	72	81.47	81.47	0	0	N _s	N _s
Cotton	1.2	72	72	81.47	81.47	0	0	N _s	N _s
Cotton	1.3	72	72	81.47	81.47	0	0	N _s	N _s
Cotton	1.4	72	72	81.47	81.47	0	0	N _s	N _s
Cotton	1.5	72	72	81.47	81.47	0	0	N _s	N _s
Cotton	2	72	72	81.47	81.47	30.93	47.4	S _{3s}	S _{3s}
Cotton	3	72	72	81.47	81.47	39.47	58.72	S _{3s}	S _{2s}
Cotton	4	72	72	81.47	81.47	38.60	58.61	S _{3s}	S _{2s}
Cotton	5	72	72	81.47	81.47	26.02	35.89	S _{3s}	S _{3s}

method for different indices gave similar results. The evaluation showed that the Tahdare zone for olive cultivation on series 1 soil was in the not suitable (N) and not subclass (N_s) categories using the storie and second root methods, respectively. The limitation faced in the area was related to soil depth and pebbles. In series 2, 3, 4 and 5, the soil was critically proportional (S₃) and the constraint was drainage except for series 2, which was hindered by climate. Evaluation of these lands on potato crop was based on the storie method for series 1, 2 and 3, which were in the subclass (N_s), and the lime was the limiting factor. For cotton cultivation in soil series 1, the land class was suitable (N), with soil depth as the constraint. Meanwhile, in series 2, 3, 4 and 5, the class of proportion was S₃, with limiting factors of soil texture and pebbles, as shown in Table

7. Elaalem (2013) compares two parametric and fuzzy methods for evaluating olive land suitability in the Jay Farah plain of Libya. The fuzzy method indicated that the majority of the study area was suitable, while the parametric method showed moderate suitability for the plant's production. Mahari and Alebachew (2013) also investigated the ability of some Ethiopian lands using the parametric method. The result showed that the most important limiting factors in the area were slope and soil physical properties, especially texture and depth. Magboul *et al.* (2015) evaluated land suitability for citrus and crop production around the Nile, North Khartoum and Sudan. The results showed that the suitability class was poor and not suitable for the crop. Furthermore, the most important factors limiting factors drainage were soil salinity and erosion. Based on these

findings, the storie method often gives more unrealistic as well as strict results and the estimates also show unsuitable class.

CONCLUSION

The evaluation results showed that the Tahdareh area for cultivation of olives and cotton in series 1 soils using the history and second root methods had an unsuitable class, with depth and gravel as the limiting factors. Series 2, 3, 4 and 5 were placed in the critical fit class S_3 with the story method, and it was hindered by drainage. However, in the second root method for the above series, a fit class S_2 was obtained for the olive crop, with soil drainage as the limiting factor. The evaluation of these lands was unsuitable for potato cultivation using the history method on series 1, 2 and 3. Suitability class S_3 was obtained for 4 with high soil lime as the constraint. The results showed that the lands assessed for olive orchards and cotton cultivation were in the critical class S_3 , and the removal of the drainage restrictions made it suitable. Most of the areas studied were not suitable for growing potatoes and their cultivation is not recommended.

CONFLICT OF INTEREST

The authors declare that this article is an original study that has not been published in another journal, and there are no conflict of interest

REFERENCES

- Al-Mashreki, M., Bin Mat Akhir, J., Abd Rahim, S., Md. Desa, K., Lihan, T. and Haider, A. (2011). Land suitability evaluation for sorghum crops in the Ibb Governorate, republic of Yemen using Remote Sensing and GIS techniques. *Aust. J. Basic Appl. Sci.*, 5(3): 359-368.
- Anaya-Romero, M., Abd-Elmabod, S. K., Muñoz-Rojas, M., Castellano, G., Ceacero, C. J., Alvarez, S., Méndez, M. and De la Rosa, D. (2015). Evaluating soil threats under climate change scenarios in the Andalusia region. *Southern Spain Land Degrad. Dev.*, 26: 441-449. <https://doi.org/10.1002/ldr.2363>
- Ashraf, S. and Normohammadan, B. (2011). Qualitative evaluation of land suitability for wheat in Northeast-Iran using FAO methods. *Indian J. Sci. Technol.*, 4(6): 703-707. [10.17485/ijst/2011/v4i6.15](https://doi.org/10.17485/ijst/2011/v4i6.15)
- Brevik, E. C. (2013). The potential impact of climate change on soil properties and processes and corresponding influence on food security. *Agriculture.*, 3: 398-417. <https://doi.org/10.3390/agriculture3030398>
- Caldiz, D. O., Gaspari, F. J., Haverkort, A. J. and Struik, P. C. (2001). Agro-ecological zoning and potential yield of single or double cropping of potato in Argentina. *Agric. Forest Meteorol.*, 109: 311-320. [https://doi.org/10.1016/S0168-1923\(01\)00231-3](https://doi.org/10.1016/S0168-1923(01)00231-3)
- Chartzoulakis, K. and Bertaki, M. (2015). Sustainable water management in agriculture under climate change. *Agric. Sci. Procedia.*, 4: 88-98. <https://doi.org/10.1016/j.aaspro.2015.03.011>
- Elaalem, M. (2013). A Comparison of parametric and fuzzy multi-criteria methods for evaluating land suitability for olive in jeffara plain of Libya. *APCBEE Procedia.*, 5: 405-409. <https://doi.org/10.1016/j.apcbee.2013.05.070>
- FAO. (1976). A framework for land evaluation. *FAO Soils Bulletin, NO 32, Rome, Italy.* 72 p. <http://www.fao.org/3/x5310e/x5310e00.htm>
- FAO. (2007). Land Evaluation: Towards a Revised Framework; *Land and Water Discussion Paper 6; FAO: Rome, Italy*, p. 107. DOI: 10.1016/j.geoderma.2008.11.001
- Givi, J. (1997). Qualitative assessment of land suitability for crops and orchards. Soil and Water Research Institute. Tehran. <http://www.swri.ir/en-US/DouranPortal/1/page/Home>
- Hanh, H. Q., Azadi, H., Dogot, T., Ton, V. D. and Lebailly, P. (2017). Dynamics of agrarian systems and land use change in North Vietnam. *Land Degrad. Dev.*, 28: 799-810. <https://doi.org/10.1002/ldr.2609>
- Hudson, G. and Birnie, R. V. (2000). A method of land evaluation including year to year weather variability. *Agric. Forest Meteorol.*, 101: 203-216. [https://doi.org/10.1016/S0168-1923\(99\)00158-8](https://doi.org/10.1016/S0168-1923(99)00158-8)
- Igue, A. M., Maier, R., Gaiser T. and Stahr, K. (2005). Land evaluation of cotton production in the Azoka of catchment in the moist Savanna of Benin. *Conference Int. Agric. Res. Dev.*, 35(3): 59-64. <https://www.ars.usda.gov/research/publications/publication/?seqNo115=365130>
- Kalogirou, S. (2002). Expert systems and GIS: an application of land suitability evaluation. *Comp. Environ. Urban Syst.*, 26: 89-112. [https://doi.org/10.1016/S0198-9715\(01\)00031-X](https://doi.org/10.1016/S0198-9715(01)00031-X)
- Kilic, S., Evrendilek, F., Senol, S. and Celik, I. (2005). Developing a suitability index for land uses and agricultural land covers: a case study in Turkey. *Environ. Monitor. Assess.*, 102: 323-335. <https://doi.org/10.1007/s10661-005-6030-6>
- Lal, R. (2009). Soils and sustainable agriculture: A review. *Agron. Sustain. Dev.*, 28: 57-64. <https://doi.org/10.1051/agro:2007025>
- Magboul, M., Sulieman, M., Ibrahim, S. and Elfaki, J. (2015). Land suitability characterization

- for crop and fruit production of some river Nile Terraces, Khartoum North, Sudan. *Int. J. Sci. Res. Pub.*, 5(1). <http://hdl.handle.net/123456789/2070>
- Mahari, A. and Alebachew, A. (2013). Land suitability evaluation for irrigation valuation for irrigation in Dejen District, Ethiopia. *Int. J. Sci. Res. Pub.*, 3(9). <http://www.ijsrp.org/research-paper-0913/ijsrp-p2137.pdf>
- Mashayekhi, F. (2015). Evaluation of qualitative and quantitative land suitability for main crops in Khoda afarin region, Moghan plain. *M.Sc. thesis, Department of Soil Science, Shahed University*. <https://research.shahed.ac.ir/wsr/webpages/Report/ThesisView.aspx?ThesisID=4377>
- Menjiver, J. C., Aquilar, J., Garcia, I. and Bouza, P. (2003). Evaluation of olive orchard soils of map (Torres, Spain). *Int. sym. Sustain. use Manage. Soil Arid Semiarid.*, 984 pp. <http://www.edafologia.net/comun/congres/juanca.htm>
- Niekerk, A. V. (2010). A comparison of land unit delineation techniques for land evaluation in the Western Cape, South Africa. *Land Use Policy*, 27: 937-945. <https://doi.org/10.1016/j.landusepol.2009.12.007>
- Norton, B. J., Sandor, J. A. and White, C. S. (2003). Hillslope soil and organic matter dynamics within native American agroecosystem of the Colorado Plateau. *Soil Sci.*, 67: 225-234. <https://doi.org/10.2136/sssaj2003.2250>
- Panagos, P., Borrelli, P., Poesen, J., Ballabio, C., Lugato, E., Meusburger, K., Montanarella, L. and Alewell, C. (2015). The new assessment of soil loss by water erosion in Europe. *Environ. Sci. Policy.*, 54: 438-447. <https://doi.org/10.1016/j.envsci.2015.08.012>
- Rahimi lake, H., Taghizadeh, M., Aknarzadeh, R. and Ramezanpour, A. (2009). Qualitative and quantitative land suitability evaluation for olive production in Roodbar regon, Iran. *Medwell J.*, 4: 52-62.
- Rossiter, D. G. (1996). A theoretical framework for land evaluation. *Geoderma.*, 72: 165-190. [https://doi.org/10.1016/0016-7061\(96\)00031-6](https://doi.org/10.1016/0016-7061(96)00031-6)
- Sam, K., Coulon, F. and Prpich, G. (2016). Working towards an integrated land contamination management framework for Nigeria. *Sci. Total Environ.*, 571: 916-925. <https://doi.org/10.1016/j.scitotenv.2016.07.075>
- Santana-Cordero, A. M., Ariza, E. and Romagosa, F. (2016). Studying the historical evolution of ecosystem services to inform management policies for developed shorelines. *Environ. Sci. Policy.*, 64: 18-29. <https://doi.org/10.1016/j.envsci.2016.06.002>
- Shepande, C. (2002). Soil and Land Use with Particular Attention to Land Evaluation for Selected Land Use Types in the Lake Neivasha Basin, Kenya. *International Institute for aerospace survey and earth sciences (ITC), Enschede, the Netherlands*, 106 pp. <ftp://majisys.itc.utwente.nl/pub/naivasha/ITC/Shepande2002.pdf>
- Sys, C., Vanranst, E. and Debaveye, J. (1993). Land evaluation. Part II. Methods in land evaluation, International training center for post graduate soil scientist, *Ghent University, Ghent*. 247. <https://lib.ugent.be/catalog/pug01:233235>
- Wahba, M. M. and Darwish, K. M. (2007). Suitability of specific crops using microleis program in Sahal Barakas, Egypt. *J. Appl. Sci. Res.*, 3(7): 531-539. <http://www.aensiweb.com/old/jasr/jasr/2007/531-539.pdf>
- ZaliVargahan, B., Shahbazi, F. and Hajrasouli, M. (2001). Quantitative and qualitative land suitability evaluation for maize cultivation in Ghobadlou region, Iran. *Ozean J. Appl. Sci.*, 4(1): 91-104. http://issc.areeo.ac.ir/article_35157.pdf