



### Role of oxidative stress on acute ischaemic stroke

Ismail Setyopranoto\*<sup>1</sup>

<sup>1</sup>Neurology Department of Medical Faculty Gajah Mada University / KSM Saraf RSUP Dr Sardjito Yogyakarta

Review Article

#### ABSTRACT

##### ARTICLE INFO

###### **Keyword:**

Acute ischemic stroke,  
cell death,  
free radical,  
oxidative stress,  
antioxidant

###### **\*Corresponding author:**

ismail.setyopranoto@ugm.ac.id

DOI : 10.20885/JKKI.Vol7.Iss4.art6

After the onset of a stroke, blood flows disrupted in areas affected by vascular occlusion limit the delivery of oxygen and metabolic substrates to neurons causing ATP reduction and energy depletion. The glucose and oxygen deficit that occurs after severe vascular occlusion is the origin of the mechanisms that lead to cell death and cerebral injury caused of oxidative stress. Oxidative stress constitutes mechanism of injury of many types of disease processes. On oxidative stress occurs on the increase in ROS and RNS. This paper will discuss about cerebral ischemia that causes activation of ROS and RNS, also the mechanisms that play a role in cell death after cerebral ischemia, for example the role of phospholipase, Haber-Weiss reaction, and lipid peroxidation. It is also described about anti-oxidants to fight free radicals, for examples glutathione peroxidase, catalase and superoxide dismutase.

*Setelah onset stroke, terjadi gangguan aliran darah di daerah yang mengalami oklusi vaskuler, sehingga terjadi gangguan pengiriman oksigen dan metabolisme substrat pada neuron, hal ini menyebabkan pengurangan ATP dan energi. Terjadinya defisit glukosa dan oksigen yang terjadi setelah oklusi vaskular merupakan mekanisme utama yang menyebabkan kematian sel dan cedera otak oleh karena stres oksidatif. Pada oksidatif stres terjadi peningkatan ROS dan RNS. Tulisan ini membahas tentang iskemia serebral yang menyebabkan aktivasi ROS dan RNS, mekanisme yang berperan dalam kematian sel setelah iskemia serebral, misalnya peran fosfolipase, reaction Haber-Weiss, dan peroksidasi lipid, serta beberapa anti-oksidan untuk melawan radikal bebas, misalnya glutathione peroksidase, katalase dan superoksida dismutase.*

#### BACKGROUND

Cerebral ischaemia is a condition in which the blood flow to the brain decreased, either due to ischaemic stroke with acute focal or global neurological deficits.<sup>1</sup> Brain ischaemia increases the levels of ions Na<sup>+</sup> intracellular (Na<sup>i</sup>) and Cl<sup>-</sup> intracellular (Cl<sup>i</sup>), as well as K<sup>+</sup> extracellular (K<sup>e</sup>) causing neuronal depolarization, release of glutamat, and opening of Ca<sup>++</sup> channel.<sup>2,3,4</sup> The activation of glutamat receptors, especially N-methyl-D-aspartate (NMDA), would be followed by the entrance of Ca<sup>++</sup> into the cells (Calcium influx).<sup>5,6</sup> In

addition to that, an increase of intraneuronal Ca<sup>++</sup> concentration would occur due to failure of active Ca<sup>++</sup> elimination to extracellular through Sodium pump, as well as the release of Ca<sup>++</sup> from mitochondria and endoplasmic reticulum to cytoplasma. The increase of Ca<sup>++</sup> would generate the activation of enzymes phospholipase A2, nitrate oxide sintase (NOS), guanilat cyclase, calsineurin, endonuclease, protease and protein kinase.<sup>7,8,9</sup> The increase of Ca<sup>++</sup> would activate phospholipase A2 which hydrolyzes phospholipid membrane that consist of linoleic acid and arachidonic acid. Hydrolyzed arachidonic acid

would form leucotriene, prostaglandin, and free radical anion superoxide ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ) and hydroxyl ( $OH^-$ ), by the assistance of lipoxygenase and cyclooxygenase ( $COX_1$  dan  $COX_2$ ).<sup>10,11,12</sup>

Free radicals anion superoxide ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ) and hydroxyl ( $OH^-$ ) are the main pathway of the formation of Reactive Oxygen Species/ ROS through the reduction of molecular oxygen into  $H_2O$ . The presence of transitional minerals like  $Fe^{2+}$ , would cause hydrogen peroxide ( $H_2O_2$ ) to go through fenton reaction, and increase the concentration of radical hydroxyl ( $OH^-$ ). This radical hydroxyl is highly reactive and is a major cause of oxidative stress. As a defense mechanism against free radicals, there are anti-oxidant enzymes like yaitu glutathione peroxidase (GPX), catalase (CAT) and superoxide dismutase (SOD). Superoxide dismutase transforms  $O_2^-$  into  $H_2O_2$ , and afterward glutathione peroxidase and catalase transforms  $H_2O_2$  into  $H_2O$ . These enzymes are able to bind antioxidant like ascorbic acid, and  $\alpha$ -tocopherol, resulting in decreased ROS concentration. During this process either an increase of free radicals concentration or the decrease of antioxidant concentration would cause oxidative stress and evoke damage to lipid, protein, or DNA, resulting into cell death through either apoptosis or necrosis.<sup>12</sup>

Oxidative stress as a result of increase ROS concentration,<sup>13</sup> might occur due to oxygen metabolism, oxygen reperfusion injury from hypoxic state, as well as hemoglobin and myoglobin oxidation.<sup>14</sup> Normally, ROS plays many roles in physiological processes, such as : the body's defense system, hormones biosynthetic, fertilization, and cellular signaling.<sup>13</sup> ROS also take part in the immune system by combating antigen during phagocytosis.<sup>14</sup> Nevertheless, the increase of ROS production have implications in many diseases, for instance hypertension, atherosclerosis, diabetes, heart failure, and stroke.<sup>13</sup>

## ISSUES PROJECTION

A lot of ischaemic stroke patients were seen to still become worsen even with appropriate care in the Stroke Unit of RSUP Dr. Sardjito Yogyakarta.

Even after standard management had been provided, risk factors had been controlled, and no complications were found.

Many research had reported the occurrence of complications post-stroke. There were even one research that reported 96% of all stroke patients who were hospitalized had more than one medical or neurological complications.<sup>15,16,17</sup> A thorough study is needed to determine its pathophysiology, therefore the possibility of oxidative stress as the cause of worsening in acute ischaemic stroke patients need to be examined.

Published research about oxidative stress and worsening of ischaemic stroke patients are still controversial due to the complexity of its mechanism. A research by Tsai et al.,<sup>18</sup> on 120 non-cardioembolic acute ischaemic stroke patients at Chang Gung Memorial Hospital's Neurological ward in Kaohsiung Taiwan, concluded that oxidative stress contribute to progressive neurological damage on certain cases of acute ischaemic stroke patients, which were shown on bad NIHSS score and stroke subtype.

Another research by İçme et al.,<sup>19</sup> on 68 subjects, which consist of 34 ischaemic stroke subjects and 34 non-stroke control subjects, at Atatürk Training and Research Hospital Ankara Turki from January 2012 until January 2013, concluded that the increase of oxidants in oxidative stress contribute on the pathogenesis of ischaemic stroke, but there were no significant correlation between Total Anti-oxidant Status (TAS), Total Oxidant Status (TOS), Oxidative Stress Index (OSI), infarction volume, NIHSS score, and oxidative stress with the severity of stroke.

İçme et al.,<sup>20</sup> also reported based on their research on 92 subjects which consist of 74 acute ischaemic stroke subjects and 18 hemorrhagic stroke subjects at Atatürk Training and Research Hospital Ankara Turki from January 2013 until November 2013, oxidative stress contribute in the pathogenesis of both ischaemic and hemorrhagic stroke, however oxidative stress had uncertain correlation with the severity of both types of stroke.

Based on these data, the mechanism of

ischaemic stroke which might contribute to the worsening of patients condition needs to be studied. The objective of this study is to determine the role of oxidative stress on acute ischaemic stroke.

**REACTIVE OXYGEN SPESIES**

Free radicals consist of several species that contain one or more unconjugated electron. Free radicals are reactive chemical spesies, such as reactive oxygen spesies and reactive nitrogen spesies. ROS consist of anion superoxide ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ) and hydroxil ( $OH^-$ ), while RNS consist of radical nitrit oxide ( $NO^-$ ) and peroxyxynitrate ( $ONOO^-$ ).<sup>12,21,22</sup>

Figure 1 showed the role of phospholipase in the formation of ROS and lipid peroxide on brain ischaemia.

Excess  $Ca^{++}$  intra cells,  $Ca^{++i}$  could be eliminated by the activation of Calcium adenosine

triphosphatase plasma membrane .

In ischaemic state where the concentration of ATP (adenosine triphosphate) is very low, this mechanism would become inactive resulting in uncontrolled increase of  $Ca^{++}$  concentration intra cells. Increase  $Ca^{++}$  inside the cells would activate various enzymes, such as nitrate oxide sintase (NOS), phospholipase A2, and phospholipase C, protease and endonuclease.

NOS activation would result in the formation of nitrNOS activation would result in the formation of nitrit oxide, while the activation of phospholipase A2 would result in the hydrolization of phospholipid membrane which then disintegrate arachidonic acid into prostaglandin, prostacycline, and tromboxane A2, as well as the formation of free radicals superoxide, hidrogen peroxide, and hydroxil as ROS.<sup>9,23,24</sup>





Figure 1. The role of phospholipase in the formation of reactive oxygen spesies

During ischaemia, ATP would be decreased causing the increase of AMP concentration and subsequently supply hypoxanthine substrates into xanthine oxidase.<sup>25</sup> Afterward, reperfusion would also supply oxygens which would act as a secondary substrate that is needed by xanthine oxidase to produce free radicals.<sup>26,27</sup> Hydrogen peroxide could also be formed from other mechanisms which is oxidoreductase in cellular peroxisome organelles.<sup>25,28</sup> Hydrogen peroxide is a free radical that does not have electron pairs. Whilst not reactive, hydrogen peroxide could easily diffuse through biological membranes,<sup>29,30</sup> and the presence of ferro ( $\text{Fe}^{2+}$ ) or cupro ( $\text{Cu}^+$ ), would cause hydrogen peroxide to transform into radical hydroxyl (reaction 1). Oxidized iron ( $\text{Fe}^{3+}$ ) would be reduced by superoxide anion (reaction 2). Reaction 1 and 2 are called Haber-Weiss reaction.<sup>22,31</sup>

Hydroxyl is a very strong oxidant of the biological system. Hydroxyl would take one electron from the thiol of an enzyme, for instance glutathione reductase and glutathione peroxidase, DNA and polyunsaturated lipid, or other hydroxylated molecules. Hydroxyl also could alter the function of mitochondria, inactivate electron-carrying-protein and mitochondrial ATPase, as well as peroxidized lipid membrane.<sup>25,32</sup>

### Haber-Weiss Reaction



Single oxygens are formed from the interaction of superoxide anion with another superoxide anion, peroxynitrate with hydrogen peroxide, or superoxide anion with hydrogen peroxide. Single oxygens are able to inactivate Calcium ATPase, denature protein, inactivate superoxide dismutase and catalase, and oxidize polyunsaturated lipid into lipid hydroperoxide and endoperoxide. With the help of ions, these peroxides could trigger the development of lipid peroxidized chains or increase the production of single oxygens.<sup>25,33</sup>

### REACTIVE NITROGEN SPECIES

Nitrite oxide as well as its enzyme, which is

nitrite oxide synthase (NOS), contribute in brain ischaemia. Normally, nitrite oxide is a benign molecular signal. Nitrite oxide is synthesized from arginine by NOS, and it also has three isoforms which are endothelial NOS (eNOS) that is produced by endothelial cells, inducible NOS (iNOS) that is produced by macrophages, and neuronal NOS (nNOS) that is produced by neurons.<sup>8,34</sup> Neuronal NOS (nNOS) and endothelial NOS (eNOS) are highly dependent on Calcium and Calmodulin. Inducible NOS (iNOS) is produced by cytokines activation through second messenger pathway and does not require the increase of cytosolic Calcium.<sup>35</sup> Inducible NOS produce nitrite oxide that is toxic in brain ischaemia.<sup>8,36</sup> Nitrite oxide is able to interact with anion superoxide to produce peroxynitrate which will transform into nitrogen dioxide (an oxide reaction similar to radical hydroxyl), oxidized protein that contain thiol, DNA base and polyunsaturated lipid, and inhibit mitochondrial electron transport hence causing a decrease of ATP production and an increase of ROS production.<sup>32,37</sup> In brain injury, polymorphonuclear leukocyte could produce nitrite oxide and superoxide. Neurons and oligodendrocytes are highly vulnerable to the effect of nitrite oxide and peroxynitrate.<sup>31,38</sup>

Nitrite oxide has high diffusion velocity so that it can bind with hemoglobin. In the presence of free radical superoxide, nitrite oxide molecules would bind with oxygen and form peroxynitrate which is cytotoxic to neurons.<sup>4,8,36</sup> The cytotoxic effect of nitrite oxide occurs in two ways: (1) nitrite oxide would decrease the amount of growth factor, and (2) increase the concentration of enzymes that disintegrate protein, both resulting in DNA damage. Nitrite oxide that is mediated by the reaction with superoxide would produce peroxynitrate, a neurotoxic free radical. Peroxynitrate would trigger protein nitrosylation, which leads to DNA damage and induce brain cell death, as well as altering brain function.<sup>11</sup>


### LIPID PEROXIDATION

When brain contains a lot of polyunsaturated fatty acids, ROS could easily increase lipid peroxidation.<sup>39</sup> Anion superoxide would cause deesterification of phospholipid membrane to release fatty acids. Radical hydroxyl extract is a

hydrogen atom that originated from methylene carbon of unsaturated fatty acids and produce radical lipid carbon-centred ( $L^{\cdot}$ ; reaction 3). Radical lipid would react with oxygen molecules to form peroxy radicals (reaction 4), which transform into lipid hydroperoxide ( $LOO^{\cdot}$ ) by separating hydrogen atom from methylene carbon that contiguous with unsaturated fatty acids (reaction 5).<sup>29,32,40,41</sup>



Lipid peroxidation could also be triggered by non-radical ROS, for instance single oxygen and hypochloride. As the presence of iron or iron complex, lipid hydroperoxide could form alkoxy radicals (reaction 6) or peroxy (reaction 7) and then generate new peroxidation chains (reaction 8 and 9), especially in acidic condition, for instance during ischaemia.<sup>22,31,42,43,44</sup>



Lipid peroxidation could alter membrane instability and permeability, as well as the function of ion pumps on the membrane border. This condition would endangered ion homeostasis, and could result in the loss of membrane integrity and cell damage.<sup>22,31,45</sup> Lipid peroxidation could also trigger disintegration of pro-inflammatory isoprostanoic mediator which is highly dangerous and release strong oxidant like 4-hydroxynonenal (4-HNE).<sup>31,46</sup>

### ACUTE ISCHAEMIC STROKE AND OXIDATIVE STRESS

Ischaemia would be followed by reperfusion that can generate ROS and NOS that are toxic to brain tissues. Reperfusion cause an increase of oxidative stress due to the restoration of oxygen into brain tissues.<sup>22,42,47</sup> Oxidative stress contributes in the pathophysiology of ischaemic stroke, and brain is the organ most sensitive to oxidative stress. The presence of free radical

activities would cause the brain to consume more lipid, increase oxygen demand, and trigger the oxidation of dopamine and glutamate. Furthermore, the concentration of catalase enzyme in the neurons would decrease, causing deterioration of glutathion peroxidase ability to eliminate hydrogen peroxide. ROS is a dangerous end-product of oxidative phosphorylation and reperfusion injury, because it could cause cellular redox which will influence the activity of protein-protein bond and DNA-protein bond of certain enzymes and transcription factors.<sup>48</sup>

Main pathway of the formation of ROS is through molecular oxygen reduction into  $H_2O$ . This reduction would generate dioxygen molecules from superoxide anion ( $O_2^{\cdot-}$ ) and hydrogen peroxide ( $H_2O_2$ ). The presence of  $Fe^{2+}$  as transitional metal might cause hydrogen peroxide to undergo fenton reaction, which will increase the concentration of radical hydroxyl ( $OH^{\cdot}$ ). Radical hydroxyl is highly reactive and is a major cause of oxidative damages. There are many instances of antioxidant enzymes, such as glutathion peroxidase (GPX), catalase (CAT) and superoxide dismutase (SOD).<sup>41,49</sup> Superoxide dismutase would change  $O_2^{\cdot-}$  into  $H_2O_2$ , and glutathion peroxidase and catalase would change  $H_2O_2$  into  $H_2O$ . These enzymes would coupled with antioxidant like ascorbic acid and  $\alpha$ -tocopherol, resulting in the decrease of ROS concentration.

The increase of free radicals concentration or decrease of antioxidant would cause oxidative stress, resulting in lipid, protein, or DNA damage, which all will lead to cell death.<sup>12</sup>

### NEUROPROTECTIVE DRUG THAT TARGETS OXIDATIVE STRESS

The objective of therapy in stroke patients is to increase clinical outcome by minimalizing the amount of damaged brain cells, treating complications, and accelerating the recovery of neurological function.<sup>50</sup> Drugs that are able to prevent or decrease the effect of oxidative stress and minimalize the amount of cell damage is called neuroprotective drugs, such as : antioxidant, agonis GABA, as well as antagonis AMPA.

Until now, there are more than 100

neuroprotective drugs had been studied and showed positive impact on ischaemic stroke in preclinical stage, however none has been clinically proven. Nevertheless, the result of these researches would be able to enhance our understanding on the biological basic of ischaemic brain injury and could become a strong keystone to enhance the management of stroke in the future.<sup>51</sup> Below are the description of these neuroprotective drugs

### **1. Ebselen**

Ebselen showed antioxidant effect by inhibiting lipid peroxidation and the expression of inducible nitric oxide synthase (iNOS) in the cerebral cortex of mice that suffered from stroke due to spontan hypertension.<sup>52</sup>

### **2. Tirilazad**

Tirilazad is a non-glucocorticoid drugs that is composed by 21-aminosteroid which prevent lipid peroxidation. A systematic review on tirilazad showed efficacy on animal models who suffered from focal ischaemia, which was shown by a decrease of infarction volume and increase of neurobehavior score.<sup>53</sup> None the less, tirilazad was not significant in regards of infarction volume in patient with acute ischaemic stroke.<sup>54</sup>

### **3. Edaravone**

Edaravone is an anti free radicals which work by preventing endothelial injury that would cause neuron damage in brain ischaemia. Edaravone will increase eNOS (NOS that is able to improve endothelial function, which would be beneficial in ischaemic stroke), as well as decreasing nNOS and iNOS (non-beneficial NOS).<sup>55</sup> Antioxidant edaravone significantly reduce infarction volume and increase neurological deficits score in mice,<sup>56</sup> reduce cell death due to oxidative in mice,<sup>57</sup> repair the size of lesion in ischaemic stroke and neurological deficits in patients with small vascular occlusion, however there had not been any significant difference on clinical outcome.<sup>58</sup>

### **4. Clomethiazole**

Clomethiazole work as neuroprotectant by being GABA agonis, in ischaemia model is well tolerated.<sup>59</sup> Nevertheless, it was reported to not have significant effects on sequele or the volume of infarction in ischaemic stroke patients.<sup>60</sup>

### **5. Aptiganel (CNS-1102)**

Aptiganel is a neuroprotectant that works as non-competitive NMDA antagonist which is effective in focal ischaemia in rat models.<sup>61</sup> Phase II/III of the research was stopped due to lack of efficacy and mortality risk.<sup>62</sup>

### **6. YM872**

YM872 is a neuroprotectant agent that works as AMPA antagonist, which significantly reduce infarct volume and improve neurological deficits in rat models with embolic stroke.<sup>63</sup> Results of phase II research have not been published.<sup>64</sup>

### **7. Magnesium Sulfat**

Ion Magnesium could affect various enzymes that contribute in a lot of celullar function, such as : cell membrane permeability, mitochondrial function, and ion membrane for cell conduction, which in hypoxia would prevent cell death.<sup>65</sup> Magnesium showed neuroprotective effects on animal models,<sup>66</sup> as well as phase II research in stroke patients.<sup>67</sup> Namun demikian Magnesium failed to show significant effect on phase III research.<sup>68</sup>

### **8. Dapsone**

Dapsone as neuroprotectant works by inhibiting glutamat exitotoxicity and inflammatory respond after brain ischaemia.<sup>69</sup> Dapsone was declared safe and effective on animal models,<sup>70</sup> and research on stroke patients had been done.<sup>69</sup> Phase III research had also been done but the results have not been published.<sup>71</sup>

### **9. Cromolyn**

Cromolyn works by inhibitting cell mast migration and degranulation, activates glial cells and prevent neural cell death. Cromolyn is also effective in reducing brain oedem and repairing blood-brain barrier permeability in animal model with stroke.<sup>72</sup> Currently clinical research phase III on acute ischaemic stroke patient is ongoing.<sup>73</sup>

According to Cheng et al.,<sup>74</sup> there are a lot of reasons that affect the success of neuroprotectant medicines in phase I and II research, none the less, it failed on clinical research, such as : (1) preclinical research use very short window periods in drug administration, while clinical research has longer window periods, (2) the target of preclinical research is ischaemic penumbra, while clinical research

targets its outcome, (3) optimal duration for neuroprotective drug administration is unknown, (4) preclinical research uses different outcome measuring, for instance the size of infarct to determine the success of therapy, while clinical research uses its clinical outcome, like Rakin scale or Barthel index modification, (5) preclinical research depend on early outcome, while clinical research depend on its end outcome, (6) stroke pathophysiological variation, most preclinical research uses the occlusion of medial cerebral artery as ischaemic stroke model, hence there are no heterogeneity of stroke pathophysiology that would affect the duration and severity of ischaemic stroke. In contrast, research in human has a wide range of heterogeneity of stroke pathophysiology, and (7) comorbidity differences, most experimental model uses young healthy rats that was not exposed to other medicines, while real stroke patients usually has various severe secondary diseases (history of ischaemic attack, cardiovascular diseases, etc) and history of other medicine usage.

## CONCLUSION

Cerebral ischaemic is the most common cause of acute ischaemic stroke. The pathophysiology of ischaemic stroke is very complex and involve various mechanism including the formation of free radicals. Imbalance of cellular production in the form of free radicals and the ability of cells to combat against it is called oxidative stress. Eventhough the mechanism is unclear, oxidative stress is one of, if not the most, important event in ischaemic stroke and contributes to the worsening of stroke event.

There are already a lot of medicines to combat against oxidative stress that are known as neuroprotective agent which have been studied and showed positive effect on ischaemic stroke in preclinical research, however none of them have shown significant clinical effect. Nevertheless, the results of these research could improve our understanding about the mechanism of ischaemic brain injury and become a strong basis to develop better stroke management in the future.

There are many potential medicines that might be able to prevent or reduce the effect of

oxidative stroke, hence minimizing the amount of cell damage. These medicines are called neuroprotective, for instance : antioxidant, GABA agonist, as well as AMPA antagonist.

## REFERENCES

1. Bandera E, Botteri M, Minelli C, Sutton A, Abrams KR, Latronico N. Cerebral Blood Flow Threshold of Ischemic Penumbra and Infarct Core in Acute Ischemic Stroke. A Systematic Review. *Stroke* 2006;37:1334-9.
2. Meldrum BS. Glutamate as a neurotransmitter in the brain: Review of physiology and pathology. *J. Nutr.* 2000;130:1007S-1015S.
3. Danysz W, Parsons CG. The NMDA receptor antagonist memantine as a symptomatic and neuroprotective treatment for Alzheimer's disease: preclinical evidence. *Int J Geriatr Psychiatry* 2002;18:S23-S32.
4. Schild L, Reiser G. Oxidative stress is involved in the permeabilization of the inner membrane of brain mitochondria exposed to hypoxia/reoxygenation and low micromolar  $Ca^{2+}$ . *FEBS Journal* 2005;272:3593-601.
5. Barone FC, Tuma RF, Legos JL, Erhardt JA, Parsons AA. Brain Inflammation, Cytokines, and p38 MAP Kinase Signaling in stroke, in Lin, RCS, *New Concepts in Cerebral Ischemia*. CRC Press 2002; p:201-44.
6. Kemp JA, McKernan RM. NMDA receptor pathways as drug targets. *Nature Neuroscience Suppl.* 2002; 5:1039-42.
7. Gwag BJ, Won SJ, Kim DY. Excitotoxicity, oxidative stress, and apoptosis in ischemic neuronal death in Cohen SN, *Management of Ischemic Stroke*. Mc Graw Hill, Health Professions Division, 2002.
8. Walford G, Loscalzo J. Nitric Oxide in vascular biology. *Journal of Thrombosis and Haemostasis* 2003;1:2112-8.
9. Berridge MJ, Bootman MD, Roderick HL. Calcium signaling: Dynamics, homeostasis and remodelling. *Molecular cell biology* 2003;4:517-29.
10. Kramer BC, Yabut JA, Cheong J, Jnobaptiste R, Robakis T, Olanow CW, dkk. Toxicity of Glutathione depletion in mesencephalic cultures: A role for Arachidonic acid and its

- lipoxygenase metabolites. *Eur J Neurosci* 2004;19:280-286.
11. Taylor JM, Crack PJ. Impact of oxidative stress on neuronal survival. *Clin and Experiment Pharmacol and Physiol*. 2004;31:397-406.
  12. Alexandrova ML, Bochev PG. Oxidative stress during the chronic phase after stroke. *Free Radic Biol Med* 2005;39:297-16.
  13. Paravicini TM, Touyz RM. NADPH Oxidase, Reactive Oxygen Species, and Hypertension. *Journal Diabetes Care* 2008;31(2):S170-S180.
  14. Finaud J, Lac G, Filaire E. Oxidative Stress, Relationship with Exercise and Training. *Journal Sports Med*. 2006;36(4):327-58.
  15. Hong KS, Kang DW, Koo JS, Yu KH, Han MK, Cho YJ, et al. Impact of neurological and medical complications on 3-month outcomes in acute ischaemic stroke. *Eur J Neurol*. 2008;15:1324-31.
  16. Indredavik B, Rohweder G, Naalsund E, Lydersen S. Medical complications in a comprehensive stroke unit and an early supported discharge service. *Stroke*, 2008;39:414-20.
  17. Sorbello D, Dewey HM, Churilov L, Thrift AG, Collier JM, Donnan G, et al. Very early mobilization and complications in the first 3 months after stroke: further results from phase II of A Very Early Rehabilitation Trial (AVERT). *Cerebrovasc Dis*. 2009;28:378-83.
  18. Tsai N-W, Chang Y-T, Huang C-R, Lin Y-J, Lin W-C, Cheng B-C, et al. Association between Oxidative Stress and Outcome in Different Subtypes of Acute Ischemic Stroke. *BioMed Research International* 2014, Article ID 256879, 7 pages. <http://dx.doi.org/10.1155/2014/256879>
  19. İçme F, Erel Ö, Avci A, Satar S, Gülen M, Acehan S. The relation between oxidative stress parameters, ischemic stroke, and hemorrhagic stroke. *Turk J Med Sci*. 2015;45:947-53.
  20. İçme F, Erel Ö, Öztürk ZS, Öz T, Avci A, Satar S, et al. The relationship of oxidative stress parameters with infarct volume and National Institutes of Health Stroke Scale in ischemic stroke. *Turk J Biochem* 2015;40(4):275-281.
  21. Sampayo JN, Gill MS, Lithgow GJ. Oxidative stress and aging – the use of dismutase superoksida/catalase mimetics to extend lifespan. *Biochemical Society Transactions* 2003;31(6):1305-1307.
  22. Madamanchi NR, Hakim ZS, Runge MS. Oxidative stress in atherogenesis and arterial thrombosis: the disconnect between cellular studies and clinical outcomes. *J Thromb Haemost*. 2005;3:254-267.
  23. Lynch DR, Guttman RP. Excitotoxicity: Perspectives based on N-Methyl-D-Aspartate receptor subtypes. *J Pharmacol Exp Ther* 2002;300:717-723.
  24. Madamanchi NR, Vendrov A, Runge MS. Oxidative Stress and Vascular Disease. *Arterioscler Thromb Vasc Biol*. 2004;25:29-38.
  25. Adibhatla RM, Hatcher JF. Phospholipase A2, reactive oxygen species, and lipid peroxidation in cerebral ischemia. *Free Radic Biol Med* 2006;40:376-387.
  26. Martelin E, Lapatto R, Raivio KO. Regulation of xanthine oxidoreductase by intracellular iron. *Am J Physiol Cell Physiol*. 2002;283:1722-28.
  27. McNally JS, Saxena A, Cai H, Dikalov S, Harrison DG. Regulation of Xanthine Oxidoreductase Protein Expression by Hydrogen Peroxide and Calcium. *Arterioscler. Thromb. Vasc. Biol*. 2005;25:1623-28.
  28. Vincent AM, Russell JW, Low P, Feldman EL. Oxidative Stress in the pathogenesis of Diabetic Neuropathy. *Endocrine Reviews* 2004;25(4):612-28.
  29. Lee J, Koo N, Min DB. Reactive Oxygen Species, Aging, and Antioxidative Nutraceu-ticals. *Compr Rev Food Sci F* 2004;3:21-33.
  30. Nohl H, Gille L, Staniek K. The mystery of reactive oxygen species derived from cell respiration. *Acta Biochim Pol* 2004;51(1):223-29.
  31. Stadtman ER. Role of Oxidant Species in Aging. *Curr. Med. Chem*. 2004;11:1105-12.
  32. Andreyev AY, Kushnareva YE, Starkov AA. Mitochondrial Metabolism of Reactive Oxygen Species. *Biochemistry* 2005;70(2):200-14.
  33. Li C, Jackson RM. Reactive Species Mech-

- anisms of Cellular Hypoxia Reoxygenation Injury. *Am J Physiol Cell Physiol.* 2002;282:C227-C41.
34. Margail I, Plotkine M, Lerouet D. Antioxidant strategies in the treatment of stroke. *Free Radic Biol Med* 2005;39:429-43.
  35. Juurlink BHJ. Management of oxidative stress in the CNS: the many roles of glutathione. *Neurotox. Res.* 1999;1:119-40.
  36. Chan KL. Role of Nitric Oxide in Ischemia and Reperfusion Injury. *Curr. Med. Chem. – Anti-Inflamm Anti-Allergy Agents* 2002;1:1-13.
  37. Drew B, Leeuwenburgh C. Aging and The Role of Reactive Nitrogen Species. *Ann. N.Y. Acad. Sci.* 2002;959:66-81.
  38. Turrens JF. Mitochondrial formation of reactive oxygen species. *J. Physiol.* 2003;552:335-344.
  39. Ishibashi N, Prokopenko N, Lefkowitz MW, Reuhl KR, Mirochnitchenko O. Glutathione peroxidase inhibits cell death and glial activation following experimental stroke. *Brain Res. Mol. Brain Res.* 2002;109:34-44.
  40. Tarpey MM, Fridovich I. Methods of Detection of Vascular Reactive Species: Nitric Oxide, Superoxide, Hydrogen Peroxide, and Peroxynitrite. *Circ. Res.* 2001;89:224-36.
  41. Warner DS, Sheng H, Batinic-Haberle I. Review - Oxidants, antioxidants and the ischemic brain. *The J.Exp.Biol.* 2004;207:3221-31.
  42. Wall J. Antioxidants in prevention of reperfusion damage of vascular endothelium. *TSMJ* 2000;1:67-71.
  43. Selim MH, Ratan RR. The role of iron neurotoxicity in ischemic stroke. *Ageing Res Rev* 2004;3:345-53.
  44. Stocker R, Keaney JF. Role of oxidative modifications in atherosclerosis. *Physrev.* 2004;84:1381-78.
  45. Fariss MW, Chan CB, Patel M, van Houten B, Orrenius S. Role of Mitochondria in Toxic Oxidative Stress. *Mol. Interv.*, 2005;5(2):94-111.
  46. Musiek ES, Gao L, Milne GL, Han W, Everhart MB, Wang D, et al. Cyclopentenone Isoprostanes Inhibit the Inflammatory Response in Macrophages. *The Jour of Biologic Chemis.* 2005;280(42):35562-35570.
  47. Hua SW, Yi ZC, Yi ZG. Antioxidants attenuate reperfusion injury after global brain ischemia through inhibiting nuclear factor-kappa B activity in rats. *Acta Pharmacol Sin.* 2003;24(11):1125-1130.
  48. Paravicini TM, Sobey CG. Cerebral vascular effects of Reactive Oxygen Species: Recent evidence for a role of NADPH – Oxidase. *Clin and Experiment Pharmacology and Physio.* 2003;30:855-859.
  49. Guo Y, Li P, Guo Q, Shang K, Yan D, Du S, dkk. Pathophysiology and Biomarkers in Acute Ischemic Stroke - A Review. *Tropical Journal of Pharmaceutical Research*, December 2013;12(6):1097-1105.
  50. Davis S, Lees K, Donnan G. Treating the acute stroke patient as an emergency: current practices and future opportunities. *Int J Clin Pract* 2006;60(4):399-407.
  51. Durukan A, Tatlisumak T. Acute ischemic stroke: overview of major experimental rodent models, pathophysiology, and therapy of focal cerebral ischemia. *Pharmacol Biochem Behav.* 2007;87:179-97.
  52. Sui H, Wang W, Wang PH, Liu LS. Protective effect of antioxidant ebselen (PZ51) on the cerebral cortex of stroke-prone spontaneously hypertensive rats. *Hypertens Res.* 2005;28:249-54.
  53. Sena E, Wheble P, Sandercock P, Macleod M. Systematic review and meta-analysis of the efficacy of tirilazad in experimental stroke. *Stroke*, 2007;38:388-394.
  54. van der Worp HB, Sena ES, Donnan GA, Howells DW, Macleod MR. Hypothermia in animal models of acute ischaemic stroke: a systematic review and meta-analysis. *Brain* 2007;130:3063-3074.
  55. Yoshida H, Yanai H, Namiki Y, Fukatsu-Sasaki K, Furutani N, Tada N. Neuroprotective effects of edaravone: a novel free radical scavenger in cerebrovascular injury. *CNS Drug Rev.* 2006;12(1):9-20.
  56. Zhang N, Komine-Kobayashi M, Tanaka R, Liu M, Mizuno Y, Urabe T. Edaravone reduces early accumulation of oxidative products and sequential inflammatory responses after transient focal ischemia in

- mice brain. *Stroke* 2005;36:2220-2225.
57. Lee BJ, Egi Y, van Leyen K, Lo EH, Arai K. Edaravone, a free radical scavenger, protects components of the neurovascular unit against oxidative stress in vitro. *Brain Res.* 2010;1307:22-7.
  58. Nakase T, Yoshioka S, Suzuki A. Free radical scavenger, edaravone, reduces the lesion size of lacunar infarction in human brain ischemic stroke. *BMC Neurol.* 2011;11:39.
  59. Chaulk D, Wells J, Evans S, Jackson D, Corbett D. Long-term effects of clomethiazole in a model of global ischemia. *Exp Neurol.* 2003;182:476-82.
  60. Lyden P, Shuaib A, Ng K, Levin K, Atkinson RP, Rajput A, et al. Clomethiazole Acute Stroke Study in Ischemic Stroke (CLASS-I): final results. *Stroke* 2002;33:122-8.
  61. Schäbitz WR, Li F, Fisher M. The N-methyl-D-aspartate antagonist CNS 1102 protects cerebral gray and white matter from ischemic injury following temporary focal ischemia in rats. *Stroke* 2000;31:1709-14.
  62. Albers GW, Goldstein LB, Hall D, Lesko LM, Aptiganel Acute Stroke Investigators. Aptiganel hydrochloride in acute ischemic stroke: a randomized controlled trial. *JAMA* 2001;286:2673-82.
  63. Suzuki M, Sasamata M, Miyata K. Neuroprotective effects of YM872 coadministered with t-PA in a rat embolic stroke model. *Brain Res.* 2003;959:169-72.
  64. A Study to Evaluate the Effects of YM872 on Stroke Lesion Volume in Acute Stroke Patients. 2002. <http://clinicaltrials.gov/ct2/show/NCT00044070>
  65. Singh H, Jalodia S, Gupta MS, Talapatra P, Gupta V, Singh I. Role of magnesium sulfate in neuroprotection in acute ischemic stroke. *Ann Indian Acad Neurol.* 2012;15(3):177-80.
  66. Marinov MB, Harbaugh KS, Hoopes PJ, Pikus HJ, Harbaugh RE. Neuroprotective effects of preischemia intraarterial magnesium sulfate in reversible focal cerebral ischemia. *J Neurosurg.* 1996;85:117-24.
  67. Bradford A, Lees K. Design of the Intravenous Magnesium Efficacy in Acute Stroke (IMAGES) trial. *Curr Control Trials Cardiovasc Med* 2000;1:184-90.
  68. Muir KW, Lees KR, Ford I, Davis S, Intravenous Magnesium Efficacy in Stroke (IMAGES) Study Investigators. Magnesium for acute stroke (Intravenous Magnesium Efficacy in Stroke trial): randomised controlled trial. *Lancet* 2004;363:439-45.
  69. Nader-Kawachi J, Góngora-Rivera F, Santos-Zambrano J, Calzada P, Ríos C. Neuroprotective effect of dapsone in patients with acute ischemic stroke: a pilot study. *Neurol Res.* 2007;29:331-334.
  70. Ríos C, Nader-Kawachi J, Rodríguez-Payán AJ, Nava-Ruiz C. Neuroprotective effect of dapsone in an occlusive model of focal ischemia in rats. *Brain Res.* 2004;999:212-5.
  71. Dapsone for Acute Ischemia Stroke Study (DAISY). 2010. <http://clinicaltrials.gov/show/NCT01144650>.
  72. Strbian D, Karjalainen-Lindsberg ML, Tatlisumak T, Lindsberg PJ. Cerebral mast cells regulate early ischemic brain swelling and neutrophil accumulation. *J Cereb Blood Flow Metab.* 2006;26:605-12.
  73. Treatment of Acute Stroke with Cromolyn (Single Dose). 2010. <http://clinicaltrials.gov/ct2/show/NCT01175525>.
  74. Cheng YD, Al-Khoury L, Zivin JA. Neuroprotection for ischemic stroke: two decades of success and failure. *NeuroRx.* 2004;1:36-45.