

REACTIONS OF $cis-[PtCl(^{15}NH_3)_2(H_2O)]^+$, THE FIRST CISPLATIN HYDROLYTES WITH THIOLS. 3: REACTIONS WITH GLUTATHIONE

Sutopo Hadi

Jurusan Kimia, FMIPA, Universitas Lampung
Jl. S. Brojonegoro No 1 Gedong Meneng, Bandar Lampung 35145
E-mail: sutopo_hadi@yahoo.com.au

ABSTRAK

Interaksi antara senyawa kompleks $cis-[PtCl(^{15}NH_3)_2(H_2O)]^+$ (2) dengan glutation (GSH) telah dilakukan pada larutan berair. Pada reaksi ini, padatan yang berbentuk polimer tidak terbentuk dan pelepasan ligan amina lebih lambat dibanding reaksi sejenis untuk kompleks $cis-[Pt(^{15}NH_3)_2(H_2O)_2]^{2+}$ (3). Dalam seri penelitian ini juga diamati bahwa walaupun reaksi yang terjadi antara kompleks 2 dengan berbagai tiol memberikan hasil yang akhir yang berbeda, akan tetapi pada tahap awal reaksi, senyawa yang terbentuk adalah sama yaitu terbentuknya spesi platina(II) dengan ligan klorida dan tiolat yang terikat secara monodentat melalui atom sulfur. Pada reaksi ini, setelah terbentuk hasil awal yaitu kompleks platina dengan glutation terikat secara monodentat $cis-[PtCl(^{15}NH_3)_2(SG)]$ (7), tidak ada senyawa kompleks lain yang teramati. Seperti pada reaksi kompleks 3 dengan GSH, senyawa bistiolat $cis-[Pt(NH_3)_2(SG)_2]$ (8) selalu teramati dalam reaksi ini. Sedangkan, kompleks sulfur dua inti berjembatan yang memberikan serapan lebar pada ^{15}N NMR hanya teramati dalam jumlah yang sangat sedikit.

Kata kunci: ^{15}N NMR, glutation, hidrolisat cisplatin

ABSTRACT

The interactions of $cis-[PtCl(^{15}NH_3)_2(H_2O)]^+$ (2) with glutathione (GSH) was investigated in aqueous solution. In this reaction, polymeric solid was not formed and the ammine liberation was slower than that for $cis-[Pt(^{15}NH_3)_2(H_2O)_2]^{2+}$ (3). It is also observed that although in the reactions of 2 with a variety of thiols gave different ultimate product in each reaction, initially the only product observed was a platinum(II) species with chloride and the thiolate ligand bound monodentate through sulfur. In this reaction, after the initial formation of a complex with glutathione bound monodentate $cis-[PtCl(^{15}NH_3)_2(SG)]$ (7), no other product was observed. As with reaction of (3) the bis (thiolate) platinum(II) complex, $cis-[Pt(NH_3)_2(SG)_2]$ (8) was always formed. The dinuclear sulphur-bridged complex, giving a broad peak in ^{15}N NMR, was only present in very tiny amounts in this reaction.

Key words: ^{15}N NMR, glutathione, cisplatin hydrolytes


INTRODUCTION

The main emphasis in the study of cisplatin, *cis*-[PtCl₂(NH₃)₂] (1), a metal based-drug, with biological system has centred on interactions with DNA, and it is widely accepted that the antitumour activity of cisplatin and related compounds is due to primarily to interactions with DNA, however, reactions with other molecules in biological fluids are likely to affect the efficiency with which platinum compounds reach that target, as well as being involved in drug toxicity. The most important appear to be those containing sulphur as a potential coordination site. These include cysteine, methionine, s-adenosyl-L-homocysteine, glutathione and a variety of proteins (Lempers & Reedijk, 1991).

Interactions between platinum and biomolecules containing sulfur have been implicated in the nephrotoxicity of cisplatin (Borch & Pleasants, 1979), antitumour resistance to cisplatin (Fram *et al.*, 1985; Hamilton *et al.*, 1985; Hospers *et al.*, 1985), cell repair mechanism (Eastman, 1987) and in the formation of crosslinks with monofunctional DNA adduct to prevent formation of bifunctional lesions (Eastman, 1987; Eastman & Barry, 1987; Bodener *et al.*, 1986).

Thiols such as cysteine (H₂cys), N-acetylcysteine (H₃accys) and glutathione (GSH) are among the more reactive biological molecules towards the antitumour drug, cisplatin *cis*-[PtCl₂(NH₃)₂] (1). These reactions play a significant role in the metabolism of cisplatin and its hydrolysis products *cis*-[PtCl(NH₃)₂(H₂O)]⁺ (2) and *cis*-[Pt(NH₃)₂(H₂O)₂]²⁺ (3) (Bancroft *et al.*, 1990).

It is established that the DNA and nucleotide reactions are primarily limited by the rate of aquations of cisplatin Bancroft *et al.*, 1990) and that the reactive species from the cisplatin is diammineaquachloroplatinum(II)(2) (Bancroft *et al.*, 1990; Barnham *et al.*, 1994; Bose *et al.*, 1997; Miller & House, 1989a,b, Miller & House, 1990, Miller & House, 1991). Cisplatin, *cis*-[PtCl₂(NH₃)₂] (1) can undergo hydrolysis reactions as shown in Scheme 1.



Scheme 1. Hydrolysis of cisplatin

It has previously been established that thiolate tends to form a bridge between the two metal ions (Odenheimer & Wolf, 1982, Dedon & Borch, 1987, Appleton *et al.*, 1989c). It is important to investigate the chemistry of the thiolate complexes which is relevant to the behaviour of cisplatin metabolites *in vivo* condition, where concentration of Pt-species is low, and there is no bridging.

These reactions were therefore explored in solution with low concentrations of the hydrolytes (1 – 10 mM). ¹⁵N NMR method may be used under these conditions if there is 100% ¹⁵N in the ammine ligands.

This paper describes the reaction of cisplatin hydrolytes, *cis*-[PtCl(¹⁵NH₃)₂(H₂O)₂]⁺ (2) with glutathione (GSH), a sulphur-containing ligand. The other papers related to this work which discuss the reaction of complexes 2 with other thiols have also been submitted to be published elsewhere (Hadi, 2006a,b,c).and reactions of 3 with different sulphur-containing ligands has been published (Hadi, 2005).

EXPERIMENTAL

Starting Materials

¹⁵N-labelled (NH₄)₂SO₄ (> 98% ¹⁵N) was obtained from Novachem (Melbourne Australia), and was used to prepare cisplatin which was prepared by the known procedure (Dhara, 1970). Glutathione was purchased from Sigma Aldrich and used without further purification.

Preparation of *cis*-[PtCl(¹⁵NH₃)₂(H₂O)]⁺ (2)

The typical procedure used to prepare *cis*-[PtCl(¹⁵NH₃)₂(H₂O)]⁺ (2) was by converting *cis*-[Pt(¹⁵NH₃)₂(H₂O)₂]²⁺ (3) to *cis*-[PtCl(¹⁵NH₃)₂(H₂O)]⁺ (2) and was based on the procedure described by Appleton *et al.* (1985). This was done by the slow addition of one mol equivalent of a solution of NaCl, by using a small syringe to deliver one drop every 2 minutes and stirring the mixture reaction continuously. This is the most critical step, since this will affect the formation of *cis*-[PtCl(¹⁵NH₃)₂(H₂O)]⁺. Normally the solution will contain mostly *cis*-[PtCl(¹⁵NH₃)₂(H₂O)]⁺ (80 – 87%), *cis*-[Pt(¹⁵NH₃)₂(H₂O)₂]²⁺ (7 – 15 %) and cisplatin (5 – 10 %) (based on the peak heights in the ¹⁵N NMR or integration in ¹H NMR). It is not possible to obtain a solution containing 100 % *cis*-[PtCl(¹⁵NH₃)₂(H₂O)]⁺.

Typical Reaction Condition

The reactions were carried out under Argon gas to minimise the oxidation of the thiols. No buffer was added as it reacts with the starting material used. The solid GSH was added to a small bottle containing a solution of 2 with pre-measured pH. Within 10 to 15 seconds the mixture reaction was transferred to a 5-mm NMR tube, then placed in the AV400 NMR spectrometer (already tuned for ¹⁵N NMR) and accumulation of 40.54 MHz ¹⁵N NMR spectra was commenced. The reaction proceeded quite fast, one

experiment was left it run for about 6 – 10 hours continuously (in multi experiment mode).

NMR Spectra

The 1D 40.54 MHz ¹⁵N NMR spectra were recorded using DEPT pulse sequence (Berners-Price & Kuchel, 1990) to increase the sensitivity in a Bruker Avance 400 MHz spectrometer with a 5 mm broadband multinuclear probe. The number of scans used to obtain spectra was normally 250 - 500. A recycle time of 3.54 s was used with pulse width of 12.55 μs (tilt angle of 45 degrees). The number of data points used was 32 K. Chemical shifts are reported relative to 2.5 M (¹⁵NH₄)₂SO₄ in 1 M H₂SO₄ (δ_N = 0.00) in coaxial capillary.


RESULTS AND DISCUSSION

The reaction of 2 with glutathione gave similar products to those with H₃accys, except that in this reaction the formation of (bis-thiolate) platinum products is more rapid. It was observed that the polymeric solid was not formed and the ammine liberation was slower compare to that for 3 (Hadi, 2005). Initially the only product observed was a platinum(II) species with chloride and the thiolate ligand bound monodentate through sulphur. The reactions that occurred are summarized in Scheme 2 and the NMR parameters for reaction products are presented in Table 1.

Table 1. ¹H and ¹⁵N NMR parameters of the complexes from reaction of 2 and glutathione

Complex	δ _H (ppm)			δ _N (ppm)		
	<i>trans</i> to S	<i>trans</i> to Cl	<i>trans</i> to O	<i>trans</i> to S	<i>trans</i> to Cl	<i>trans</i> to O
<i>cis</i> -[PtCl(¹⁵ NH ₃) ₂ (H ₂ O)]		4.22	4.17		-65.87	-89.22
<i>cis</i> -[PtCl(¹⁵ NH ₃) ₂ (SG)]	3.83	4.08		-43.18	-62.45	
<i>cis</i> -[Pt(NH ₃) ₂ (SG) ₂](8)	3.77			-41.59		

All reactions were carried out in H₂O/5%D₂O.



Scheme 2. Reaction of (2) with GSH

Previous extensive studies (Appleton *et al.*, 1988) in related system have shown that the sulphydryl group of H₃accys is the preferred binding site of platinum. The presence of free ammonium ion was readily detected from the ¹⁵N NMR spectrum of the solution if the reference capillary containing ¹⁵NH₄⁺ was removed. The signal from free ammonium ion was detected after 30 minutes and continued to grow with time. This ammonium ion is from ammine loss which occurred due to the high *trans* effect of thiolate sulphur.

When solid GSH was added to the solution of 2 in 1 : 1 mol ratio, the ¹⁵N NMR spectrum 1 h after mixing (Fig. 1), showed one pair of new peaks with equal intensities (δ_N – 43.18. and –62.45 ppm) and one other peak at δ_N –41.59 ppm. The pair of peaks corresponded to a species with ammine *trans* to sulfur and chloride respectively (Appleton *et al.*, 1985, Appleton *et al.*, 1988, Appleton *et al.*, 1989a, Appleton *et al.*, 1989b) and was assigned to complex *cis*-[PtCl(¹⁵NH₃)₂(SG)](7) (Scheme 2). The other peak, δ_N –41.59 ppm, present must be due to the formation of bis (thiolate) platinum(II) *cis*-[Pt(NH₃)₂(SG)₂](8), this platinum(II) species was also observed in the reaction of 3 with GSH (Hadi, 2006a).




Fig. 1. 40.54 MHz ¹⁵N DEPT NMR spectrum of a solution obtained from the reaction of 7.4 mM 2 and GSH in 1:1 mol ratio, 1 h after mixing

No further change with time was observed. After 12 hours, the peaks remaining were from unreacted (I), free ammonium ion released, and bis (thiolate) complex (8). When the reaction was carried out with 1 : 2 mol ratio (or GSH in excess), there was no difference in the product formation, except the formation of complex 8 was much faster.

CONCLUSIONS

Compounds containing monodentate S-bound thiolate are moderately stable in solution at low concentration. Ammine loss *trans* to sulfur is rapid, especially in the chelate complex. The results presented here, is believed, represent the first adequate characterization of these species in such

reaction. The other reactions of the compound 2 and 3 with other sulphur-containing ligand will be published elsewhere.

ACKNOWLEDGMENTS

The author deeply thanks to Prof. T.G. Appleton (University of Queensland, Australia) for his help, guidance and discussion. I am also grateful for the award of Scholarship from DUE Project Universitas Lampung, Indonesia and Department of Chemistry, University of Queensland, Australia. Thanks also go to Ms. L. Lambert and Dr. T. Lee for their assistance in the N.M.R. experimentation.

REFERENCES

- Appleton, T.G., Hall, J.R., Ralph, S.F. (1985). Reactions of Platinum(II) Aqua Complexes. 3. Multinuclear (^{15}N , ^{195}Pt , ^{13}C and ^1H) NMR Study of the Reactions of Aqua and Hydroxo Complexes with Glycine and (Methyl-imino)diacetic Acid. *Inorg. Chem.*, 20, 673-677.
- Appleton, T.G., Connor, J.W., Hall, J.R. (1988). S,O versus S,N-Chelation in the Reactions of the *cis*-Diamminediaqua-platinum(II) Cation with Methionine and S-Methylcysteine. *Inorg. Chem.*, 27, 130-137.
- Appleton, T.G., Hall, J.R., Prenzler, P.D. (1989a). Reaction of the *cis*-Diamminediaquapl原因(II) Cation with N-Acetylglycine. *Inorg. Chem.*, 28, 815-819.
- Appleton, T.G., Hall, J.R., Ralph, S.F., Thompson, C.S.M. (1989b). NMR Study of the Acid-Base Equilibria and other Reactions of Ammineplatinum Complexes with Aqua and Hydroxo Ligands. *Inorg. Chem.*, 28, 1989-1993.
- Appleton, T.G., Connor, J.W., Hall, J.R., Prenzler, P.D. (1989c). NMR Study of the Reactions of the *cis*-Diamminediaqua-platinum(II) Cation with Glutathione and Amino Acids Containing a Thiol Group. *Inorg. Chem.*, 28, 2030-2037.
- Bancroft, D.P., Lepre, C.A., Lippard, S.J. (1990). ^{195}Pt NMR Kinetic and Mechanistic Studies of *cis*- and *trans*-Diammine-dichloroplatinum(II) Binding to DNA. *J. Am. Chem. Soc.*, 112, 6860-6871.
- Barnham, K.J., Djuran, M.I., Murdoch, P.S., Sadler, P.J. (1994). Intermolecular Displacement of S-Bound L-Methionine on Platinum(II) by Guanosine-5'-monophosphate: Implications for the Mechanism of Action of Anticancer Drugs. *J. Chem. Soc. Chem. Comm.*, 721-722.
- Berners-Price, S.J., Kuchel, P.W. (1990). Reactions of *Cis*- and *Trans*-[PtCl₂(NH₃)₂] with Reduced Glutathione Inside Human Red Blood Cells, Studied by ^1H and ^{15}N -{ ^1H } DEPT NMR. *Inorg. Biochem.*, 3, 327-345.
- Bodenner, D.L., Dedon, P.C., Keng, P.C., Katz, J.C., Borch, R.F. (1986). Effect of diethyldithiocarbamate on *cis*-Diammine-dichloroplatinum(II)-induced Cytotoxicity, DNA Cross-Linking, and γ -Glutamyl Trans-peptidase Inhibition. *Cancer Res.*, 46, 2745-2750.
- Borch, R.F. & Pleasants, J.M. (1979). Inhibition of *cis*-platinum nephrotoxicity by diethyldithiocarbamate rescue in a rat model. *Proc. Natl. Acad. Sci. U.S.A.*, 26, 6611-6614.
- Bose, R.N., Ghosh, S.K., Moghaddas, S. (1997). Kinetic Analysis of the *cis*-Diamminedichloroplatinum(II)-Cysteine Reaction: Implications to the Extent of Platinum-DNA Binding. *J. Inorg. Biochem.*, 65, 199-205.
- Dedon, P.C., Borch, R.F. (1987). Characterization of the Reactions of Platinum Antitumour Agents with Biologic and Nonbiologic Sulfur-containing Nucleophiles. *Biochem. Pharmacol.*, 36, 1955-1964.
- Dhara, S.C. (1970). A Rapid Method for the Synthesis of *cis*-[Pt(NH₃)₂Cl₂]. *Indian. J. Chem.*, 8, 193-194.
- Eastman, A. (1987). Cross-Linking of Glutathione to DNA by Cancer Chemotherapeutic Platinum Coordination Complexes. *Chem. Biol. Interact.*, 61, 241-248.
- Eastman, A., Barry, M.A. (1987). Interaction of *trans*-Diamminedichloroplatinum(II) with DNA: Formation of Monofunctional Adducts and Their Reaction with Glutathione. *Biochemistry*, 26, 3303-3307.
- Fram, R.J., Cusick, P.S., Wilson, J.M., Marinus, M.G. (1985). Mismatch Repair of *cis*-

Diamminedichloroplatinum(II)-Induced DNA Damage. *Mol. Pharmacol.*, 28, 51-55.

Diammine(aqua)chloroplatinum(II) Cation. *Inorg. Chim. Acta*, 166, 189-197.

Hadi, S. (2005). Reactions of Cisplatin Hydrolytes with Thiols 1: Reaction of $cis-[Pt(^{15}NH_3)_2(H_2O)_2]^{2+}$ with N-Acetyl-L-Cysteine. *J. Sains Tek.*, 11(2): 111-117.

Miller, S.E., House, D.A. (1990). The Hydrolysis Products of *cis*-Diamminedichloroplatinum(II). 3. Hydrolysis Kinetics at Physiological pH. *Inorg. Chim. Acta*, 173, 53-60.

Hadi, S. (2006a). Reactions of Cisplatin Hydrolytes with Thiols 3: Reactions of $cis-[Pt(^{15}NH_3)_2(H_2O)_2]^{2+}$ with Glutathione. Submitted to *Makara Seri Sains, Universitas Indonesia*.

Miller, S.E., House, D.A. (1991). The Hydrolysis Products of *cis*-Diammine-dichloroplatinum(II). 4. The Anation Kinetics of $cis-Pt(X)(NH_3)_2(OH_2)^+$ ($X = Cl, OH$) with glycine, monohydrogen malonate and chloride. *Inorg. Chim. Acta*, 187, 125-132.

Hadi, S. (2006b) Reactions of $cis-[PtCl(^{15}NH_3)_2(H_2O)]^+$, The First Cisplatin Hydrolytes with Thiols 1: Reactions with L-cysteine, Submitted for Publication

Odenheimer, B., Wolf, W. (1982). Reactions of Cisplatin with Sulfur-containing Amino Acids and Peptide 1. Cysteine and Glutathione. *Inorg. Chim. Acta*, 66, L41-L42.

Hadi, S. (2006c) Reactions of $cis-[PtCl(^{15}NH_3)_2(H_2O)]^+$, The First Cisplatin Hydrolytes with Thiols 2: Reactions with N-acetylcysteine, Submitted for Publication

Hamilton, T.C., Winker, M.A., Louie, K.G., Behrens, B.C., Tsuruo, T., Grtzing, K.G., McKay, W.M., Young, R.C., Ozols, R.F. (1985). Augmentation of Adriamycin, Melfhalin, and Cisplatin Cytotoxicity in Drug-Resistant and -Sensitive Human Ovarian Carcinoma Cell Lines by Buthionine Sulfoximine Mediated Glutathione Depletion. *Biochem. Pharmacol.*, 34, 2583-2586.

Hospers, G.A.P., Mulder, N.H., Dejong, B., de Ley, L., Uges, D.R.A., Fichtinger-Schepman, A.M.J., Scheper, R.J., de Vries, E.G.E. (1988). Characterization of a Human Small Cell Lung Carcinoma Cell Line with Acquired Resistance to *cis*-Diamminedichloroplatinum(II) *in Vitro*. *Cancer Res.*, 48, 6803-6807.

Lempers, E.L.M. & Reedijk, J. (1991). Interactions of Platinum Amine Compounds with Sulfur-Containing Biomolecules and DNA Fragments. *Adv. Inorg. Chem.*, 37, 175-217.

Miller, S.E., House, D.A. (1989). The Hydrolysis Products of *cis*-Diamminedichloroplatinum(II). 1. The Kinetics of Formation and Anation of the *cis*-Diammine(aqua)-chloroplatinum(II) Cation in Acidic Aqueous Solution. *Inorg. Chim. Acta*, 161, 131-137.

Miller, S.E., House, D.A. (1989). The Kinetics of Formation and Anation of the *cis*-