

THE PROFILE OF VAGINAL TEMPERATURE AND CYTOLOGY OF VAGINAL SMEAR IN BALI CATTLE DURING ESTRUS CYCLE PHASE

Indira P.N., Kustono and Ismaya

*Master Program of Animal Science, Department of Animal Physiology and Reproduction,
Faculty of Animal Science, Gadjah Mada University,
Jl Fauna No 3 Karangmalang, Yogyakarta - Indonesia
Corresponding E-mail: indira.putri.n@mail.ugm.ac.id*

Received July 04, 2014; Accepted August 27, 2014

ABSTRAK

Penelitian ini bertujuan untuk mengevaluasi kondisi fisiologis sapi Bali selama birahi. Sembilan ekor sapi Bali digunakan dalam penelitian sebagai sampel. Metode deskriptif digunakan dalam penelitian ini. Pengukuran suhu vagina dilakukan dengan termometer digital dan sitologi vagina dengan bantuan *cottonbud* yang diusapkan pada *glass* objek. Sampel smear vagina diwarnai dengan pewarnaan Giemsa yang selanjutnya diamati dengan menggunakan mikroskop. Hasil penelitian menunjukkan bahwa suhu vagina adalah $38,39 \pm 0,29$ °C; $38,05 \pm 0,15$ °C; $37,4 \pm 0,74$ °C; $37,86 \pm 0,3$ °C yang meliputi fase diestrus, metestrus, diestrus, dan proestrus. Berdasarkan sitologi vagina smear, fase estrus memiliki lebih banyak sel kornifikasi superfisial dan sel *intermediate* yang menyebabkan konsentrasi estrogen meningkat. Berdasarkan hasil analisis profil suhu vagina, dapat disimpulkan bahwa sitologi dari vagina *smear* setiap jenis fase siklus estrus memiliki karakteristik sendiri dan pada gambaran setiap sel terdapat perbedaan fase siklus estrus.

Kata kunci: sapi Bali, siklus estrus, sitologi vagina smear, suhu

ABSTRACT

The objective of the study was to evaluate physiological condition of Bali cattle during estrus. Nine cattles were used in the research. The method of the research was descriptive analysis. The measurement of vaginal temperature was conducted by using digital thermometer and cytology of vaginal smear was done by using cotton bud swabbed on object glass. It was washed by methanol and was stained with Giemsa staining and was observed by microscope. The results indicated that vaginal temperature were 38.39 ± 0.29 °C; 38.05 ± 0.15 °C; 37.4 ± 0.74 °C; 37.86 ± 0.3 °C at estrus, metestrus, diestrus, proestrus phase, respectively. On the basis of the cytology of vaginal smear, estrus phase had more superficial cornification and intermediate cells causing estrogen concentration increased. In conclusion, the profile of vaginal temperature showed that each kind of estrus cycle phase had characteristic within estrus. The cytology of vaginal smear showed various types of cells preferences in different phase of estrus cycle.

Keywords: Bali cattle, estrus cycle, temperature, cytology vaginal smear

INTRODUCTION

Bali cattle is potential and popular for developing germplasm in Indonesia because they have high productivity not only carcass but also body weight. Bali cattle has special genetic that is easy to adapt to bad environment. It was stated by Handiwirawan and Subandriyo (2007), Bali cattle

were called pioneer cattle from the aspect of reproduction, Bali cattle has high fertility rate, which is about 83-86%.

Currently, Bali cattle has some problems such as genetic decline, high inbreeding, low adaptation (Sariubang *et al.*, 1998) Low estrus detection efficiency is known as one of the main causes of poor fertility rate of Bali cattle (Noor *et*

al., 2002). This lead to prolonged calving interval and calving to conception intervals, and also to low herd technical and economic performance, even in developed countries (Bage *et al.*, 2002). This situation results of poor reproductive practices that commonly observed in livestock production systems in the tropics as well as in Indonesia (Hardjopranojoto, 1995).

Visual methods for estrous detection are well documented, but method usage and their efficiency are still controversial (Fahey *et al.*, 2002). The silent heat of Bali cattle is a supplementary difficulty to visual estrus detection (Salem *et al.*, 2006). Hormones profile monitoring appear to be very expensive for low income livestock production systems (Wattemann *et al.*, 2003). Cytology vaginal smear changes during estrus cycle have been studied in sheep (Bearden *et al.*, 2004), but little is known about this issue in cattle, especially in Bali cattle. On the basis of those reason, a research contribute to heat detection efficiency by evaluating cytology vaginal smear combined to vaginal temperature changes during estrus cycle were conducted. It can protect the cattle from repeating breeding. This research was aimed to help animal farm and researcher to know more about the physiological condition of Bali cattle during estrus. It was expected that this study will increase the success of breeding of Bali cattle for both natural and insemination.

MATERIALS AND METHODS

Materials

Nine cattles were used in the research which was 3 years old with average body weight of 313.67±20.66 kg. The cattles were fed elephant grass and polar concentrate. The cattle drank mineral water added with *ad-libitum*.

Procedures

The measurement of vaginal temperature was conducted by using digital thermometer (length of 3 to 5 cm) until constant temperature was obtained. Before and after the measurement of temperature, thermometer was sterilized using cotton added with 70% of alcohol. The cotton bud was used for wrapping of cytology vaginal smear from vaginal epithelia tissues into object glass. The object glass was then stained with Giemsa method for 30 to 45 minutes. After being dried, it was observed by the microscope.

Data Analysis

The vaginal temperature was analyzed by using SPSS ver 20 program for Windows (IBM®, 2012). Cytology of vaginal smear was analyzed descriptively according to method of Astuti (2007).

RESULTS AND DISCUSSION

The vaginal temperature of nine Bali cattles are presented the Table 1. The table showed that the vaginal temperature ranged from 37.4 °C to 38.39 °C within estrus cycle phase. The highest temperature was in estrus phase. Kyle *et al.* (1998) reported that in their research, vaginal temperature gradually decreased many days before estrus and it increased during estrus phase. The vaginal temperatures were changed because of the effect of hormonal system. The hormone system of ovarium especially estrogen and luteinizing can create blood vascularization around genital organ on increasing reproduction. Thus, the thermals were streamed by bloods and vaginal temperatures increased because of vascularization process (Hovinen *et al.*, 2008). Estrus cycle phase was affected by hormonal change that happened during ovulation with the increase of the luteinizing hormone on bloods. Acosta *et al.* (2003) also reported the significant correlation between estradiol plasma concentration and luteinizing hormone which increased the vascularization within preovulation cycle on cattle. Other factors were vaginal mucus that was changed within estrus cycle (Rorie *et al.*, 2002). However, this research was still difficult to compare with other different treatments or qualitative research. In addition, many animals had the highest variation on measurement (Lehrer *et al.*, 1995).

The wrapping method of vaginal smear used Giemsa staining method to find out estrus cycle. The result of vaginal smear is presented in Figure

Table 1. The Vaginal Temperature of Bali Cattles

Phase	Vaginal Temperature (°C)
Diestrus	37.40 ± 0.74
Proestrus	37.86 ± 0.30
Estrus	38.39 ± 0.29
Metestrus	38.05 ± 0.15

1. It can be seen in Figure 1 that the difference of cells in each phase and estrus phase was marked by many superficial cells and intermediate cells. Vaginal epithelial cells proportions recorded during estrous phases was similar to report of Marondes *et al.* (2002). The epithelium of vagina was sensitive for estradiol 17- β concentration during its development. The receptor of estradiol 17- β can be obtained on vagina tissues (Bologna, 2001). The part of vaginal epithelium consisted of four cells type where almost three cells on vaginal epithelium always had cornification that were parabasal cell, intermediate cell and superficial cell. However, cornification of basal cell only happened during trauma and radiation (Bologna, 2001). Intermediate and parabasal cells were indicated by vaginal smear on metestrus, diestrus, and proestrus phase. Perez *et al.* (2009) reported

that superficial parabasal cells were specifically marked to detect the activity of ovarium before puberty. Similar finding was also reported by Schuttle (2010) stating that superficial cells in vaginal smear were specifically marked for the condition of estrogen hormone. Superficial cells were found on vaginal smear during estrus phase while estrogen concentration was increased (Junaidi, 2005). Superficial cells came out of basal membrane and they developed than other cells. Thus, this cells were mature because of cornification after development process stimulated by estradiol 17- β (Bologna, 2001). Mehta *et al.* (2000) reported that changing steroid hormone is related with vaginal smear's changes. This was supported by Garofalo and Tasendo (2000) who stated about the effect of estradiol 17- β hormone and the cytology of vaginal smear,




Figure 2. The cytology of Vaginal Smear with Giemsa Stain during Estrus Cycle Phase in Bali Cattle. 1:Leukocyte, 2: Superficial cell, 3: Intermediate cell, 4: Parabasal cell

which is typically similar with estrus behavior.

CONCLUSION

Vaginal temperature gave complete information about physiology within estrus cycle phase and the cytology of vaginal smear can explain that estrus phase had more superficial cornification and intermediate cells.

ACKNOWLEDGEMENT

The authors would like to highly appreciate to the chair of Field for Education, Research, and Farming Development, Gadjah Mada University for giving permission to use animals and barn facilities during this research. The present research is granted and funded by Indonesian Endowment Fund for Education, Ministry of Finance, Republic of Indonesia.

REFERENCES

- Acosta, T.J., K.G. Hayashi, M. Mohtani and A. Miyamoto. 2003. Local changes in blood flow within the preovulatory follicle wall and early corpus luteum in cows. *Reproduction*. 125:759-767
- Astuti, M. 2007. Pengantar ilmu statistik untuk peternakan dan kesehatan hewan. Binasti Publisher. Bogor.
- Bage, R., H. Gustafsson, B. Larsson, M. Forsberg and H. Rodriguez-Martinez. 2002. Repeat breeding in dairy heifers: Follicular dynamics and oestrous cycle characteristic in relation to sexual hormone patterns. *Theriogenology* 57:2257-2269
- Bearden, H.J., J.W. Fuquay and S.T. Willard. 2004. *Applied Animal Reproduction*. 6th ed. Pearson Prentice Hall, USA.
- Bologna, J.W. 2001. Genitourinary problems associated with menopause. *Anim. Reprod. Sci.* 90:51-55
- Fahey, J., K. O'Sullivan, J. Crilly and J.F. Mee. 2002. The effect of feeding and management practices on calving rate in dairy herds. *Anim. Reprod. Sci.* 74:133-150
- Garofalo, E. G. and C. Tasendo. 2000. Uterine estrogen and progesterone receptors in prepubertal ewes: distribution in myometrium, endometrium and caruncle. *Vet. Res.* 27:177-183
- Handiwirawan, E. and Subandriyo. 2007. Perbaikan Mutu Genetik Sapi Bali. Balai Penelitian Ternak. Bogor.
- Hardjopranojoto, H.S. 1995. Ilmu kemajiran ternak. Airlangga University Press, Surabaya.
- Hovinen, M., J. Silvonene, S. Taponen, L. Hanninen, M. Pastell, A. M. Aisla and S. Pyorala. 2008. Detection clinical mastitis with the help of a thermal camera. *J. Dairy Sci.* 91:4592-4598
- IBM®. 2012. SPSS Statistics Editions. IBM Corporation Software Group. United States America.
- Junaidi, A. 2005. Reproduksi dan obstetri pada anjing. Gadjah Mada University Press. Yogyakarta.
- Kyle, B.L., A.D. Kennedy and J.A. Small. 1998. Measurement of vaginal temperature by radioelementary for the prediction of estrus in beef cows. *Theriogenology* 49:1437-1449
- Lehrer, A.R., Lewis, G.S., Mc Millan, W.H. and E. Aizinbud. 1995. Bio-impedance monitoring of genital tissues of cows as an aid in cattle reproductive management—a review. *Proceedings of the New Zealand Society of Animal Production* 55:221-223
- Marcondes, F.K., F.J., Bianchi and A.P. Tanno. 2002. Determination of the estrous cycle phase of rats: some helpful considerations. *J. Brazilian Archiv. Biol. Technol.* 4(A):600-614
- Mehta, R. R., J.M. Jenco, L.V. Gaynor and R.T. Jr. Chartterton. 2000. Relationships between ovarian morphology, vaginal cytology, serum progesterone, and urinaria immunoreactive pregnanediol during the menstrual cycle of the cynomolgus monkey. *Biol. Reprod.* 47:408-417
- Noor, R. R, A. Farajallah and M. Karmita. 2002. Pengujian kemurnian sapi Bali dengan analisis hemoglobin dengan metode Isoelectric Focusing. *Hayati* 8: 107-111
- Perez, M., Mendoza, M. E. and M.C. Romano. 2009. Exfoliative vaginal cytology and plasma levels of estrone and oestradiol 17 in young and adult goats. *Small Rum. Res.* 33:153-158
- Rorie, R.W., T.R. Bilby and T.D. Lester. 2002. Application of electronic estrus detection technologies to reproductive management of cattle. *Theriogenology* 57:137-148
- Salem, M.B., M. Djemali, C. Kayouli and A. Majdoub. 2006. A review of environmental and management factors affecting the reproductive performance of Holstein-

- Friesian dairy herds in Tunisia. *Livest. Res.* 18(4):123-129
- Sariubang, M., D. Pasambe and Chalidjah. 1998. Pengaruh kawin silang terhadap performa hasil turunan pertama (F1) pada sapi Bali di Sulawesi Selatan. *Prosiding Seminar Nasional Peternakan dan Veteriner, Bogor.* June 19-20, 1998. P. 731-737
- Schuttle, A.P. 2010. Technique and cytology morphology. *J. Small. Anim. Pract.* 18:301-306
- Wattemann, R.P., C.A. Lents, N.H., Ciccioli, F.J. White and I. Rubio. 2003. Nutritional and suckling-mediated anovulation in beef cows. *J. Anim.Sci.* 81 (E. Suppl.2):E48-E59