

## Production, Competition Indices, and Nutritive Values of *Setaria splendida*, *Centrosema pubescens*, and *Clitoria ternatea* in Mixed Cropping Systems in Peatland

A. Ali<sup>a,d,\*</sup>, L. Abdullah<sup>b</sup>, P. D. M. H. Karti<sup>b</sup>, M. A. Chozin<sup>c</sup>, & D. A. Astuti<sup>b</sup>

<sup>a</sup>Major Program of Animal Nutrition and Feed Science, Graduate School, Bogor Agricultural University

<sup>b</sup>Department of Nutrition and Feed Technology, Faculty of Animal Science, Bogor Agricultural University  
Jln. Agatis, Kampus IPB Darmaga Bogor 16680, Indonesia

<sup>c</sup>Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University  
Jln. Meranti Kampus IPB Darmaga, Bogor 16680, Indonesia

<sup>d</sup>Department of Animal Science, Faculty of Agriculture and Animal Science, UIN Suska Riau  
Jln. H. R. Soebrantas No.155 KM 18, Simpang Baru Panam, Pekanbaru 28293, Indonesia

(Received 11-09-2013; Reviewed 09-10-2013; Accepted 03-12-2013)

### ABSTRACT

This research was conducted to evaluate production, different competition indices and nutritive value of *Setaria splendida*, *Centrosema pubescens*, and *Clitoria ternatea* in monoculture and mix cropping system on peat soil land. The experiment was set up in a randomized complete block design with five treatments and three replications. The five treatments were: *S. splendida* sole cropping (SS), *C. pubescens* sole cropping (CP), *C. ternatea* sole cropping (CT), *S. splendida* and *C. pubescens* mix cropping (SS/CP) and *S. splendida*/*C. ternatea* mix cropping (SS/CT). The DM yield of *S. splendida* in mixed cropping with *C. pubescens* increased 43.4% and in mix cropping with *C. ternatea* increased 15.7% compared to sole *S. splendida*. The value of land equivalent ratio of SS/CP ( $LER_{SS/CP}$ ) was >1. The  $LER_{SS/CT}$  value was <1. The crowding coefficient value of *S. splendida* ( $K_{SS}$ ) was higher than  $K_{CP}$  and  $K_{CT}$ . The total value of  $K_{SS/CP}$  and  $K_{SS/CT}$  were >1. The competition ratio (CR) values of *S. splendida* in both mix cropping were >1. The aggressivity (A) values of *S. splendida* in both mix cropping were positive. The crude protein, NDF and ADF content of forage were not affected by mix cropping system. In conclusion, mix cropping in peatland do not affect productivity and nutritive value of *S. splendida*, *C. pubescens*, and *C. ternatea*. *S. splendida* is more effective in exploiting environmental resources when intercropped with *C. pubescens* compared to *C. ternatea* on peatland.

**Key words:** cropping system, forage, production, quality

### ABSTRAK

Penelitian bertujuan untuk mengevaluasi produksi, indeks kompetisi yang berbeda dan nilai nutrisi *Setaria splendida*, *Centrosema pubescens*, dan *Clitoria ternatea* dengan sistem pertanaman tunggal dan campuran di lahan gambut. Penelitian menggunakan rancangan acak kelompok yang terdiri atas 5 perlakuan dan 3 ulangan, yaitu *S. splendida* yang ditanam tunggal (SS), *C. pubescens* yang ditanam tunggal (CP), *C. ternatea* yang ditanam tunggal (CT), *S. splendida* dan *C. pubescens* yang ditanam campuran (SS/CP), serta *S. splendida* dan *C. ternatea* yang ditanam campuran (SS/CT). Produksi bahan kering *S. splendida* yang ditanam campuran dengan *C. pubescens* meningkat 43.4% dan pertanaman campuran dengan *C. ternatea* meningkat 15.7% dibandingkan pertanaman tunggal. Nilai land equivalent ratio SS/CP ( $LER_{SS/CP}$ ) adalah >1.  $LER_{SS/CT}$  adalah <1. Nilai *crowding coefficient* (K) *S. splendida* ( $K_{SS}$ ) adalah lebih tinggi dari  $K_{CP}$  and  $K_{CT}$ . Nilai total K plot SS/CP dan SS/CT adalah >1. Nilai *competition ratio* (CR) *S. splendida* pada kedua pertanaman campuran adalah >1. Nilai *agressivity* (A) *S. splendida* pada kedua pertanaman campuran adalah positif. Kandungan PK, NDF, dan ADF hijauan tidak terpengaruh oleh sistem pertanaman campuran. Dapat disimpulkan bahwa pertanaman campuran di lahan gambut tidak mempengaruhi produktivitas dan nilai nutrisi *S. splendida*, *C. pubescens*, dan *C. ternatea*. *S. splendida* lebih kompetitif dan dominan dibanding *C. pubescens* dan *C. ternatea* di lahan gambut.

**Kata kunci:** hijauan, sistem penanaman, produksi, kualitas

\*Corresponding author:  
E-mail: ali\_arsyadi@yahoo.com

## INTRODUCTION

The availability of low cost feed resources is important for successful ruminant production in smallholder farming system in the tropics (Noula *et al.*, 2004). The major limitation of livestock production is the lack of suitable fodder crops that can produce green forage year round (Mutimura & Everson, 2012). One of the logical approaches to increasing forage production to overcome these problems is pasture improvement by grass and legume mix cropping system (Ajayi *et al.*, 2007). Mix cropping is one of the multiple cropping system, has been practiced by farmer for many years in various ways and most areas (Zhang & Li, 2003). The advantage of grass/legume mix cropping in pasture is the possibility of N addition from the legume to the grass. Depending on the nitrogen content of the soil and the mix of legumes and grasses in a pasture, during the growing season legumes transfer approximately 40 kg N ha<sup>-1</sup> to neighbouring plants (Pirhofer-Walzl *et al.*, 2012). Grass/legume mixtures are more sustainable and better overcome unfavorable conditions as compared to their pure cultivation (Peyraud *et al.*, 2009). They are more productive than pure stands and each species contributes the productivity in varying degrees (Vasileva & Vasilev, 2012).

The mayor limitation of pasture development is the lack of mineral soil for forage. The development pasture can be cultivated in peatland (*Organosol*). Organosol are formed by progressive accumulation plant materials over time (Agus & Subiksa, 2008). Limitations in obtaining high forage production in peatland due to differences in the characteristics of peat soil with mineral soil. Naturally, the peat has a low fertility rate because of the low nutrient content, and generally, the peat has a relatively high level of acidity. The purpose of this study was to evaluate the dry matter production, different competition indices and nutritive value of *S. splendida*, *C. Pubescens*, and *C. ternatea* in sole and mixed cropping system in peatland.

## MATERIALS AND METHODS

This study was conducted at research farm of Faculty of Agriculture and Animal Science of UIN Suska Riau and Laboratory of Research Center of Biological Resources and Biotechnology, PAU, Bogor Agricultural University from October 2011 to November 2012. The forages investigated were *S. splendida* (grass), *C. Pubescens*, and *C. ternatea* (legumes). The experiment was set up in randomized complete block design with five treatments and three replicates. Five treatments were: *S. splendida* sole cropping (SS), *C. pubescens* sole cropping (CP), *C. ternatea* sole cropping (CT), *S. splendida*/*C. pubescens* mix cropping (SS/CP) and *S. splendida*/*C. ternatea* mix cropping (SS/CT).

### Plot, Planting Density and Fertilizing

This experiment was conducted in peatland type soil (type *sapric*). The pH and mineral soil content of experimental land in the form of N,C,C/N, K and P were 5.54, 0.14%, 7.20%, 51.43, 2.48 me/100g, and 30.18

ppm, respectively. The size of experimental land was 11.5x17 m and divided in three blocks. Each block was sub-divided into five plots (each plot size of 2.5 x 5 m), namely SS, CP, CT, SS/CP and SS/CT. The forages were cultivated in September 2011. The plot had a planting density of 50 plants/plot (planting space was 50 x 50 cm) and was maintained under rain-fed condition. The proportion of grass and legume in mix culture plots was 50 : 50. The basal fertilizer was organic fertilizer (cattle manure) applied at the rate 10 t/ha and was applied two weeks before planting, and inorganic fertilizers (NPK) at the rate of 50 kg/ha/yr of was applied two weeks after planting (surrounding the plant).

### Propagating, Pruning, Harvesting and Sample Procedure

*S. splendida* was propagated by cutting while, *C. pubescens* and *C. ternatea* was propagated by seed. Pruning was done after 2 mo of grown. The forages were trimmed approximately 20 cm above the ground by using a pair of garden shear. This would allow a new and uniform re-growth from where the experimental samples were later harvested. Grass and legume foliage were harvested six time a year with 60 d cutting interval. The plants were cut approximately 20 cm from the ground from each plot (n= 24 plant) and directly weighed to determine the fresh yield.

### Competition Indices

The competitive behaviour of component forages in grass/legume mix cropping system was determined in terms of land equivalent ratio (LER), competition ratio (CR), aggressivity (A) according to the methods described by Dhima *et al.* (2007) and crowding coefficient (K) was estimated according to the methods of Banik *et al.* (2006). The LER, which measures the effectiveness of mixed cropping in using the environmental resources, compared to sole cropping. The LER values were calculated as follows:  $LER = (LER_{grass} + LER_{legume})$ , where  $LER_{grass} = (Y_{gm}/Y_{ls})$ , and  $LER_{legume} = (Y_{lm}/Y_{gs})$ , where  $Y_{gs}$  and  $Y_{ls}$  are the yields of grass and legume as sole crops respectively, and  $Y_{gm}$  and  $Y_{lm}$  are the yields of grass and legume as mixtures, respectively. When  $LER > 1$ , there is yield advantage. The CR gives a clear idea about which forage is more competitive in association. The CR values were calculated as:  $CR_{grass} = (LER_{grass}/LER_{legume}) \times (Z_{lp}/Z_{gp})$ , and  $CR_{legume} = (LER_{legume}/LER_{grass}) \times (Z_{gp}/Z_{lp})$ , where  $Z_{gp}$  and  $Z_{lp}$  are the proportion of grass and legume in the mixture. If  $CR_{grass} > 1$ , grass is more competitive than legume and if the value is  $< 1$ , grass is less competitive than legume. The reverse is true for  $CR_{legume}$ . The aggressivity (A) is a measure of competitive relationships between two forages in mixed cropping. This was calculated as follows:  $A_{grass} = (Y_{gm}/Y_{gs} \times Z_{gp}) - (Y_{lm}/Y_{ls} \times Z_{lp})$  and  $A_{legume} = (Y_{lm}/Y_{ls} \times Z_{lp}) - (Y_{gm}/Y_{gs} \times Z_{gp})$ . Thus if  $A_{grass} = 0$ , both crops are equally competitive, If  $A_{grass}$  is positive, then the grass is dominant, and if  $A_{grass}$  is negative, the grass is weak. The relative crowding coefficient (K), which is a measure of the relative dominance of one species over the other in a mixture, was calculated as:  $K =$

( $K_{grass} \times K_{legume}$ ), where  $K_{grass} = Y_{gm} \times Z_{lp} / [(Y_{gs} - Y_{gm}) \times Z_{gp}]$ , and  $K_{legume} = Y_{lm} \times Z_{gp} / [(Y_{ls} - Y_{lm}) \times Z_{lp}]$ , when the value of  $K$  is  $> 1$ , there is yield advantage; when  $K$  is  $= 1$ , there is no yield advantage; and, when  $K < 1$  there is a disadvantage.

**Chemical Analysis**

Fresh samples of grass and legume from each plot (about 500 g) were dried in air-forced oven at 60 °C for 48 h, and ground to pass through a 1 mm sieve for chemical analysis. The dry matter (DM) and crude protein (CP) were determined according to the AOAC (2005) procedure. Neutral detergent fiber (NDF) and acid detergent fiber (ADF) are estimated according to the method of Van Soest *et al.* (1991).

**Statistical Analysis**

Data was analyzed by analysis of variance (ANOVA) based on a completely randomized block design with five treatments and three blocks. Comparison of means were considered significantly different at 5% level ( $P < 0.05$ ) using Duncan’s Multiple Range Test (DMRT).

**RESULTS AND DISCUSSION**

**Dry Matter Yield of Forages**

**Dry matter (DM) yield per plant.** The effect of mixed cropping on DM yield per plant (g/yr) of *S. splendida*, *C. pubescens* and *C. ternatea* are shown in Table 1. The DM yield of *S. splendida* was not significantly increased by mixed cropping. The DM yield of *S. splendida* in *C. pubescens* increased 43.4% and in *C. ternatea* increased 15.7% compared sole *S. splendida*. Intercropped with *S. splendida* decreased 23.4% DM yield of *C. pubescens*. The decreasing in DM yield due to the impact of the interspecific competition which led *C. pubescens* yield less than those of sole cropping system. Otherwise, *S. splendida* yield higher than those of sole *C. pubescens*. This condition is regularly found in mixed cropping system. A competition increases yield of dominant species, but decreases yield of sub-ordinate species (Marty *et al.*,

Table 1. Dry matter (DM) yield (g/year) per plant of *S. splendida*, *C. pubescens*, and *C. ternatea* under monoculture and mixed cropping system

Forages	DM yield (g/yr)
<i>S. splendida</i>	325± 93
<i>S. splendida</i> in <i>C. pubescens</i> intercrop	466±215
<i>S. splendida</i> in <i>C. ternatea</i> intercrop	376±152
<i>C. pubescens</i>	145± 13
<i>C. pubescens</i> in <i>S. splendida</i> intercrop	111± 27
<i>C. ternatea</i>	205± 77 <sup>a</sup>
<i>C. ternatea</i> in <i>S. splendida</i> intercrop	73± 28 <sup>b</sup>

Note: Mean in *C. ternatea* row with different superscripts differ significantly ( $P < 0.05$ ).

2009). Meanwhile, *S. splendida*/*C. ternatea* intercropping was amensalistic, where the DM yield of *C. ternatea* was significantly ( $P < 0.05$ ) decreased (64.4%) as a result of interaction between *S. splendida* and *C. ternatea* but no significant effect was found on DM yield of *S. splendida*. Limited growth of *C. ternatea* in mix cropping system may be caused by extended growth of *S. splendida* roots than those of *C. ternatea*. It led extensive nutrient uptake by *S. splendida* from the soil than *C. ternatea*. The negative effect on mixed-culture cropping is a competition in uptaking same nutrient, water, light and the other sources (Forrester *et al.*, 2006; Thorsted *et al.*, 2006).

**Dry matter yield per plot.** The effect of cropping system (sole and mix cropping) on DM yield per plot are shown in Table 2. It was recorded that SS/CP and SS/CT mixed cropping in peatland did not influence ( $P > 0.05$ ) the DM yield of forage per plot. Intercropping with *S. splendida* caused the growth of *C. pubescens* and *C. ternatea* to be lower than the sole crop. Growth of *C. pubescens* and *C. ternatea* was hindered by the presence of competition in uptaking nutrient elements in soil (Oseni, 2010). *S. splendida* grew quickly and forming stolons so that more nutrient were absorbed than *C. pubescens* and *C. ternatea*. Intercropping with *C. pubescens* or *C. ternatea* caused *S. splendida* to grow at a faster rate than the monoculture, so the lack of production on the mixed culture plot was filled by the production of *S. splendida*. It has been established that the legume component in a grass/legume mixture would improve availability of nutrient in the soil especially nitrogen content through nitrogen fixation (Dhalika *et al.*, 2006; Pozdissek *et al.*, 2011) and via the mineralization of the N-rich legume litter (Marty *et al.*, 2009)

**Competition Indices**

**Land equivalent ratio (LER).** The value of  $LER_{SS/CP}$  was  $> 1$  (1.10) (Table 3). This value indicated that mixed cropping was more effective over sole or mono cropping with regard to the use of environmental sources for plant growth (Mahapatra, 2011). The value of  $LER_{SS/CT}$  was  $< 1$  (0.76) (Table 3) which indicated the disadvantage of mixed cropping over sole *S. splendida* and *C. ternatea* (Yilmaz *et al.*, 2008; Dhima *et al.*, 2007).

Table 2. Dry matter (DM) yield of forages per plot based on cropping system

Cropping system	Plot	DM yield (t/ha/yr)
Monoculture	<i>S. splendida</i> (SS)	13.01±3.72 <sup>a</sup>
	<i>C. pubescens</i> (CP)	5.84±0.53 <sup>b</sup>
	<i>C. ternatea</i> (CT)	8.20±3.07 <sup>ab</sup>
Mixture	SS/CP	11.55±3.87 <sup>ab</sup>
	SS/CT	9.00±3.49 <sup>ab</sup>

Note: Mean in the same column with different superscripts differ significantly ( $P < 0.05$ ).

Table 3. Competition indices (LER, K, CR and A) of *S. splendida* (SS), *C. pubescens* (CP), and *C. ternatea* (CT) under monoculture and mixed cropping

Competition indices	SS : CP		Total	SS : CT		Total
Land Equivalent Ratio (LER)	0.72	0.38	1.10	0.58	0.18	0.76
Crowding Coefficient (K)	2.53	0.62	1.56	1.38	0.22	0.30
Competition Ratio (CR)	1.88	0.53	-	3.25	0.31	-
Aggressivity (A)	0.67	-0.67	-	0.80	-0.80	-

**Crowding coefficient (K).** The value of  $K_{SS}$  was higher than  $K_{CP}$  and  $K_{CT}$  in mixed cropping (Table 3), indicating an advantage of *S. splendida* over the other legume in the mixed cropping system. The presence of legume improved the growth of *S. splendida* (Mucheru-Muna *et al.*, 2010). Legume root systems did not give tough competition to the grass (Refliaty *et al.*, 2009), as the grass acquire N from legumes through the decomposition and mineralization of N-rich legume plant litter and N transfer from the legume to grass (Pirhofer-Walzl *et al.*, 2012). The total K value on SS/CP plot was >1 and on SS/CT was < 1 (Table 3), indicating that *S. splendida* intercropped with *C. pubescens* contributed to the high productivity per unit of land compared to intercropping with *C. ternatea* (Yilmaz *et al.*, 2008).

**Competition ratio (CR).** The value of  $CR_{SS}$  in *S. Splendida* intercropped with *C. pubescens* and *C. ternatea* were >1 (Table 3), indicating that *S. splendida* was more competitive than legumes, and caused *C. pubescens* and *C. ternatea* growth to be inhibited. Zhang & Li (2003) reported that interspecific competition may occur when two crops are grown together on the same field. Such competition usually decreases survival, growth or reproduction of at least one species.

**Aggressivity (A).** Based on Table 3, the positive value of aggressivity for *S. splendida* in both mixed cropping in peatland showed that *S. splendida* was more dominant over *C. pubescens* and *C. ternatea*. Dominance of *S. splendida* was also reflected in the value of K and CR. This was probably due to more extensive root system of grass than legume (Chen *et al.*, 2004).

### Proportion of Grass and Legume on Mix Cropping System

The proportion of grass to legume in mix cropping system based on SS/CP and SS/CT increased from harvest 1 (December 2011) to harvest 6 (October 2012) (Figure 1a and 1b). The reason for increasing proportion of *S. splendida* may be attributed to nitrogen fixing ability of legume (Yilmaz *et al.*, 2008) and *S. splendida* was more competitive and aggressive than both the legumes (Table 3). Based on DM yield/yr, the proportion grass to legume on SS/CT plot was higher than the SS/CP plot. The proportion of grass to legume on SS/CT and SS/CP plots were 81.13% versus 18.87% and 79.57% versus 20.43%, respectively. This suggested that *C. pubescens* was more competitive than *C. ternatea* when intercropped with *S. splendida* in peatland.

### Nutrient Composition

**Dry matter (DM) content.** The DM content of forage was not affected by cropping system but influenced by the forage type (Table 4). The result of study found that DM content of legume was significantly ( $P < 0.05$ ) higher than grass. This result agree with Ajayi *et al.* (2007) who found that the DM content of grass (*Panicum maximum*) was lower than legumes (*Aeschynomene histrix* and *Stylosanthes guianensis*).

**Crude protein (CP) content.** The CP content of forages was not affected by cropping system but influenced by the forage type (Table 4). This result was contrary with those reported by several researchers (Karadag &



Figure 1a. Proportion of *S. splendida* (■) and *C. pubescens* (□) in mix cropping plot at each harvestFigure 1b. Proportion of *S. splendida* (■) and *C. ternatea* (□) in mixed cropping plot at each harvest

Table 4. The content of dry matter (DM), crude protein (CP), neutral detergent fibre (NDF), and acid detergent fibre (ADF) of *S. splendida*, *C. pubescens*, and *C. ternatea* under monoculture and mixed cropping

Forage	DM (%)	%DM		
		CP (%)	NDF (%)	ADF (%)
<i>S. splendida</i>	14.8±0.3 <sup>b</sup>	7.5±0.5 <sup>c</sup>	72.4±2.1 <sup>a</sup>	37.4±1.1 <sup>b</sup>
<i>S. splendida</i> in <i>C. pubescens</i> intercrop	14.2±0.4 <sup>b</sup>	7.4±0.9 <sup>c</sup>	68.2±0.8 <sup>ab</sup>	36.0±0.5 <sup>b</sup>
<i>S. splendida</i> in <i>C. ternatea</i> intercrop	14.8±0.7 <sup>b</sup>	7.4±0.8 <sup>c</sup>	72.6±1.2 <sup>a</sup>	37.6±0.9 <sup>b</sup>
<i>C. pubescens</i>	24.3±0.4 <sup>a</sup>	17.5±2.2 <sup>a</sup>	67.8±2.1 <sup>ab</sup>	45.6±1.6 <sup>a</sup>
<i>C. pubescens</i> in <i>S. splendida</i> intercrop	23.7±1.3 <sup>a</sup>	17.4±0.9 <sup>a</sup>	66.0±0.7 <sup>b</sup>	45.8±1.7 <sup>a</sup>
<i>C. ternatea</i>	23.0±1.3 <sup>a</sup>	14.8±2.8 <sup>b</sup>	55.7±1.4 <sup>c</sup>	39.2±0.6 <sup>b</sup>
<i>C. ternatea</i> in <i>S. splendida</i> intercrop	24.1±0.9 <sup>a</sup>	15.1±3.1 <sup>b</sup>	52.1±8.2 <sup>c</sup>	37.4±3.9 <sup>b</sup>

Note: Mean in the same column with different superscripts differ significantly (P<0.05).

Buyukburc, 2004; Albayrak *et al.*, 2011; Eskandari, 2012; Njad *et al.*, 2013) who reported that grass intercropped with legume resulted in improved soil fertility through the addition nitrogen by fixation and excretion from the legume component and also increased CP content of nearby grass. The CP content of *S. splendida* that were planted in peatland was slightly lower than the average CP content of *S. splendida* reported by Heuzé & Tran (2012) who showed that the CP content of *S. splendida* was 9.1%. The CP content of *C. pubescens* and *C. ternatea* were higher significantly (P<0.05) than *S. splendida*. This indicated that legumes have a higher nutrient content than grass (Albayrak *et al.*, 2011) particularly CP content (Paulson *et al.*, 2008). These study also showed that CP content of both legumes planted in peatland was generally lower than those found by several researches. Evitayani *et al.* (2004), Omele *et al.* (2011) and Martens *et al.* (2012) reported that CP content of *C. pubescens* was 18.9%, 25.5%, and 23.6%, respectively. However, Mahala *et al.* (2012), Nasrullah *et al.* (2003) and Heinritz *et al.* (2012) reported that CP content of *C. ternatea* was 17%, 18.28%, and 19%, respectively. These evidences showed that CP content is influenced by forage type, environmental condition and land condition (Jayanegara & Sofyan, 2008). Soil conditions will affect the availability of nutrients and soil organic matter (Sabrina *et al.*, 2013) which led to varied growth and forage quality.

*S. splendida*/*C. pubescens* and *S. splendida*/*C. ternatea* mixed cropping did not increase CP production per unit area of land. Even, the CP production of SS/CT plot tended to be lower than SS plot (Table 5). This was due to the low dry matter production of *C. pubescens* and *C. ternatea* intercropped with *S. splendida* (Table 1).

**Neutral detergent fibre (NDF).** Based on Table 4, the NDF content of forages was not affected by cropping system but influenced by the forage type. NDF content of *C. ternatea* (52.1%-55.7%) obtained in this study were significantly (P<0.05) lower than NDF content of *S. splendida* (68.2%-72.6%) and *C. pubescens* (66%-67.8%). In general, NDF content of *C. pubescens* was lower than of *S. splendida*. These results indicated that the NDF content of legumes was generally lower than those of grass. This was attributed to the different characteristics

Table 5. Crude protein (CP) production per plot of forages based on cropping system

Cropping system	Plot	CP production	
		(g/plot/harvest)	(kg/plot/yr)
Monoculture	<i>S. splendida</i> (SS)	207	1.243
	<i>C. pubescens</i> (CP)	213	1.280
	<i>C. ternatea</i> (CT)	261	1.567
Mixture	SS/CP	223	1.339
	SS/CT	157	0.943

of the cell wall (Paulson *et al.*, 2008). Legumes have particularly large amounts of pectin in primary walls, resulting in more pectin in legume forages than grasses in both leaves and stems (Jung & Casler, 2006; Paulson *et al.*, 2008). Most tissue types in grasses lignify as they mature whereas most legume tissues do not lignify (Paulson *et al.*, 2008). Based on the value of NDF, the *C. ternatea* intake was higher than *S. splendida* and *C. pubescens* when fed to ruminants. NDF value reflects the total fraction of fibre (cellulose, hemicellulose, and lignin). Thus, the NDF content will be negatively correlated to the intake (Mertens, 2009). When NDF content of forage is high, it causes a decrease in intake (Milic *et al.*, 2011). The average NDF content of *S. splendida*, *C. Pubescens*, and *C. ternatea* found in this study were generally higher than the findings of other researchers. Heuzé & Tran (2012) reported that NDF content of *S. splendida* was 69.5%. Evitayani *et al.* (2004) reported that NDF content of *C. pubescens* that grows in North Sumatra was 51.1%. Nasrullah *et al.* (2003) found that NDF content of *C. ternatea* which grows naturally in South Sulawesi was 42.30%. Variation in NDF content is caused by genotype and environmental factors (Adesogan *et al.*, 2002; Nasrullah *et al.*, 2003; Evitayani *et al.*, 2004).

**Acid detergent fibre content.** This study found that the ADF content of forages was not affected by cropping system but influenced by the forage type (Table 4). This was probably due to the relative stability of the cellulose content of grasses and legumes (Weiss *et al.*, 2002). The

results showed that the ADF content of *C. pubescens* was significantly ( $P < 0.05$ ) higher than *C. ternatea* and *S. splendida*. In relation to the digestibility *C. pubescens* have lower digestibility than *C. ternatea* and *S. splendida* (Albayrak *et al.*, 2011). The results also illustrated that the ADF content of legumes was not always lower than the grass. Grasses and legumes are likely to have the same ADF value, but NDF content of grass generally always substantially higher than the legume (Weiss *et al.*, 2002).

## CONCLUSION

Mixed cropping in peatland (type sapric) does not influence productivity and nutritive value of *S. splendida*, *C. Pubescens*, and *C. ternatea*. *S. splendida* is more effective in exploiting environmental resources when intercropped with *C. pubescens* compared *C. ternatea* on peatland. Based on value of K, CR, and A, *S. splendida* is more competitive and dominant than *C. pubescens* and *C. ternatea* on peatland.

## REFERENCES

- Adesogan, A. T., L. E. Sollenberger, Y. C. Newman, & J. M. B. Vendramini. 2002. Factors Affecting Forage Quality. Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. [edis.ifas.ufl.edu/pdffiles/AG/AG16100.pdf](http://edis.ifas.ufl.edu/pdffiles/AG/AG16100.pdf). [1 September 2013].
- Agus, F. & I. G. M. Subiksa. 2008. Lahan Gambut: Potensi untuk Pertanian dan Aspek Lingkungan. Balai Penelitian Tanah. Badan Penelitian dan Pengembangan Pertanian. Bogor. Indonesia.
- Ajayi, F. T., O. J. Babayemi, & A. A. Taiwo. 2007. Effect of *Stylosanthes guianensis* and *Aeschynomene histrix* on the yield, proximate composition and in-situ dry matter and crude protein degradation of *Panicum maximum* (Ntchisi). *Lives. Res. for Rur. Dev.* 19 (3). [www.lrrd.org/lrrd19/3/ajay19032.htm](http://www.lrrd.org/lrrd19/3/ajay19032.htm). [24 July 2013].
- Albayrak S., M. Turk, O. Yuksel, & M. Yilmaz. 2011. Forage yield and the quality of perennial legume-grass mixtures under rainfed conditions. *Not. Bot. Hort. Agrobot. Cluj.* 39:114-118.
- AOAC. 2005. Official Methods of Analysis. AOAC International. 18<sup>th</sup> ed. Assoc. Off. Anal. Chem, Arlington, USA.
- Banik, P., A. Midya, B. K. Sarkar, & S. S. Ghose. 2006. Wheat and chickpea intercropping systems in an additive series experiment: Advantages and weed smothering. *Eur. J. Agron.* 24: 325-332. <http://dx.doi.org/10.1016/j.eja.2005.10.010>
- Chen, C., M. Westcott, K. Neill, D. Wichman, & M. Knox. 2004. Row configuration and nitrogen application for barley-pea intercropping in Montana. *Agron. J.* 96: 1730-1738. <http://dx.doi.org/10.2134/agronj2004.1730>
- Dhalika, T., Mansyur, H. K. Mustafa, & H. Supratman. 2006. Proportion of African star grass and sentro in mix pasture system on production and quality of forages. *J. Ilmu Ternak.* 6: 163-168.
- Dhima, K. V., A. A. Lithourgidis, I. B. Vasilakoglou, & C. A. Dordas. 2007. Competition indices of common vetch and cereal intercrops in two seeding ratio. *Field Crop Res.* 100: 249-256. <http://dx.doi.org/10.1016/j.fcr.2006.07.008>
- Eskandari, H. 2012. Yield and quality of forage produced in intercropping of maize (*Zea mays*) with cowpea (*Vigna sinensis*) and mungbean (*Vigna radiate*) as double cropped. *J. Basic. Appl. Sci. Res.* 2: 93-97.
- Evitayani, L. Warly, A. Fariani, T. Ichinohe, & T. Fujihara. 2004. Study on nutritive value of tropical forages in North Sumatra, Indonesia. *Asian-Aust. J. Anim. Sci.* 17: 1518-1523.
- Forrester, D. I., J. Bauhus, A. L. Cowie, & J. K. Vanclay. 2006. Mixed-species plantations of *Eucalyptus* with nitrogen-fixing trees: A review. *Forest Ecol. Manag.* 233: 211-230. <http://dx.doi.org/10.1016/j.foreco.2006.05.012>
- Heinritz, S. N., S. Hoedtke, S. D. Martens, M. Peters, & A. Zeyner. 2012. Evaluation of ten tropical legume forages for their potential as pig feed supplement. *Lives. Res. for Rur. Dev.* 24 (1). <http://www.lrrd.org/lrrd24/1/cont2401.htm>. [28 August 2013].
- Heuzé, V. & G. Tran. 2012. Giant setaria (*Setaria sphacelata var. splendida*). *Feedipedia.org*. <http://www.feedipedia.org/node/380>. [30 Juli 2013].
- Jayanegara, A. & A. Sofyan. 2008. Penentuan aktivitas biologis tanin beberapa hijauan secara *in vitro* menggunakan hohenheim gas test dengan polietilen glikol sebagai determinan. *Med. Pet.* 31: 44-52.
- Jung, H. G. & M. D. Casler. 2006. Maize stem tissues: Cell wall concentration and composition during development. *Crop. Sci.* 46:1793-1800. <http://dx.doi.org/10.2135/cropsci2005.02-0085>
- Karadag, Y. & U. Buyukburc. 2004. Forage qualities, forage yields and seed yields of some legume-triticale mixtures under rainfed conditions. *Acta Agri. Scan. B-S P* 54: 140-148.
- Mahala, A.G., S.O. Amasiab, Monera, A. Yousif, & Elsadig. 2012. Effect of plant age on DM yield and nutritive value of some leguminous plants (*Cyamopsis tetragonoloba*, *Lablab purpureus* and *Clitoria ternatea*). *Int. Res. J. Agric. Sci.* 2: 502-508.
- Mahapatra, S. C. 2011. Study of grass-legume intercropping system in term of competition indices and monetary advantage index under acid lateritic soil of India. *Am. J. Exp. Agric.* 1: 1-6.
- Martens, D., S. D. T. T. Tiemann, J. Bindelle, M. Peters, & C. E. Lascano. 2012. Alternative plant protein sources for pigs and chickens in the tropics—nutritional value and constraints: A review. *J. Agr. Rural Develop. Trop. Subtrop.* 113: 101-123.
- Marty, C., A. Pornon, N. Escaravage, P. Winterton, & T. Lamaze. 2009. Complex interactions between a legume and two grasses a subalpine meadow. *Am. J. Bot.* 96: 1814-1820. <http://dx.doi.org/10.3732/ajb.0800405>
- Mertens, D. R. 2009. Impact of NDF content and digestibility on dairy cow performance. *WCDS Advance in Dairy Technology.* 21: 191-201.
- Milić, D., Đ. Karagić, S. Vasiljević, A. Mikić, B. Mijić, & S. Katić. 2011. Leaf and stem chemical composition of devergent alfalfa cultivars. *Biotechnology in Animal Husbandry* 27: 1505-1511. <http://dx.doi.org/10.2298/BAH1104505M>
- Mucheru-Muna, M., P. Pypers, D. Mugendi, J. Kung'u, J. Mugwe, R. Merckx, & B. Vanlauwe. 2010. A staggered maize-legume intercrop arrangement robustly increases crop yields and economic returns in the highlands of Central Kenya. *Field Crops Res.* 115: 132-139. <http://dx.doi.org/10.1016/j.fcr.2009.10.013>
- Mutumura, M. & T. M. Everson. 2012. On-farm evaluation of improved *Brachiaria* grass in low rainfall and aluminium toxicity prone areas of Rwanda. *Int. J. Biodivers. Conserv.* 4: 137-154.
- Nasrullah, M. Niimi, R. Akashi, & O. Kawamura. 2003. Nutritive evaluation of forage plants grown in South Sulawesi, Indonesia. *Asian-Aust. J. Anim. Sci.* 16: 693-701.
- Njad, A. K., S. Mohammadi, N. Khaliliaqdam, M. P. Yousef, & N. J. Nejad. 2013. Barley-clover intercropping: advantages and competition indices. *Adv. Crop. Sci.* 13: 497-505.
- Noula, F. S., O. O. Akinbamijo, O. B. Smith, & V. S. Pandey.

2004. Horticultural residues as ruminant feed in per-urban area of the Gambia. *Livestock Research for Rural Development* 16 (6). <http://www.cipav.org.co/lrrd/lrrd16/6/noua16037.htm>. [20 July 2013].
- Omole, A. J., A. M. Ogungbesan, J. A. Fayenuwo, & Y. A. Popoola.** 2011. Comparative utilization of *Centrosema pubescens* or *Mucuna purensis* as a substitute for pawpaw leaf by growing snails. *Lives. Res. for Rur. Dev.* 23 (6). <http://www.lrrd.org/lrrd23/3/omol23063.htm>. [18 August 2013].
- Oseni, T. O.** 2010. Evaluation of sorghum-cowpea intercrop productivity in savanna agro-ecology using competition indices. *J. Agric. Sci.* 2: 229-234.
- Paulson, J., H. Jung, M. Raeth-Knight, & J. Linn.** 2008. Grass vs Legume Forage for Dairy Cattle. University of Minnesota Extension. <http://www.extension.umn.edu> [30 Juli 2013].
- Peyraud, J. L., A. Le Gall, & A. Lüscher.** 2009. Potential food production from forage legume-based-systems in Europe: An overview. *Irish J. Agric. Food Res.* 48: 115-135.
- Pozdissek, J., B. Henriksen, A. Ponizil, & A. K. Loes.** 2011. Utilizing legume-cereal intercropping for increasing self-sufficiency on organic farms in feed for monogastric animal. *Agron. Res.* 9: 343-356.
- Pirhofer-Walzl, K., J. Rasmussen, H. Høgh-Jensen, J. Eriksen, K. Søegaard, & J. Rasmussen.** 2012. Nitrogen transfer from forage legumes to nine neighbouring plants in a multi-species grassland. *Plant Soil.* 350:71-84. <http://dx.doi.org/10.1007/s11104-011-0882-z>
- Refliaty, Y. Farni, & S. Intan.** 2009. Pengaruh leguminosa cover crop (LCC) terhadap sifat fisik *ultisol* bekas alang-alang dan hasil jagung. *J. Agronomi* 13: 51-56.
- Sabrina D.T., M. M. Hanafi, A.W. Gandahi, M. T. M. Mo-hamed, & N. A. A. Aziz.** 2013. Effect of mixed organic-inorganic fertilizer on growth and phosphorus uptake of setaria grass (*Setaria splendida*). *Aust. J. Crop Sci.* 7:75-83.
- Thorsted, M. D., J. Weiner, & J. E. Olesen.** 2006. Above and below-ground competition between intercropped winter wheat *Triticum aestivum* and white clover *Trifolium repens*. *J. Appl. Ecol.* 43: 237-245. <http://dx.doi.org/10.1111/j.1365-2664.2006.01131.x>
- Van Soest, P. J., J. B. Robertson, & B. A. Lewis.** 1991. Methods for dietary fiber, neutral detergent fiber and non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74: 3583-3593. [http://dx.doi.org/10.3168/jds.S0022-0302\(91\)78551-2](http://dx.doi.org/10.3168/jds.S0022-0302(91)78551-2)
- Vasileva, V. & E. Vasilev.** 2012. Study on productivity of some legume crops in pure cultivation and mixtures. *Agric. Conspec. Sci.* 77: 91-94.
- Weiss, W. P., M. L. Eastridge, & J. F. Underwood.** 2002. Forages for Dairy Cattle. Ohio State University Extension. <http://ohioline.osu.edu/as-fact/0002.html>. [19 August 2013].
- Yilmaz, S., M. Atak, & M. Erayman.** 2008. Identification of advantage of maize-legume intercropping over solitary cropping through competition indices in the East Mediterranean Region. *Turk. J. Agric. For.* 32: 111-119.
- Zhang, F. & L. Li.** 2003. Using competitive and facilitative interactions in intercropping system enhances crop productivity and nutrient-use efficiency. *Plant Soil* 248: 305-312. <http://dx.doi.org/10.1023/A:1022352229863>