

## The Prevalence and Risk Factors of GERD among Indonesian Medical Doctors

Ari F Syam<sup>1\*</sup>, Florentina CP Hapsari<sup>2</sup>, Dadang Makmun<sup>1</sup>

1. Division of Gastroenterology, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia, Jakarta 10310, Indonesia

2. Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia, Jakarta 10310, Indonesia

\*E-mail: ari\_syam@hotmail.com

---

### Abstract

**Background:** Based on our knowledge, the study of gastrointestinal reflux disease (GERD) among certain profession has never been conducted. The aim of this study is to determine the prevalence and risk factors of GERD among Indonesian doctors. **Methods:** A consecutive study involving 515 doctors was conducted in October 2015. The GerdQ score was used to the diagnosis of GERD and determined its impact on daily life. All possible risk factors were also analysed. **Results:** A total of 515 subjects completed the questionnaire. The mean age of them was  $41.37 \pm 11.92$  years old. Fifty-five percent of them were male and 60.6% general practitioners. The prevalence of GERD was 27.4% of which 21.0% was had GERD with low impact on daily life, and 6.4% was GERD with high impact on daily life. The statistically significant risk factors of GERD was found in age  $>50$  y.o ( $p = 0.002$ ; OR = 2.054), BMI  $>30$  kg/m<sup>2</sup> ( $p = 0.016$ ; OR = 2.53), and smokers ( $p = 0.031$ ; OR = 1.982). Sex and education level were not found significant statistically as the risk factors of GERD. **Conclusions:** The prevalence of GERD among Indonesian physician was 27.4%. We found that age over 50 y.o, obesity and smoking habit were the risk factors of GERD in Indonesian doctors.

**Keywords:** GERD, physicians, prevalence, risk factors

---

### Introduction

Gastroesophageal reflux disease (GERD) is one of the common gastrointestinal problems worldwide with a range of GERD prevalence was 18.1-27.8% in North America, 23.0% in South America, 8.8-25.9% in Europe, 11.6% in Australia and 8.7-33.1% in the Middle East. Lower prevalence was found in East Asia region, ranging from 2.5 to 7.8%.<sup>1</sup> Several studies in Iran population reported the prevalence of GERD was 6.3-18.3%. Also, a study in Pakistan showed higher prevalence, 24.0%.<sup>2,3</sup> Even the prevalence in East Asia region was relatively low, a hospital-based study in Indonesia found the prevalence of GERD among dyspeptic patient who underwent endoscopic procedure was 32.4%, higher.<sup>4</sup> Time trend studies have also reported the increase of GERD prevalence.<sup>2</sup>

GERD is known as a condition caused by abnormal reflux of gastric contents into the oesophagus so that the patients may complain heartburn, a burning sensation in the middle of the chest, and regurgitation, a moving upward sensation of stomach content. The inappropriate lower oesophageal sphincter closure categorised as the cause of gastric contents reflux.<sup>5,6</sup> Besides the gastrointestinal symptoms, GERD can also manifest in vary analysed oesophageal symptoms, such as dysphasia, chronic cough, laryngitis, and asthma-like-symptoms.<sup>7</sup>

Eighty percent of GERD symptoms may be persistent, and it implicates to increase the risk of several complications such as oesophageal ulcer, oesophageal stricture, Barrett's oesophagus, and adenocarcinoma.<sup>8,9</sup> Moreover, the persistent symptoms can reduce the work productivity and quality of life.<sup>10</sup>

The GERD diagnosis can be established in the setting of typical symptoms of heartburn and regurgitation.<sup>11</sup> Many studies develop easier tools to help the physicians in the diagnosing patient with GERD symptoms. The most recent instrument was GERD Questionnaire (GerdQ) which was created from three different validated questionnaires evaluated in the DIAMOND study.<sup>12,13</sup> GerdQ has been used worldwide as an easy way to predict the probability of GERD in patients with gastrointestinal symptoms. Validation study of GERDQ in the Indonesian language has been done before which showed that Indonesian version of GERDQ is reliable to be used to diagnose GERD based on the reported symptoms with the same cut-off point.<sup>6</sup>

Medical doctor has been known as one of high work-hour professions, with a range of 51 to 67 hours per week from various studies.<sup>14-16</sup> The general practitioner, internal medicine resident, and internal medicine specialist and subspecialist work for long hours in a quite stressful environment and spend most of their time

in hospitals.<sup>16</sup> That kind of physician could be highly potential burnout. A Recent study showed that stress and shift duty influenced the amount of daily intake among health provider. Moreover, they were likely to eat fast food and snack, while fruits and vegetables were the less likely to be eaten under stress.<sup>17</sup> A study in Bahrain also revealed that 39% of the physician being overweight and 33% obese.<sup>18</sup> Meanwhile, previous studies have been proved that obesity was related to GERD.<sup>19</sup> Besides, low prevalence of *Helicobacter pylori* infection can influence the prevalence rate of GERD. In the previous study, several ethnics group found to have a lower risk of *H. pylori* infection.<sup>20</sup> Therefore, ethnicity and *H. pylori* infection can also be the factors that contribute the prevalence of GERD.

A Recent study among the certain population in Indonesia has never been conducted. This study was designed to determine the prevalence of GERD among Indonesian medical doctors using GerdQ as the diagnostic tool. We also analysed the demographic characteristics and possible risk factors of GERD.

## Methods

**Study design.** The cross-sectional study involving 515 medical doctors was conducted in October 2015. The subjects fulfilled questionnaire containing demographic questions (i.e. sex, age, weight, height, ethnic, and educational level), validated Indonesian language GERD Questionnaire, and several questions about smoking habit and endoscopy history. People who fulfilled all questions would proceed as research subjects. We ruled out all incomplete questionnaire. All information about the subjects was confidential. The GerdQ score was used to diagnose GERD and determined its impact on daily life. All possible risk factors were also analysed to find its association with GERD. The protocol has been approved by the committee of the Medical Research Ethics of the Faculty of Medicine, Universitas Indonesia.

**The GERD Questionnaire (GerdQ).** The GerdQ is a self-assessment questionnaire that evaluates the presence of cardinal symptoms of GERD, heartburn and regurgitation, and its impact on daily life.<sup>12</sup> This questionnaire has been developed from three established GERD questionnaire instruments: the Reflux Disease Questionnaire, Gastrointestinal Symptom Rating Scale, and the Gastrointestinal Symptom Scale through Diamond Study.<sup>6,21</sup> The GERDQ is a simple communication tool which has been developed for physicians to identify patients with GERD.

The GerdQ consists of six questions about related symptoms in the last seven days as follows: heartburn, regurgitation, upper stomach pain, nausea, sleep difficulties due to heartburn and regurgitation, and

medication history to relieve the GERD symptoms. The first and last two questions are positive predictors of GERD, where a more frequent of symptoms, the higher the score. Meanwhile, the questions about nausea and abdominal pain are the negative predictors of GERD which is the score is in reverse. The last two questions (sleep difficulty and medication history) assess the influence of symptoms on daily life. GerdQ score of more than 8 is determined as high probability of GERD. If the total score for the last two questions (question number 5 and 6) is more than 2 (two), it is determined as GERD with high impact on daily life.

**Data analysis.** The survey results were input and analysed using SPSS 22.0 for windows. Descriptive statistics were presented as mean, the standard of deviation (SD), and proportion (%). The BMI was classified into two groups:  $<30 \text{ kg/m}^2$  and  $\geq 30.0 \text{ kg/m}^2$ .<sup>22</sup> Total GerdQ score were calculated by summing all of the scores for each question in the questionnaire, then classified as the mention above. Independent factors were analysed using multivariate analysis with backwards logistic regression methods. Statistically significant was determined by *p* value under 0.05 with 95% confidential interval (CI).

## Results

**Basic characteristics.** The demographic distribution was presented in Table 1. As many as 515 subjects were recruited to this study. The mean of age was  $41.37 \pm 11.92$  years old, 55.7% was male, and the mean of body mass index (BMI) was  $23.62 \pm 4.41 \text{ kg/m}^2$ . General practitioner/bachelor graduates were found in 60.6% of subjects and the rest of them were specialist and subspecialist or PhD graduates.

**GerdQ results.** The prevalence of GERD in Indonesian medical doctors was 27.4% of which 21.0% was GERD with low impact on daily life, and 6.4% was GERD with high impact on daily life (Table 2). 72.6% of subjects had a GerdQ score below eight so that they oesophageal as low probability of GERD.

The heartburn symptoms appear more frequent than regurgitation. Heartburn symptoms found in 17.4% of subjects at least twice a week, meanwhile regurgitation was found in 15% of subjects for the same frequency. Sleep disturbance was experienced in 27.3% of subjects at least once a week. The heartburn and regurgitation symptoms implicated 35% of subjects took medication to relieve the symptoms at least once a week. Table 3 shows the proportion for each GerdQ question in all subjects.

**Risk factors of GERD.** Our study was analysed for potential risk factors of GERD (Table 4). Previous studies reported age, sex, education level, high body mass index, and smoker possibly increased the risk of

**Table 1. Basic Characteristics of Subjects**

Characteristics	Frequency (n)	Percent (%)
Mean of age (y.o)	41.37+11.92	
Mean of weight (kg)	64.87+13.50	
Mean of height (cm)	165.6+8.56	
Mean of body mass index (kg/m <sup>2</sup> )	23.62+4.41	
<b>Age Groups</b>		
21-30 y.o	106	20.6
31-40 y.o	180	35.0
41-50 y.o	120	23.3
>50 y.o	109	21.2
<b>Sex</b>		
Male	287	55.7
Female	228	44.3
<b>BMI Groups</b>		
<18.5 kg/m <sup>2</sup>	55	10.7
18.5-22.9 kg/m <sup>2</sup>	171	33.2
23.0-24.9 kg/m <sup>2</sup>	106	20.6
25.0-29.9 kg/m <sup>2</sup>	152	29.5
>30kg/m <sup>2</sup>	31	6.0
<b>Ethnics</b>		
Javanese	188	36.5
Sundanese	90	17.5
Batak	41	8.0
Minang	34	6.6
Chinese	33	6.4
Betawi	33	6.4
Bugis	16	3.1
Minahasa	14	2.7
Aceh	12	2.3
Palembang	11	2.1
Others	43	8.3
<b>Educational Status</b>		
Bachelor degree	312	60.6
Master degree/ specialist	194	37.7
Doctoral Degree/ Subspecialist	9	1.7
<b>Smoking habit</b>		
Non-smoker	467	90.7
Smoker	29	5.6
Ex-smoker	19	3.7

GERD. Then multivariate analysis was performed using backwards logistic regression analysis that can show only the statistic significant probability among the independent factors. Table 5 shows the result of

multivariate analysis of GERD risk factors. The statistically significant risk factors of GERD was found in age over 50 years old ( $p = 0.002$ ; OR 2.054; 95% CI 1.299-3.247). BMI over 30 kg/m<sup>2</sup> ( $p = 0.016$ ; OR 2.53; 95% CI 1.191-5.375), and smokers ( $p = 0.031$ ; OR 1.982; 95% CI 1.063-3.694) also was showed statistical significance as the risk factors of GERD.

**Table 2. Prevalence of GERD in Each Group of Age**

Age	Low probability of GERD (n,%)	GERD with low impact on daily life (n,%)	GERD with high impact on daily life (n,%)
21-30 years old	77 (72.0)	24 (22.4)	6 (5.6)
31-40 years old	138 (76.7)	32 (17.7)	10 (5.6)
41-50 years old	92 (76.7)	21 (17.5)	7 (5.8)
>50 years old	67 (62.0)	31 (28.7)	10 (9.3)
Total	374 (72.6)	108 (21.0)	33 (6.4)

**Table 3. Frequency of Symptoms during Previous Week in GERD Patients**

Questions	Frequency of symptoms (n,%)			
	0 days	1 day	2-3 days	4-7 days
How often did you have a burning feeling behind your breastbone (heartburn)?	310 (60.2)	115 (22.3)	63 (12.2)	27 (5.2)
How often did you have stomach contents moving upwards to your throat or mouth (regurgitation)?	275 (53.4)	163 (31.7)	55 (10.7)	22 (4.3)
How often did you have a pain in the centre of the upper abdomen?	21 (4.1)	60 (11.7)	156 (30.3)	278 (54.0)
How often did you have nausea?	30 (5.8)	71 (13.8)	145 (28.2)	269 (52.2)
How often did you have difficulty sleeping because of your heartburn and/or regurgitation?	374 (72.6)	89 (17.3)	31 (6.0)	21 (4.1)
How often did you take additional medication for heartburn and/or regurgitation, other than what we physician told to you take?	335 (65.0)	100 (19.4)	56 (10.9)	24 (4.7)

**Table 4. The Association of GERD and Several Possible Risk Factors**

Characteristics	GERD (n,%)	Non-GERD (n,%)	p (CI 95%)	OR (CI 95%)
<b>Sex</b>				
Male	88 (30.7)	199(69.3)	0.061	1.460 (0.982-2.171)
Female	52 (23.2)	175(76.8)		
<b>Age</b>				
≤50 y.o	100 (24.6)	307 (75.4)	0.008	2.011 (1.195-3.382)
>50 y.o	42 (38.5)	67(61.5)		
<b>Education Status</b>				
Bachelor/Doctor	86 (27.6)	226(72.4)	0.852	1.039 (0.694-1.555)
Master/Specialist	52 (26.8)	142(73.2)		
PhD/Subspecialist	3 (33.3)	6(66.7)		
<b>Body Mass Index</b>				
<30 kg/m <sup>2</sup>	127 (26.2)	357 (73.8)	0.035	2.315 (1.109-4.832)
≥30 kg/m <sup>2</sup>	14 (45.2)	17 (54.8)		
<b>Smoking Habit</b>				
Smoker	20 (41.7)	28 (58.3)	0.026	2.043 (1.110-3.759)
Non Smoker	121 (25.9)	346(74.1)		

**Table 5. Result of Multivariate Analysis of GERD Risk Factors**

Variables	p	OR (CI 95%)
Age >50 y.o	0.002	2.054 (1.299-3.247)
BMI >0 kg/m <sup>2</sup>	0.016	2.53 (1.191-5.375)
Smoker	0.031	1.982 (1.063-3.694)

## Discussion

The prevalence of GERD among medical doctors using GerdQ tools 27.4% (141 subjects). It consists of 21% (108 subjects) of GERD with low impact on daily life and 6.4% of GERD with high impact on daily life. Age > 50 years old, obesity and smoking habit were found statistically significant as the risk factors of GERD among medical doctors in Indonesia.

Our findings were similar to the prevalence in North America but higher than the findings in South America, Europe, Australia, and East Asia.<sup>1</sup> Similar study in Pakistan, Iran, and Korea reported GERD in the population was 6.3-18.3%, 24.0%, and 25% respectively.<sup>2,3,23</sup> Lower prevalence was also found in studies in India (7.6%), Japan (1.6%-16.3%), and China (3.1%).<sup>1,24,25</sup> A comparative study conducted by Mahadeva *et al.* reported the British have more prevalent GERD symptoms than South-east Asian population. They predicted the difference is probably due to both ethnic differences and environmental influences.<sup>26</sup> Therefore, the previous study conducted in Indonesia reported that the prevalence among dyspeptic patients was 32.4%, higher than our findings.<sup>4</sup> It might

be due to the study was conducted only among dyspeptic patients while our study was conducted in the general population. A hospital-based study in Jakarta revealed an increasing trend of GERD prevalence from 6% to 26% within 5 years.<sup>3</sup>

In our study, heartburn and regurgitation symptoms were presented in 39.8% and 46.6% of subjects respectively in vary frequency (from once a week to every day). This result was consistent with the previous study in Indonesia.<sup>3</sup> Because of these symptoms, 4.1% of the subjects had sleep disturbance every day, and 35% need medication to relieve the symptoms at least once a week.

Many factors were expected contribute to these findings, such as of obesity, alcohol consumption, and smoking habit. Medical doctors as one of high workload and high work-hour profession may potentially influence their eating habit. Though the GERD event among medical doctor has not been clear, some studies reported unhealthy diet, shift-work, and obesity among health profession that might be contributing to the GERD event.<sup>17,27</sup>

More male (55.7%) than female (44.3%) were included in this study. Most of the patients with GERD were male (88/62.9%), but it was not significantly difference ( $p = 0.061$ ; OR = 1.460; 95% CI = 0.982-2.171). The association between GERD and gender has not shown consistency. Similar to our study, the study in Chinese population did not show any significant difference in gender related to GERD.<sup>28</sup> On the contrary, the studies in Japan and Greece reported that male is more prevalent than female.<sup>29,30</sup>

We found that age over 50 years old was a significant risk factor of GERD among doctors ( $p = 0.002$ ; OR = 2.054; 95% CI = 1.299-3.247). This study revealed that the older the age, the higher the prevalence of GERD with high impact on daily life.<sup>2</sup> The prevalence of GERD with high impact on daily life in a group of age > 50 years old was 1.75 times higher than 21-30 years old group. Further study is needed to elaborate the reason behind this finding.

Obesity in Asian population was defined as BMI over 24.9 kg/m<sup>2</sup> (24.9-29.9 kg/m<sup>2</sup> as obesity grade 1 and over 30.0 kg/m<sup>2</sup> as obesity grade 2).<sup>22</sup> We found that subjects with BMI  $\geq 30$  kg/m<sup>2</sup> have significantly higher risk of GERD ( $p = 0.016$ , OR = 2.53; 95% CI = 1.191-5.375). This finding was similar to a study in the Danish population.<sup>19</sup> The linkage between GERD and obesity is not clearly understood. It was predicted due to the insufficiency of the barrier to gastric reflux and the increasing intra-abdominal pressure. The alteration of bile and pancreatic function in obese patients are correlated with the composition of the refluxate products which are more toxic for esophagus.<sup>31</sup>

In the present study, GERD was significantly more prevalent in smokers ( $p = 0.031$ ; OR = 1.982; 95% CI = 1.063-3.694). Our results were similar to study in Japan, Northern India, and Denmark.<sup>19,32,33</sup> A study in Japan also reported the risk of GERD among smokers is doubled if it overlaps with other functional gastrointestinal syndromes such as functional dyspepsia and irritable bowel syndrome.<sup>32</sup> Furthermore, a study in Japanese men revealed that an increase in some pack-years of cigarette smoking was significantly associated with an increased risk of GERD.<sup>34</sup> Different mechanisms have been suggested to justify the association between smoking and GERD. Cigarette smoking can reduce the lower oesophageal sphincter (LES) pressure and decrease salivary bicarbonate secretion so that the acid clearance is prolonged and the neutralising effect of saliva on intra oesophageal acid is reduced.<sup>35</sup>

## Conclusions

The prevalence of GERD among Indonesian medical doctors was 27.4%. We conclude that age over 50 years old, obesity (BMI > 30 kg/m<sup>2</sup>), and smoking habit were the risk factors of GERD in Indonesian medical doctors. Further study is needed to support these findings and explore other potential risk factors of GERD.

## Conflict of Interest Statement

The Authors declare that there is no conflict of interest.

## Acknowledgements

We thank gastroenterologists at RS Cipto Mangunkusumo and Indonesian Society of Gastroenterology who fully

support to this study. This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## References

1. El-Serag HB, Sweet S, Winchester CC, Dent J. Update on the epidemiology of gastro-oesophageal reflux disease: a systematic review. *Gut*. 2014;63:871-80.
2. Jung H-K. Epidemiology of gastroesophageal reflux disease in Asia: A systematic review. *J Neurogastroenterol Motil*. 2011;17:14-27.
3. Jafri N, Jafri W, Yakooob J, Islam M, Manzoor S, Jalil A, et al. Perception of gastroesophageal reflux disease in urban population in Pakistan. *J Coll Physicians Surg Pak*. 2005;15:532-4.
4. Aulia C. Prevalence of non-erosive reflux disease in Pondok Indah Hospital: A preliminary study. *Acta Med Indones*. 2005;37:79-81.
5. Mansour-Ghanaei F, Joukar F, Atshani SM, Chaghavand S, Souti F. The epidemiology of gastroesophageal reflux disease: A survey on the prevalence and the associated factors in a random sample of the general population in the northern part of Iran. *Int J Mol Epidemiol Genet*. 2013;4:175-82.
6. Simadibrata M, Rani A, Adi P, Djumhana A, Abdullah M. The gastro-oesophageal reflux disease questionnaire using Indonesian language : a language validation survey. *Med J Indones*. 2011;20:4-6.
7. Saritas Yuksel E, Vaezi MF. Extraesophageal manifestations of gastroesophageal reflux disease: cough, asthma, laryngitis, chest pain. *Swiss Med Wkly*. 2012;142:1-8.
8. Nouraie M, Razjouyan H, Assady M, Malekzadeh R, Nasseri-Moghaddam S. Epidemiology of gastroesophageal reflux symptoms in Tehran, Iran: a population-based telephone survey. *Arch Iran Med*. 2007;10:289-94.
9. Thrift AP, Kramer JR, Qureshi Z, Richardson PA, El-Serag HB. Age at onset of GERD symptoms predicts risk of Barrett's esophagus. *Am J Gastroenterol*. 2013;108:915-22.
10. Wahlqvist P, Karlsson M, Johnson D, Carlsson J, Bolge SC, Wallander M a. Relationship between symptom load of gastro-oesophageal reflux disease and health-related quality of life, work productivity, resource utilization and concomitant diseases: Survey of a US cohort. *Aliment Pharmacol Ther*. 2008;27:960-70.
11. Katz PO, Gerson LB, Vela MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. 2013;108:308-28.
12. Jones R, Junghard O, Dent J, Vakil N, Halling K, Wernersson B, et al. Development of the GerdQ, a tool for the diagnosis and management of gastro-oesophageal reflux disease in primary care. *Aliment Pharmacol Ther*. 2009;30:1030-8.
13. Ang TL, Fock KM, Ng TM, Teo EK, Chua TS, Tan J. A comparison of the clinical, demographic and psychiatric profiles among patients with erosive and non-erosive reflux disease in a multi-ethnic Asian country. *World J Gastroenterol*. 2005;11:3558-61.
14. Wen J, Cheng Y, Hu X, Yuan P, Hao T, Shi Y. Workload, burnout, and medical mistakes among physicians in China: A cross-sectional study. *Biosci Trends*. 2016;10:27-33.

15. Staiger DO, Auerbach DI, Buerhaus PI. Trends in the work hours of physicians in the United States. *JAMA*. 2010;303:747-53.
16. Gopal R, Glasheen JJ, Miyoshi TJ, Prochazka AV. Burnout and internal medicine resident work-hour restrictions. *Arch Intern Med*. 2005;165:2595-00.
17. Almajwal AM. Stress, shift duty, and eating behavior among nurses in Central Saudi Arabia. *Saudi Med J*. 2016;37:191-8.
18. Borgan SM, Jassim GA, Marhoon ZA, Ibrahim MH. The lifestyle habits and wellbeing of physicians in Bahrain: A cross-sectional study. *BMC Public Health*. 2015;15:655.
19. Hansen JM, Wildner-Christensen M, Schaffalitzky de Muckadell OB. Gastroesophageal reflux symptoms in a Danish population: A prospective follow-up analysis of symptoms, quality of life, and health-care use. *Am J Gastroenterol*. 2009;104:2394-403.
20. Syam AF, Miftahussurur M, Makmun D, Nusi IA, Zain LH, Zulkhairi, et al. Risk factors and prevalence of *helicobacter pylori* in five largest islands of Indonesia: A preliminary study. *PLoS One*. 2015;10:e0140186.
21. Tielemans MM, Jansen JB, van Oijen MG. Open access capture of patients with gastroesophageal reflux disease using an online patient-reported outcomes instrument. *Interact J Med Res*. 2012;1:e7.
22. World Health Organization. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2015;363:157-63.
23. Hung L-J, Hsu P-I, Yang C-Y, Wang E-M, Lai K-H. Prevalence of gastroesophageal reflux disease in a general population in Taiwan. *J Gastroenterol Hepatol*. 2011;26:1164-8.
24. Bhatia SJ, Reddy DN, Ghoshal UC, Jayanthi V, Abraham P, Choudhuri G, et al. Epidemiology and symptom profile of gastroesophageal reflux in the Indian population: report of the Indian Society of Gastroenterology Task Force. *Indian J Gastroenterol*. 2011;30:118-27.
25. He J, Ma X, Zhao Y, Wang R, Yan X, Yan H, et al. A population-based survey of the epidemiology of symptom-defined gastroesophageal reflux disease: the Systematic Investigation of Gastrointestinal Diseases in China. *BMC Gastroenterol*. 2010;10:94.
26. Mahadeva S, Raman MC, Ford AC, Follows M, Axon AT, Goh KL, et al. Gastro-oesophageal reflux is more prevalent in Western dyspeptics: a prospective comparison of British and South-East Asian patients with dyspepsia. *Aliment Pharmacol Ther*. 2005;21:1483-90.
27. Knutsson A, Bøggild H. Gastrointestinal disorders among shift workers. *Scand J Work Environ Health*. 2010;36:85-95.
28. Lin M, Gerson LB, Lascar R, Davila M, Triadafilopoulos G. Features of gastroesophageal reflux disease in Women. *Am J Gastroenterol*. 2004;99:1442-7.
29. Mishima I, Adachi K, Arima N, Amano K, Takashima T, Moritani M, et al. Prevalence of endoscopically negative and positive gastroesophageal reflux disease in the Japanese. *Scand J Gastroenterol*. 2005;40:1005-9.
30. Pelechas E, Azoicăi D. Preventive Medicine - Laboratory Gastroesophageal Reflux Disease : Epidemiological Data, Symptomatology, and Risk Factors. *Rev Med Chir Soc Med Nat*. 2013;117:183-8.
31. Fornari F, Callegari-Jacques S, Dantas R, Scarsi A, Ruas L, de Barros S. Obese Patients Have Stronger Peristalsis and Increased Acid Exposure in the Esophagus. *Dig Dis Sci*. 2011;56:1420-6.
32. Fujiwara Y, Kubo M, Kohata Y, Machida H, Okazaki H, Yamagami H, et al. Cigarette smoking and its association with overlapping gastroesophageal reflux disease, functional dyspepsia, or irritable bowel syndrome. *Intern Med*. 2011;50:2443-7.
33. Sharma P, Ahuja V, Madan K, Gupta S, Raizada A, Sharma M. Prevalence, severity, and risk factors of symptomatic gastroesophageal reflux disease among employees of a large hospital in Northern India. *Indian J Gastroenterol*. 2011;30:128-34.
34. Watanabe Y, Fujiwara Y, Shiba M, Watanabe T, Tominaga K, Oshitani N, et al. Cigarette smoking and alcohol consumption associated with gastro-oesophageal reflux disease in Japanese Men. *Scand J Gastroenterol*. 2003;38:807-11.
35. Festi D, Scafoli E, Baldi F, Vestito A, Pasgui F, Di Biase AR, et al. Body weight, lifestyle, dietary habits, and gastroesophageal reflux disease. *World J Gastroenterol*. 2009;15:1690-701.