

An updated checklist of the herpetofauna of the Belum-Temengor forest reserves, Hulu Perak, Peninsular Malaysia

Amirah Hurzaid¹, MohdAzmeer Abu Bakar¹, Dionysius Sharma², Nurolhuda Nasir², Reuben Sharma³, Ahmad RidzuanYeop Aznan⁴ and Ibrahim Jaafar¹

¹Biological Sciences Program, School of Distance Education, Universiti Sains Malaysia, 11800 Minden, Penang, Malaysia.

²WWF- Malaysia, 49, Jalan SS 23/15, Taman SEA, 47400 Petaling Jaya, Selangor, Malaysia.

³Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia.

⁴Yayasan EMKAY, Unit 3A-3, Level 4, Jalan PJU 8/5g, Bandar Damansara Perdana, 47820 Petaling Jaya, Selangor, Malaysia.

Corresponding Author: amirahhurzaid@gmail.com

Abstract. A herpetofaunal survey was carried out in Temengor Forest Reserve, Peninsular Malaysia during the second Temengor Scientific Expedition conducted from 1st to 10th October 2012. This study represents the first records of amphibians and reptiles of Sungai Enam Basin and an updated record of Belum-Temengor Forest Reserve. In this survey, a total of 27 species of herpetofauna was recorded comprising 12 species of amphibians from four families and 15 species of reptiles from six families. No new records of frog or lizard species are reported in this study. However, three new records of snakes are reported from the area.

Key words: Herpetofauna, Malaysia, Temengor Forest Reserve.

Introduction

The 2nd Temengor Scientific Expedition to Belum running from 1st to 10th October 2012 was based at Sungai Enam, a small, 4 km long tributary of the Singor River, which flows into the Temengor Reservoir. The vegetation of this area is characteristic of Lowland Dipterocarp Forest. The first herpetofauna survey of Belum-Temengor Forest Reserves was conducted by Kiew et al. (1995) during the Malaysian Nature Society's Heritage and Scientific Expedition to Belum. They found 24 species of amphibians, representing 26% of the amphibian fauna of Peninsular Malaysia. In the same expedition, Diong et al. (1995) reported on the lizards while Lim et al. (1995a, b) reported on the snakes and turtles, respectively.

The second survey conducted opportunistically during a mammal survey of the northern part of the Belum Forest Reserve by Norsham et al. (2000) recorded the presence of 26 species of amphibians and reptiles (nine species of frogs, 10 species of lizards and geckos, one species of varanid and skink, four species of snakes and one species of turtle). Sukumaran (2002) documented 26 species of frogs and seven species of reptiles from five and four families respectively within the Perak Integrated Timber Complex (PITC) located in Temengor Forest Reserve. Two years later, a comprehensive survey by Grismer et al. (2004) was conducted at the PITC. They discovered 32 species of amphibians and 25 species of reptiles representing five and seven families, respectively. In their survey, they added five new records of frogs including an undescribed species of flying frog (*Rhacoporus*), one caecilian, 15 species of lizards and four species of snakes. In this study, we added three new records of snakes; *Typhlopsmuelleri*, *Calamariaschlegeli* and *Rhabdophischrysargos* from two families.

Materials and Methods

Study area

The Sungai Enam base camp (5°30.65'N 101°27.79'E), with an elevation of 450m a.s.l., is located at Sungai Enam, a small, 4 km long tributary of the Singor River, which then flows into Temenggor Reservoir (Fig. 1). The base camp is about 500 m from the river mouth; about 200 m of the lower part of the river, the riverbed is made up of sandy and silty soil, whereas about 3.8 km upstream is of rocky terrain with gravel, cobblestones, boulders and occasional large boulders (Fig. 2). Bedrock is also present, forming some rock pools and some small riffles. The banks of the river were somewhat overgrown with lowland dipterocarp vegetation with trees, shrubs, saplings, ferns and lianas overhanging the river.

Herpetofaunal survey

The survey was conducted over a period of four days from 2-5 October 2013, accumulating to 24 active survey hours (hours surveyed actively by 4 to 7 number of people). Areas around the base camp, the surrounding forest, established trails, river banks and streams were actively searched for herpetofauna. Amphibians were captured mainly at night by hand or with fishnets by field parties working from 2000 hrs to 2300 hrs with the help of handheld battery powered torchlights. Surveys for reptiles were a combination of active searches and chance observations. Lizards were captured by hand or by sweep nets, while smaller snakes and turtles were caught by hand, nets and poles. The specimens collected were placed in separate plastic bags and labeled for subsequent processing at the base camp. All captured specimen were photographed and identified by referring to Berry (1975), Ibrahim et al. (2008), Cox et al. (1998) and Das (2010), and the amphibians names were updated according to Frost et al. (2011). All amphibian specimens were injected with benzocaine solution into its dorsal lymph sac to humanely kill them before dissection (Norhayati Ahmad, pers. comm).




Figure 1: Map showing the location of study area (modified from Rainwalker Ecosystems, 2012).

Specimens were then weighed to the nearest 0.1g using an AND Electronic Precision Model EK- 1200A (1200g capacity). Basic morphometric data; snout- vent length (SVL), mouth width (MW), tibia length (TL) and tympanum diameter (TD) were measured to the nearest 0.01mm using dial calipers (\pm 0.05mm). Voucher specimens of species collected were preserved in 10% formalin and later stored in 70% alcohol and housed at The Amphibian and Reptile Collections University Sains Malaysia (ARC, USM). For the reptiles, the voucher specimens were housed at the Faculty of Veterinary Medicine, Universiti Putra Malaysia for future reference.

Results and Discussion

A total of 27 species of amphibians and reptiles were identified. These comprised twelve species of frogs and toads, belonging to four families; eight species of lizards and geckos from two families; six species of snakes from four families; and a species of softshell turtle of one family. The updated checklists of amphibians (Table 1) and reptiles (Table 2) are presented below with annotations for each species recorded.

Table 1: Amphibians of Belum-Temengor Forest Reserves.

| Species | Kiew et al. 1995 | Norsham et al. 2000 | Sukumaran 2002 | Grismer et al. 2004 | This study 2012 |
|-------------------------------------|------------------------|---------------------------|-------------------|---------------------------|-----------------------|
| AMPHIBIANS (ANURANS) | | | | | |
| Bufonidae | | | | | |
| 1 <i>Phrynoideasaspera</i> | X | X | X | X | X |
| 2 <i>Ingerophrynusparvus</i> | X | X | X | X | X |
| 3 <i>Leptophryneborbonica</i> | | | X | X | X |
| 4 <i>Pedostibesoshii</i> | | X | X | X | |
| Dicroglossidae | | | | | |
| 5 <i>Fejervaryaaff. Limnocharis</i> | X | X | X | X | X |
| 6 <i>Limnonectesblythii</i> | X | X | X | X | X |
| 7 <i>Limnonecteskuhlii</i> | X | | X | X | |
| 8 <i>Limnonecteslaticeps</i> | X | | X | X | X |
| 9 <i>Limnonectesparamacrodon</i> | | X | | | |
| 10 <i>Limnonectesplicatellus</i> | | | X | X | |
| 11 <i>Occidozygalaavis</i> | X | | | | |
| 12 <i>Limnonecteshascheanus</i> | | | | X | |
| Megophryidae | | | | | |
| 13 <i>Leptobrachiumhendricksoni</i> | X | | | | |
| 14 <i>Leptolalaxheteropus</i> | X | | X | X | X |
| 15 <i>Megophrysnasuta</i> | | | X | X | X |
| 16 <i>Megophrysaceras</i> | | | X | X | |
| Microhylidae | | | | | |
| 17 <i>Kalophrynuspleurostigma</i> | X | | | | |

| | | | | | | | |
|--------------------------------|------------------------------------|---|---|--|---|--|---|
| 18 | <i>Microhylaberdmorei</i> | | | | | | X |
| 19 | <i>Microhylabutleri</i> | | | | X | | X |
| 20 | <i>Microhylaheymonsi</i> | X | | | X | | X |
| Ranidae | | | | | | | |
| 21 | <i>Amolopslarutensis</i> | X | X | | X | | X |
| 22 | <i>Hylaranalaterimaculata</i> | | | | X | | |
| 23 | <i>Hylaranaerythraea</i> | X | X | | | | X |
| 24 | <i>Hylaranaglandulosa</i> | X | | | X | | X |
| 25 | <i>Hylaranahosii</i> | X | X | | X | | X |
| 26 | <i>Hylarananicobariensis</i> | | | | | | X |
| 27 | <i>Hylarananigrovittata</i> | X | | | X | | X |
| 28 | <i>Hylaranalabialis (raniceps)</i> | X | | | X | | X |
| 29 | <i>Hylaranasignata</i> | X | | | X | | X |
| Rhacoporidae | | | | | | | |
| 30 | <i>Nyctixaluspictus</i> | | | | X | | X |
| 31 | <i>Philautusparvulus</i> | | | | X | | X |
| 32 | <i>Philautuspetersi</i> | X | | | | | |
| 33 | <i>Polypedatesleucomystax</i> | | | | X | | X |
| 34 | <i>Polypedatesmacrotis</i> | X | | | | | |
| 35 | <i>Rhacoporusnigropalmatus</i> | | | | X | | |
| 36 | <i>Rhacoporusprominanus</i> | X | | | X | | X |
| 37 | <i>Rhacoporus sp.</i> | | | | | | X |
| 38 | <i>Theلودermahorridum</i> | | | | | | X |
| AMPHIBIANS (CAECILIANS) | | | | | | | |
| Ichthyophidae | | | | | | | |
| 39 | <i>Ichthyophis sp.</i> | | | | | | X |

Table 2: Reptiles of Belum- Temengor Forest Reserves.

| Species | Diong et al. 1995 | Lim et al. 1995a | Lim et al. 1995b | Norsham et al. 2000 | Sukumaran 2002 | Grismer et al. 2004 | This study 2012 |
|---------------------------|------------------------------|------------------------|------------------------|---------------------------|-------------------|---------------------------|-----------------------|
| REPTILES (LIZARDS) | | | | | | | |
| Agamidae | | | | | | | |
| 40 | <i>Aphanotisfusca</i> | X | | | | | X |
| 41 | <i>Acanthasauraarmata</i> | X | | | | X | |
| 42 | <i>Bronchoelacristatella</i> | X | | | | X | |
| 43 | <i>Calotesemma</i> | X | | | | | |
| 44 | <i>Draco cristatellus</i> | | | | | X | |
| 45 | <i>Draco fimbriatus</i> | | | | | X | |

| | | | | | | |
|--------------------------|------------------------------------|---|---|---|---|---|
| 46 | <i>Draco formosus</i> | | | | X | |
| 47 | <i>Draco maximus</i> | X | | | X | X |
| 48 | <i>Draco melanopogon</i> | X | | X | X | X |
| 49 | <i>Draco quinquefasciatus</i> | X | | | X | |
| 50 | <i>Draco sumatranus</i> | X | | X | | |
| 51 | <i>Gonocephalusabbotti</i> | X | | X | X | X |
| 52 | <i>Gonocephalusbellii</i> | X | | | | |
| 53 | <i>Gonocephalusgrandis</i> | X | | X | X | X |
| Eublepharidae | | | | | | |
| 54 | <i>Aeluroscalabotesfelinus</i> | | | | X | |
| Gekkonidae | | | | | | |
| 55 | <i>Cosymbotescraspedotus</i> | | | | X | |
| 56 | <i>Cyrtodactylusconsobrinus</i> | X | | | | X |
| 57 | <i>Cyrtodactylusquadrivirgatus</i> | X | | | X | X |
| 58 | <i>Gekkosmithii</i> | X | | | | X |
| 59 | <i>Gehyramutilata</i> | | | | X | |
| 60 | <i>Ptychozoonhorsfieldi</i> | X | | | | |
| 61 | <i>Ptychozoonkuhlii</i> | | | | X | |
| Scincidae | | | | | | |
| 62 | <i>Dasiaolivacea</i> | X | | | X | |
| 63 | <i>Eutropismultifasciatus</i> | X | | | X | |
| 64 | <i>Sphenomorphus maculates</i> | X | | | | |
| 65 | <i>Sphenomorphus cf. butleri</i> | | | | X | |
| Varanidae | | | | | | |
| 66 | <i>Varanusdumerilii</i> | | | | X | |
| 67 | <i>Varanusrudicollis</i> | X | | | | |
| 68 | <i>Varanussalvator</i> | X | | | | |
| REPTILES (SNAKES) | | | | | | |
| Typhlopidae | | | | | | |
| 69 | <i>Ramphotyphlopslineatus</i> | | X | | | |
| 70 | <i>Typhlopsmuelleri</i> | | | | | X |
| Pythonidae | | | | | | |
| 71 | <i>Python brongersmai</i> | | X | | | |
| 72 | <i>Python reticulatus</i> | | X | | | |
| Colubridae | | | | | | |
| 73 | <i>Ahaetullaprasina</i> | | X | | | |
| 74 | <i>Aplopeltura boa</i> | | | X | | |
| 75 | <i>Boigacynodon</i> | | X | | | |
| 76 | <i>Boigadendrophila</i> | | X | | X | X |
| 77 | <i>Calamariaschlegelii</i> | | | | | X |
| 78 | <i>Dendrelaphisformosus</i> | | X | | | |

| | | | | |
|---------------------------|------------------------------------|---|---|---|
| 79 | <i>Elapheflavolineata</i> | X | | |
| 80 | <i>Elapheprasina</i> | X | | |
| 81 | <i>Enhydrisindica</i> | X | | |
| 82 | <i>Gonylosoma cf. longicauda</i> | | X | |
| 83 | <i>Gonyosomaoxycephalum</i> | X | | |
| 84 | <i>Lepturophisalbofuscus</i> | X | | |
| 85 | <i>Lycodonsubcintus</i> | | X | X |
| 86 | <i>Oligodonpurpurescens</i> | X | | |
| 87 | <i>Oreocalamushanitschi</i> | X | | |
| 88 | <i>Psammodynastespulvurulentus</i> | X | | |
| 89 | <i>Pseudorhabdionlongiceps</i> | X | | |
| 90 | <i>Rhabdophischrysargos</i> | | | X |
| 91 | <i>Xenochrophistrianguligerus</i> | | X | |
| Elapidae | | | | |
| 92 | <i>Bungaruscandidus</i> | X | | |
| 93 | <i>Calliophisbivirgata</i> | X | | |
| 94 | <i>Najasumatrana</i> | X | | |
| Viperidae | | | | |
| 95 | <i>Pariashageni</i> | X | | |
| 96 | <i>Pariassumatranus</i> | | X | |
| 97 | <i>Popeiapopeiorum</i> | X | | |
| 98 | <i>Tropidolaemuswagleri</i> | X | | X |
| REPTILES (TURTLES) | | | | |
| Emydidae | | | | |
| 99 | <i>Batagurborneoensis</i> | X | | |
| 100 | <i>Cuoraamboinensis</i> | X | | |
| 101 | <i>Cyclemysdentata</i> | X | | |
| 102 | <i>Heosemyspinosa</i> | X | | |
| 103 | <i>Notochelysplatynota</i> | X | | |
| Trionychidae | | | | |
| 104 | <i>Amydacartilaginea</i> | X | | |
| 105 | <i>Doganiasubplana</i> | X | | X |

The present effort represents the fifth herpetological survey of the Belum- Temengor Forest Reserves, but constitutes the first survey of the Sungai Enam Basin, Temengor. In this study, we added three new records of snakes namely *Typhlopsmuelleri*, *Calamariaschlegelii* and *Rhabdophischrysargos*. This brings the total number of herpetofauna documented in the Belum-Temengor area to 105. This low herpetofaunal diversity may be largely attributed to the limited sampling period (four search days) and restricted to three main trails (Trail 1, Trail 2 and Waterfall 4) of the area. In addition, herpetological surveys in general require longer periods of time due to the elusive and secretive nature of many species of snakes and lizards which they tend to slip away

unnoticed at the approach of man (Ibrahim et al., 2013). Hence, cross sectional surveys normally tend to under- represent the actual herpetofaunal diversity, especially so in the dense tropical forest. According to Inger (2003), numerous factors such as duration of sampling period, area of coverage, sampling technique, topography, weather, types of microhabitat and activity pattern are known to influence the number of species recorded in each area. Besides, continuous surveys are also necessary to increase the chances of detection and increase the total number of herpetofauna recorded (Shahriza et al., 2012). For example, an early record of amphibian's diversities of Peninsular Malaysia by Berry (1975) found 83 species. Thirty years later, this number increased to 100 species (Inger, 2005) and seven more were identified in the following five years (Chan et al. 2010). This scenario clearly reveals that Malaysian forests are very rich in herpetofaunal assemblage. Similarly, in this study, the number of herpetofauna of the Belum- Temengor Forest Reserves is still increasing and additional new records can be expected with continuous intensive surveys. Most reptiles caught especially snakes are juveniles which indicated that the area is still viable to the snakes although the base camp site is already disturbed and slowly been developed into an ecotourism site.

Acknowledgements

We wish to thank the Pulau Banding Foundation and Yayasan EMKAY for the opportunity to participate in the expedition. We also wish to express our gratitude to the staffs and crew of PulauBanding Foundation, Yayasan EMKAY and Belum Rainforest Resort for excellent logistics and hospitality provided, and all expedition members who have helped us with the additional herpetofaunal specimens and information especially from few other research teams including FRIM small mammals team, insect team, DWNP Perak Branch survey team and USM fish team. This study was also partially funded by UniversitiSains Malaysia's Research University Grant No. 5001/PJJAUH/834059 awarded to the last author.

References

- Berry P. Y. 1975. The Amphibian Fauna of Peninsular Malaysia. Tropical Press, Kuala Lumpur.
- Chan K. O., Daicus B., Norhayati, A. 2010. A revised checklist of the Amphibians of Peninsular Malaysia. *Russian Journal of Herpetology*, 17(3): 202-206.
- Cox M. J., van Dijk P. P., Nabhitabhata J., Thirukhupl K. 1998. A Photographic Guide to Snakes and Other Reptiles of Peninsular Malaysia, Singapore and Thailand. New Holland Publishers, London.
- Das I. 2010. A Field Guide to the Reptiles of South-East Asia. New Holland Publishers, London.
- Diong C. H., Kiew B. H., Lim B. L. 1995. An annotated checklist of the lizard fauna in the Temengor Forest Reserve, Hulu Perak, Malaysia. *Malayan Nature Journal*, 48: 353-356.
- Frost D. R. 2011. Amphibian Species of the World: an Online Reference. Version (31 January, 2011). Electronic Database accessible at <http://research.amnh.org/vz/herpetology/amphibia/Am. Mus. Nat. Hist., New York, USA>.
- Grismer L.L., Sukumaran J., Grismer J.L., Youmans T.M., Wood P. L., Johnson R. 2004 Report on the Herpetofauna from the Temengor Forest Reserve, Perak, West Malaysia. *Hamadryad* 29 (1): 15-32
- Ibrahim J., Shahrul A.M.S., Norhayati A., Chan K.O., Muin, M.A. 2008. The Common Amphibians and Reptiles of Penang Island. Penang State Forestry Department, Penang.

- Ibrahim J., Amirah H., Shahriza S., Nurhafizah I., Zalina A., Yap C. H., Nurliza A. M., Nurhafizah C. Z. 2013. Additions to the herpetofauna of Jerejak Island, Penang, Peninsular Malaysia. *Malayan Nature Journal*, 64(4): 213-232.
- Inger, R. F. 2003. Sampling Biodiversity in Bornean Frogs. *The Natural History Journal of Chulalongkorn University*, 3(1): 9-15.
- Inger R. F. 2005. The Frog Fauna of the Indo- Malayan Region as it Applies to Wallace's Line. In:Tuen, A. A. and Das, I (eds). *Wallace in Sarawak-150 Years Later. An International Conference on Biogeography and Biodiversity*, pp. 82-90. Institute of Biodiversity and Environmental Conservation, University Malaysia Sarawak.
- Kiew B. H., Diong C. H., Lim B. L. 1995. An Annotated Checklist of the Amphibian Fauna in the Temengor Forest Reserve, Hulu Perak, Malaysia. *Malayan Nature Journal*, 48: 347-351.
- Lim B. L., Ratnam L., NorAzman H. 1995a. Snakes examined from the Sungai Singgor area of Temengor, Hulu Perak, Malaysia. *Malayan Nature Journal*, 48: 57- 364.
- Lim B. L., Ratnam L., NorAzman H. 1995b. Turtles from the Sungai Singgor area of Temengor, Hulu Perak, Malaysia. *Malayan Nature Journal*, 48:366- 369.
- Norsham Y., Bernard H., Chew K. L., Yong H. S., Yap M. N., Lim B. L. 2000. An Annotated Checklist of Herpetofauna in the Northern Part of Belum Forest Reserve, Perak Peninsular, Malaysia. *Malayan Nature Journal*, 54: 245-251.
- Shahriza S., Ibrahim J., ShahrulAnuar M. S., Abdul Muin M. A. 2012. Herpetofauna of Peta Area of Endau-Rompin National Park, Johor, Malaysia. *Pertanika J. Trop. Agric. Sci.*, 35 (3): 553 – 567.
- Sukumaran J. 2002. The amphibian fauna of a forested area in Temengor, Perak, Malaysia, with the first record of *Philautus parvulus* (Boulenger, 1893) in the Malay Peninsula. *Hamdryad*, 27:1-10.