



# AGRIVITA

## Journal of Agricultural Science

### Distribution of the Endophytic Fungi in Apple Leaves

Aminudin Afandhi<sup>\*)</sup>, Fery Abdul Choliq, Havinda Anggrilika W.S. and Hagus Tarno

Pest and Plant Disease Department, Faculty of Agriculture, Universitas Brawijaya, Indonesia

#### ARTICLE INFO

##### Keywords:

Abundance  
Apple leaf  
Diversity  
Endophytic fungi

##### Article History:

Received: August 3, 2017

Accepted: December 8, 2017

<sup>\*)</sup> Corresponding author:

E-mail: aaf\_fp@ub.ac.id

#### ABSTRACT

Endophytic fungi have roles to protect the plant from herbivorous insects and diseases, and to support the absorption process of nutrients needed by plants for photosynthesis. Related to the important role of endophytic fungi, research was aimed to identify fungal endophytes associated with young, mature and old leaves on apple and to evaluate the effect of leaf ages on the abundance and diversity of endophytic fungi. The research was conducted in Biological Control Laboratory, Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Brawijaya from May to September 2016. Apple leaves sampling was done diagonally, and taken from nine apple trees. Based on macroscopic and microscopic characteristics, 38 isolates were identified. In the old apple leaves, it obtained 17 isolates that consisted of 5 genera, mature leaves obtained 14 isolates consisted of 6 genera, and young leaves 7 isolates obtained consists of 2 genera. The mature leaves have highest abundance and diversity of endophytic fungi compared to young and old leaves. Endophytic fungi apple leaf dominated by *Aspergillus*. Mature and young leaves were different from old one based on Bray-Curtis similarity.

#### INTRODUCTION

Endophytic fungi are microorganisms that are in the plant tissue during certain periods of their life cycle, without causing damage to the plants (Hilarino et al., 2011). It play an important role in balancing ecosystems and benefiting the host such as increasing plant growth (Zheng et al., 2017).

Abundance and diversity of endophytic fungi species increases with increasing leaf age (López-González et al., 2017). Hilarino et al. (2011) claimed that differences in the composition of endophytic fungi on the leaves of *Bauhinia brevipes* influenced by the age of the leaf. Endophytic fungi colonization is higher in the leaves of certain associated with anatomical structures and susceptibility to infection, changes in biochemical influences (Fernandes, Oki, Sanchez-Azofeifa, Faccion, & Amaro-Arruda, 2011) and leaf nutrient content such as chlorophyll.

Understand the abundance and diversity of endophytic fungi on apple leaves is the key to know in the ecology and the association between endophytic fungi and plants. The research of endophytic fungi have been many developed knowledge about influence of plant resistance to climate, pathogens and insect pests. Endophytic fungi that colonize a host plant will affect the presence of insects, especially pest insects into the host. Endophytic fungi *Nigrospora oryzae* and *Cladosporium Arenicola* isolated from plant *Tinospora cordifolia* able to suppress the development of the larvae of *Spodoptera litura* (Thakur, Singh, Kaur, & Kaur, 2014). In addition, endophytic fungi are also capable of inhibiting pathogens thus showing potential as a biocontrol agent (López-González et al., 2017). The important role can be influenced by abundance and diversity of endophytic fungi in the ecosystem as a whole.

ISSN: 0126-0537 Accredited by DIKTI Decree No: 60/E/KPT/2016

**Cite this as:** Afandhi, A., Choliq, F. A., Anggrilika W. S., H., & Tarno, H. (2018). Distribution of the endophytic fungi in apple leaves. AGRIVITA Journal of Agricultural Science, 40(1), 91–100. <http://doi.org/10.17503/agrivita.v40i1.1563>

Apple as one of popular fruit in Indonesia, is cultivated in the highlands of Indonesia such as in Batu City. One obstacle is the cultivation of apple pests and diseases. Pests and diseases occurs after the leaves removed. Endophytic fungi of apple leaves in Brazil are *Alternaria* sp., *Botryosphaeria* sp., *Cladosporium* sp., *Colletotrichum* sp., *Epicoccum* sp., *Fusarium* sp., and *Xylaria* sp. (Camatti-Sartori et al., 2005). Endophytic fungi have roles to protect the plant from herbivorous insects and diseases, to support the absorption process of nutrients needed by plants for photosynthesis. Meanwhile, the results of plant photosynthesis can be used fungus for life sustainability. Related to the important roles of endophytic fungi on a host plant, this research aimed to identify fungal endophytes associated with young, mature and old leaves on apple and to evaluate the effect of leaf ages on the abundance and diversity of endophytic fungi.

## MATERIALS AND METHODS

### Sampling and Isolation of Endophytic Fungi

Sampling was conducted on apple orchard in Bumiaji Village, Bumiaji Subdistricts, Batu City (Coordinates: S 7°51'32.4" and E 112°32'20.1"), altitude 1062 m above sea level (masl). Isolation and identification conducted in Biological Control Laboratory of Department of Pests and Plant Diseases, Faculty of Agriculture, Universitas Brawijaya. The study was conducted from May to September 2016.

For sampling purposes, nine apple trees were done diagonally that leaves samples drawn more equally. Each one tree taken 6 leaves, 2 leaves were taken from the young leaves of the sequence number 2 of shoots, 2 mature leaves were taken from the leaves of the sequence number 4 of shoots, and 2 old leaves were taken from the leaves of the sequence number 6 of shoots. There were 54 strands of total leaf samples. Parts of plants were taken to the exploration process were in a healthy condition. Leaf samples were collected in an envelope and taken to a laboratory to be isolated.

Early stages of isolation to prepared a leaves sample apple was washed with running water until clean of dirt, then dried on sterile wipes. Sterilization using method of Greenfield et al. (2015) which leaves soaked in the solution of 0.5 % NaOCl for 3 minutes, 70 % alcohol for 1 minute and rinsed with sterile distilled water for 3 times each 15 seconds and then dried using sterile wipes. Each leaf was cut 1 cm x 1 cm as much as 6 leaf pieces, so the Petri

dish contained 6 leaf pieces. Then a leaf samples were isolated in a Sabauraud Dextrose Agar Yeast media (SDAY) in 9 cm diameter Petri dishes and coded. Materials for the manufacture of media SDAY namely dextrose 10 g, peptone 2.5 g, to 20 g, yeast extract 2.5 g, chloramphenicol 0.5 g and distilled water 1 l. Last rinse of distilled water was taken 1 ml and isolated in a new SDAY media, this treatment was used as an insulating material sterility testing of microorganisms other than endophyte.

### Identification of the Isolates

Identification of isolates morphological macroscopic and microscopic. Identification was done at the genus level. Observations were made every day for 7 days after the inoculation, if found new colonies that grow so do the purification of the new SDAY media (Subculture 1). Macroscopic identification was done by observing the macroscopic characteristics includes color colony, colony surface texture, diameter and colonies form. Microscopic identification was done by taking most isolates selected using a needle ose then placed on an object glass and closed with a cover glass. Observations were made under a microscope with parameters including hyphae (sectional or not and branched or not), the shape of conidiophores, conidia (shape and size) and specific characteristics to determine the type of fungus by using reference books *Illustrated Genera of Imperfect Fungi*.

### Statistical Analysis

#### Abundance of Endophytic Fungi

The rate of colonization percentage of endophytes was calculated by using equation 1 (Chougule & Andoji, 2015).

$$\text{Rate of colonization} = \frac{\text{NSCSF}}{\text{TNSO}} \times 100\% \quad (1)$$

where: NSCSF is Number of Segmented Colonized by Single Fungus; TNSO is Total Number of Segmented observed

#### Diversity of Endophytic Fungi

Several indices to describe alfa diversity such as Simpson Index, species diversity indices and species Evenness indices were calculated (Habib, Heshmati, & Siroosi, 2014; Shannon, 1948; Tarno, Septia, & Aini, 2016). Simpson index (D), species diversity indices (H') and Species Evenness indices (E) were described in Equation 2, 3, and 4 respectively

$$D = \sum_{i=1}^s P_i^2 \quad \dots\dots\dots (2)$$

$$H' = -\sum_{i=1}^s (P_i)(\log_2 P_i) \quad \dots\dots\dots (3)$$

Remarks: Simpson index, index of species diversity, proportion of individuals of species  $i$  in the community, number of species in the sample, and logarithm are  $D$ ,  $H'$ ,  $P_i$ ,  $S$  and  $\log$  respectively.

$$E = \frac{H'}{\ln(S)} \quad \dots\dots\dots (4)$$

Remarks: index of species diversity, exponential logarithm, and proportion of individuals of species are  $H'$ ,  $\ln$  and  $S$ , respectively.

Value of Shannon-Winner Indices ( $H'$ ) are describes as low level of diversity (with value  $<1$ ), low individual distribution of each species, middle level of diversity (1-3), middle level of individual distribution for each species, and high level of diversity ( $>3$ ), high level of individual distribution for each species (Tarno, Septia, & Aini, 2016).

Based on value of Evenness index, there are three categories i.e.: low ( $0.00 < E < 0.50$ ) described as under pressure community, medium level ( $0.50 < E < 0.75$ ) described as unstable community, and high level ( $0.75 < E < 1.00$ ) described as stable community. In addition, beta diversity was also calculated by using Bray-Curtis similarity index to describe similarity between three different of ages of leaf on apple based on all observed endophytic fungal species. Alfa and beta diversities were calculated by using Vegan package of R Software (R Core Team, 2013).

## RESULTS AND DISCUSSION


### Abundance Endophytic Fungi

Six of endophytic fungal genera were identified from old, mature, and young apple leaves i.e.: *Aspergillus*, *Penicillium*, *Fusarium*, *Acremonium*, *Curvularia* and *Alternaria*. The results showed that,


in the old apple leaves, it obtained 17 isolates that consisted of five genera, mature leaves obtained 14 isolates consisted of six genera, and young leaves 7 isolates obtained consists of two genera. Abundance of endophytic fungi on host plants affected by the isolated organ, the sampling locations and age of leaves. Abundance of endophytic fungi obtained from a sample of leaves colonized endophytic fungi, endophytic fungi colonization percentage and the percentage of total colonization of endophytic fungi.

Each leaves sample with different criteria namely the old, mature and young apple leaves colonized types of endophytic fungi were different (Fig. 1). Colonization of endophytic fungi varies according to leaf age (Sanchez-Azofeifa, Oki, Fernandes, Ball, & Gamon, 2012). *Aspergillus* sp. 1 and *Aspergillus* sp. 3 samples colonized old, mature and young apple leaves. *Aspergillus* sp. 2 colonized the old and mature leaves samples. *Penicillium* sp. 1, *Penicillium* sp. 2, *Fusarium* sp. 1, *Acremonium* sp. 1, and *Curvularia* sp. 1 colonized the old leaves sample. *Penicillium* sp. 3, *Penicillium* sp. 4, *Fusarium* sp. 2, *Acremonium* sp. 2, *Acremonium* sp. 3, *Curvularia* sp. 2, and *Alternaria* sp. 1 colonized the mature leaves sample. Meanwhile, *Alternaria* sp. 2 and *Alternaria* sp. 3 colonized the young leaves.

*Aspergillus* sp. 1 was found in all ages of leaf samples with the highest population. *Aspergillus* sp. 3 and A. sp.2 followed based on population of species (Fig. 2). It's related to report on paddy that *Aspergillus* is one of the most recovered genus (Zakaria, Yaakop, Salleh, Zakaria, & Pinang, 2010). In addition, *Fusarium* was the most common endophytic fungal genus recovered on paddy, followed by *Aspergillus*, *Curvularia*, *Penicillium*, *Gilmaniella* and *Arthrobotrys foliicola* (Zakaria, Yaakop, Salleh, Zakaria, & Pinang, 2010). In case of *Fusarium* and *Curvularia* had the commonly occurrences in the seeds (Zakaria, Yaakop, Salleh, Zakaria, & Pinang, 2010). *Aspergillus* spp., *Penicillium* spp., *Fusarium* spp., and *Acremonium* spp. were reported found on *Pterocarpus indicus* (Tarno, Septia, & Aini, 2016).



**Fig. 1.** Population of colonized endophytic fungi in each age of leaf; *Aspergillus* sp1 (Asp.1), *Aspergillus* sp2 (Asp.2), *Aspergillus* sp3 (Asp.3), *Penicillium* sp1 (Pen.1), *Penicillium* sp2 (Pen.2), *Penicillium* sp3 (Pen.3), *Penicillium* sp4 (Pen.4), *Fusarium* sp1 (Fus.1), *Fusarium* sp2 (Fus.2), *Acremonium* sp1 (Acr.1), *Acremonium* sp2 (Acr.2), *Acremonium* sp3 (Acr.3), *Curvularia* sp1 (Cur.1), *Curvularia* sp2 (Cur.2), *Alternaria* sp1 (Alt.1), *Alternaria* sp2 (Alt.2), and *Alternaria* sp3 (Alt.3).



**Fig. 2.** Boxplot of population of each endophytic fungus species on all ages of leaves; *Aspergillus* sp1 (Asp.1), *Aspergillus* sp2 (Asp.2), *Aspergillus* sp3 (Asp.3), *Penicillium* sp1 (Pen.1), *Penicillium* sp2 (Pen.2), *Penicillium* sp3 (Pen.3), *Penicillium* sp4 (Pen.4), *Fusarium* sp1 (Fus.1), *Fusarium* sp2 (Fus.2), *Acremonium* sp1 (Acr.1), *Acremonium* sp2 (Acr.2), *Acremonium* sp3 (Acr.3), *Curvularia* sp1 (Cur.1), *Curvularia* sp2 (Cur.2), *Alternaria* sp1 (Alt.1), *Alternaria* sp2 (Alt.2), and *Alternaria* sp3 (Alt.3).


Percentage colonization of endophytic fungi in old, mature and young apple leaves can be seen on Fig. 3. In the old leaves obtained five genera are *Aspergillus*, *Penicillium*, *Fusarium*, *Acremonium* and *Curvularia*. In the mature leaves obtain six genera are *Aspergillus*, *Penicillium*, *Fusarium*, *Acremonium*, *Curvularia* and *Alternaria*. While the young leaves obtained two genera are *Aspergillus*, and *Alternaria*. Colonization of endophytic fungi genus *Aspergillus* in old, mature and young apple leaves showed the highest percentage (Fig. 4). Colonization of endophytic fungi on old, mature and young apple leaves showed their dominance of the genus *Aspergillus*. In this study, samples of old, mature and young apple leaf isolated endophytic fungi have different total colonization. The results showed that the mature apple leaves have the highest percentage of total colonization of 66.67 %, followed by 50 % of the old leaves and young leaves 38.89 %. It shows that the mature leaves were higher compared to the old and young apple leaves based on colonization or abundance. The colonization is influenced by the age of leaves.

Abundance and diversity of endophytic fungal species increases with increasing leaf age (López-González et al., 2017).


### Diversity of Endophytic Fungi

Diversity of endophytic fungal species was analyzed by using Shannon-Winner ( $H'$ ) and Simpson Diversity Indexes ( $D$ ). In addition, Evenness index ( $E$ ) was also used to analyze diversity. 14, 17 and 7 isolates of endophytic fungi collected from mature, old and young leaves resulted six, five and two endophytic fungal genera, respectively.


Based on  $H'$ , mature and young leaves resulted the highest (more than 2.0) and lowest values (more than 1.0) (Fig. 5). Mature, old and young leaves produced index's value between 1 and 3, and it's mean that diversities on each age of leaf on the middle level of diversity status (middle level of individual distribution for each species) (Tarno, Septia, & Aini, 2016). Evenness index showed that all three ages of leaf produced the high value (more than 0.75) or stable communities.  $H'$ ,  $D$  and  $E$  indexes showed that all ages of leaf on the middle level diversity and stable communities.



**Fig. 3.** Percentage of endophytic fungi colonization; *Aspergillus* sp1 (Asp.1), *Aspergillus* sp2 (Asp.2), *Aspergillus* sp3 (Asp.3), *Penicillium* sp1 (Pen.1), *Penicillium* sp2 (Pen.2), *Penicillium* sp3 (Pen.3), *Penicillium* sp4 (Pen.4), *Fusarium* sp1 (Fus.1), *Fusarium* sp2 (Fus.2), *Acremonium* sp1 (Acr.1), *Acremonium* sp2 (Acr.2), *Acremonium* sp3 (Acr.3), *Curvularia* sp1 (Cur.1), *Curvularia* sp2 (Cur.2), *Alternaria* sp1 (Alt.1), *Alternaria* sp2 (Alt.2), and *Alternaria* sp3 (Alt.3).



**Fig. 4.** Total percentage of colonization of endophytic fungi



**Fig. 5.** Diversity indices of endophytic fungi on three ages of leaf

Abundance and diversity of endophytic fungi from mature leaves higher than the old and young leaves of apple allegedly relates to compounds of photosynthesis that supports the growth and development of endophytic fungi. Getting old apple leaves are green, green leaves tend to lack the chlorophyll higher than young leaves. The higher the content of chlorophyll in the leaves, the higher the rate of photosynthesis. Results of photosynthesis used for plant growth and development but it is

also used endophytic fungi to survive in the plant tissue. Thus, the leaves have a higher chlorophyll content is characterized by dark green color, the number and types of endophytic fungi on the leaves higher. Hilarino et al. (2011) in his study claimed that differences in the composition of endophytic fungi on the leaves of *Bauhinia brevipes* influenced by the age of the leaf. The leaves are aging tend to receive more nutrients, so the old leaves to get more chlorophyll, leaf color more green old age.


Aminudin Afandhi et al.: Diversity of Endophytic Fungi in Apple Leaves.....

Fernandes, Oki, Sanchez-Azofeifa, Faccion, & Amaro-Arruda (2011) adds that the old leaves more favorable to the colonization of the fungus, leaf biochemical changes influence affecting the distribution of endophytic colonization. Endophytic fungi are mutualistic symbiosis with the host plant. Endophytic fungi help the absorption of nutrients needed by plants for photosynthesis and photosynthesis results can be used fungi in order to survive. Abundance and diversity of endophytic fungi in old leaves lower than mature leaves, old leaves suspected decreased chlorophyll content so that colonization of endophytic fungi that are in the old leaves is reduced. Decreased levels of chlorophyll in the leaves affects the rate of photosynthesis, photosynthetic rate lower if the compounds that plants need and endophytic fungi on the wane. Plant and fungal endophyte can't absorb the nutrients optimally. Endophytic fungi survive in host plants are influenced by environmental factors such as temperature and humidity, chemicals and leaf anatomic structures. Thus, affects the abundance and diversity of endophytic fungi (Sanchez-Azofeifa, Oki, Fernandes, Ball, & Gamon, 2012). Changes in endophytic fungi communities are related to leaf age (Fernandes, Oki, Sanchez-Azofeifa, Faccion, & Amaro-Arruda, 2011, Hilarino et al., 2011) and even to plant tissue types (Sanchez-Azofeifa, Oki, Fernandes, Ball, & Gamon, 2012). Many biotic and abiotic factors are involved in colonization, including chemical and plant structure (Sanchez-Azofeifa, Oki, Fernandes, Ball, & Gamon, 2012), and environmental conditions where endophytic fungi are present (Fernandes, Oki, Sanchez-Azofeifa, Faccion, & Amaro-Arruda, 2011).

Endophytic fungi in addition to helping the plant in the absorption of nutrients, it also protects the plants against pathogens and insect herbivores. Plant defense system is stronger when various types of endophytic fungi are mutualistic symbiosis with plants. Many reports of endophytic fungi are mutualistic to the plant than endophytes are pathogen. Host plant and endophytic fungi mutualism association by way of endophytic fungi obtain nutrients and survive in plants tissue, while the host plants get nutrients in the defense system, growth, and reproduction. Thus, leaves

that have an abundance and diversity of high endophyte fungus is thought to have an important and beneficial role of host. Compounds mycotoxins produced by endophytic fungi, capable of being weapons of host defense against insect herbivores and pathogens. Although endophytic fungi cannot directly kill insect herbivores, but can suppress the presence of insect herbivores on host plants. According Irmawan (2007) stated that endophytic fungi on plants grasses produce mycotoxins such as alkaloids capable of protecting the host from attack herbivorous invertebrates, nematodes and pathogens. Endophytic fungi as bio indicators of the health of plants, endophytic fungi that colonize a host plant will affect the existence of herbivorous insects and pathogens. An important role is influenced by the diversity and abundance of endophytic fungi in host. Endophytic fungi *Nigrospora oryzae* and *Cladosporium Arenicola* isolated from plant *Tinospora cordifolia* able to suppress the development of the larvae of *Spodoptera litura* (Thakur, Singh, Kaur, & Kaur, 2014). Endophytic fungi were able to suppress the presence of plant pathogens, fungal endophyte usually antagonistic to plant pathogenic fungi. Test antagonistic endophytic fungi *Hyalodendron* sp. and *Cephalosporium* sp. derived from potato plants could reduce *Phytophthora infestans* attacks by 66.56 % and 61.52 % (Tirtana, Sulistyowati, & Cholil, 2013). Endophytic fungi have been considered as a promising source for the development of biological control agents. In addition, the endophytic fungi lately also reportedly able to produce metabolites which served to protect host plants from extreme environmental conditions and can promote plant growth. Endophytic fungi produce phytohormones such as indole-3-acetic acid (IAA), cytokines, and other growth promoters' compound. Endophytic fungi produces an antifungal antibiotic as pathogenic to humans is *Candida albicans* and *Trichopyton* spp, paclitaxel and its derivatives are useful as anticancer substance was first discovered in endophytic microorganisms. Paclitaxel is a diterpenoid compound obtained in *Taxus* plants. Endophytic fungi benefit host plants by producing secondary bioactive metabolites (Mei & Flinn, 2010).





**Fig. 6.** Similarity analysis of three ages of leaf based on its endophytic communities

Endophytic fungi of the genus *Aspergillus* predominantly found in old, mature and young apple leaves. Ariyono, Djauhari, & Sulistyowati (2014) states that the kale leaves land in organic fields are dominated by the genus *Aspergillus*. *Aspergillus* sp. is a cosmopolitan fungi has a wide host range, being in nature as a saprophyte fungi, parasites and spores can easily spread through the air. Fungus *Penicillium* sp., *Fusarium* sp., *Acremonium* sp. and *Curvularia* sp., found on old leaves and mature, but not found in young leaves apple. Endophytic fungi *Fusarium* sp., *Penicillium* sp., and *Curvularia* sp. endophyte found in the leaves of potato (Tirtana, Sulistyowati, & Cholil, 2013). While, according to Hilarino et al. (2011), states that the fungus endophyte *Acremonium* sp. obtained from the leaves *Bauhinia brevipes*. Fungus *Alternaria* sp., found in the mature leaves and young but not found on old leaves apple. Camatti-Sartori et al. (2005) states that the endophytic fungus *Alternaria* sp. reported as an endophyte in apple leaves in Brazil. Endophytic fungi obtained from old, mature and young apple leaves have different types as it relates to micro ecological and physiological condition of each leaf that supports the growth of endophytic fungi, thus affecting the number and types of endophytic fungi were found. Endophytic fungi isolated from old, mature and young apple leaves has a type and the number of different isolates, allegedly influenced by the survivability of endophytic fungi against micro ecological and physiological condition of each leaf.

Endophytic fungi survive in host plants are influenced by environmental factors such as temperature and humidity, chemicals and leaf anatomic structures. Thus, it affects the abundance and diversity of endophytic fungi (Sanchez-Azofeifa,

Oki, Fernandes, Ball, & Gamon, 2012). The abundance and diversity of endophytic fungi in host plants illustrates the components of biodiversity. In addition, as bio indicators of the health of plants. Each host plants have diverse environmental conditions for growing a fungal endophyte. In almost all tissues of healthy plants are endophytic microorganisms that colonize. Many studies suggest benefits of colonization of endophytic fungi in plant tissues such as strengthening plant resistance to insect herbivores, pathogens, defense against extreme weather, produces growth regulators, antibiotics, anticancer and antioxidant. However, so far the interaction of each type of endophytic fungi that colonize a host plant has not been clearly reported. Several types of endophytic fungi that colonize a host plant allegedly symbiotic mutualism, so as to protect crops from adverse environmental effects.

Bray–Curtis similarity was adopted in this analysis to evaluate the similarity between all ages of leaf. It was widely employed in multivariate analysis of assemblage data, for sound biological reasons (Clarke, Somerfield, & Chapman, 2006). Based on Fig. 6, there are two clusters of diversity between all three ages of leaf. Mature and young leaves are on the same cluster, and it showed different from old leaves. Composition of endophyte fungal species between mature and young leaves were different to the old one.

## CONCLUSION

The mature leaves have highest abundance and diversity of endophytic fungi compared to young and old leaves. In the old apple leaves, it obtained 17 isolates that consisted of 5 genera, mature leaves obtained 14 isolates consisted of 6 genera,



and young leaves 7 isolates obtained consists of 2 genera. *Aspergillus* sp. dominated on all ages of apple leaf. Mature and young leaves were different from old one based Bray-Curtis similarity.

## REFERENCES

- Ariyono, R. Q., Djauhari, S., & Sulistyowati, L. (2014). Keanekaragaman jamur endofit daun kangkung darat (*Ipomoea reptans* Poir.) pada lahan pertanian organik dan konvensional [The diversity of the endophytic fungi of land kangkung (*Ipomoea reptans* Poir.) on organic and conventional farmland]. *Jurnal HPT*, 2(1), 19-28. Retrieved from <http://jurnalhpt.ub.ac.id/index.php/jhpt/article/view/63>
- Camatti-Sartori, V., Da Silva-Ribeiro, R. T., Valdebenito-Sanhueza, R. M., Pagnocca, F. C., Echeverrigaray, S., & Azevedo, J. L. (2005). Endophytic yeasts and filamentous fungi associated with southern Brazilian apple (*Malus domestica*) orchards subjected to conventional, integrated or organic cultivation. *Journal of Basic Microbiology*, 45(5), 397-402. <http://doi.org/10.1002/jobm.200410547>
- Chougule, P. M., & Andoji, Y. S. (2015). Isolation and frequency distribution of Endophytic fungi inhabiting *Justicia adhatoda* L. *Research Journal of Life Sciences, Bioinformatics, Pharmaceutical and Chemical Sciences*, 1(4), 215-218. Retrieved from <http://rjlbpcs.com/wp-content/uploads/2015/Nov-Dec/4-Isolation-and-frequency-distribution-of-Endophytic-fungi-inhabiting.pdf>
- Clarke, K. R., Somerfield, P. J., & Chapman, M. G. (2006). On resemblance measures for ecological studies, including taxonomic dissimilarities and a zero-adjusted Bray-Curtis coefficient for denuded assemblages. *Journal of Experimental Marine Biology and Ecology*, 330(1), 55-80. <https://doi.org/10.1016/j.jembe.2005.12.017>
- Fernandes, G. W., Oki, Y., Sanchez-Azofeifa, A., Faccion, G., & Amaro-Arruda, H. C. (2011). Hail impact on leaves and endophytes of the endemic threatened *Coccoloba cereifera* (Polygonaceae). *Plant Ecology*, 212(10), 1687-1697. <http://doi.org/10.1007/s11258-011-9941-z>
- Greenfield, M., Pareja, R., Ortiz, V., Gómez-Jiménez, M. I., Vega, F. E., & Parsa, S. (2015). A novel method to scale up fungal endophyte isolations. *Biocontrol Science and Technology*, 25(10), 1208-1212. <http://doi.org/10.1080/09583157.2015.1033382>
- Habib, R. K., Heshmati, G., & Siroosi, H. (2014). Comparison of ecological patches' potentials and functions in rangeland ecosystems (case study: Qahavand Rangelands, Hamedan Province, Iran). *Journal of Rangeland Science*, 4(3), 234-245. Retrieved from [http://www.rangeland.ir/article\\_513017.html](http://www.rangeland.ir/article_513017.html)
- Hilarino, M. P. A., Silveira, F. A. de O. e, Oki, Y., Rodrigues, L., Santos, J. C., Corrêa Junior, A., ... Rosa, C. A. (2011). Distribution of the endophytic fungi community in leaves of *Bauhinia brevipes* (Fabaceae). *Acta Botanica Brasilica*, 25(4), 815-821. <http://doi.org/10.1590/S0102-33062011000400008>
- Irmawan, D. E. (2007). *Kelimpahan dan keragaman cendawan endofit pada beberapa varietas padi di Kuningan, Tasikmalaya dan Subang, Jawa Barat* [Abundance and diversity of endophytic fungi on several rice varieties in Kuningan, Tasikmalaya and Subang, West Java]. Retrieved from <http://repository.ipb.ac.id/jspui/bitstream/123456789/44597/1/A07dei.pdf>
- López-González, R. C., Gómez-Cornelio, S., De la Rosa-García, S. C., Garrido, E., Oropeza-Mariano, O., Heil, M., & Partida-Martínez, L. P. (2017). The age of lima bean leaves influences the richness and diversity of the endophytic fungal community, but not the antagonistic effect of endophytes against *Colletotrichum lindemuthianum*. *Fungal Ecology*, 26, 1-10. <http://doi.org/10.1016/j.funeco.2016.11.004>
- Mei, C., & Flinn, B. S. (2010). The use of beneficial microbial endophytes for plant biomass and stress tolerance improvement. *Recent Patents on Biotechnology*, 4(1), 81-95. <http://doi.org/10.2174/187220810790069523>
- Odum, E. P. (1993). *Dasar-dasar ekologi* [Fundamentals of Ecology]. (T. Samingan, Trans.). Yogyakarta: Gadjah Mada University Press. (Original work published 1953)
- R Core Team. (2013). *R: A language and environment for statistical computing*. Vienna, AT: R Foundation for Statistical Computing.
- Sanchez-Azofeifa, A., Oki, Y., Fernandes, G. W., Ball, R. A., & Gamon, J. (2012). Relationships between endophyte diversity and leaf optical properties. *Trees - Structure and Function*, 26(2), 291-299. <http://doi.org/10.1007/s00468-011-0591-5>

Aminudin Afandhi et al.: Diversity of Endophytic Fungi in Apple Leaves.....

- Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*, 27, 379–423. <http://doi.org/10.1145/584091.584093>
- Sieber, T. N. (2007). Endophytic fungi in forest trees: Are they mutualists? *Fungal Biology Reviews*. <http://doi.org/10.1016/j.fbr.2007.05.004>
- Tarno, H., Septia, E. D., & Aini, L. Q. (2016). Microbial community associated with ambrosia beetle, *Euplatypus parallelus* on sonokembang, *Pterocarpus indicus* in Malang. *AGRIVITA Journal of Agricultural Science*, 38(3), 312–320. <http://doi.org/10.17503/agrivita.v38i3.628>
- Thakur, A., Singh, V., Kaur, A., & Kaur, S. (2014). Suppression of cellular immune response in *Spodoptera litura* (Lepidoptera: Noctuidae) larvae by endophytic fungi *Nigrospora oryzae* and *Cladosporium uredinicola*. *Annals of the Entomological Society of America*, 107(3), 674–679. <http://doi.org/10.1603/AN13164>
- Tirtana, Z. Y. G., Sulistyowati, L., & Cholil, A. (2013). Eksplorasi jamur endofit pada tanaman kentang (*Solanum tuberosum* L.) serta potensi antagonismenya terhadap *Phytophthora infestans* (Mont.) de Barry penyebab penyakit hawar daun secara in vitro [Exploration of endophytic fungi in potato plants (*Solanum tuberosum* L.) and potential antagonism against *Phytophthora infestans* (Mont.) de Barry causes leaf blight disease in vitro]. *Jurnal HPT*, 1(3), 91-101. Retrieved from [http://download.portalgaruda.org/article.php?article=190870&val=6471&title=EKSPLORASI%20JAMUR%20ENDOFIT%20PADA%20TANAMAN%20KENTANG%20\(Solanum%20tuberosum%20L\)%20SERTA%20POTENSI%20ANTAGONISMENYA%20TERHADAP%20Phytophthora%20infestans%20\(Mont.\)%20de%20Barry%20PENYEBAB%20PENYAKIT%20HAWAR%20DAUN%20SECARA%20IN%20VITRO](http://download.portalgaruda.org/article.php?article=190870&val=6471&title=EKSPLORASI%20JAMUR%20ENDOFIT%20PADA%20TANAMAN%20KENTANG%20(Solanum%20tuberosum%20L)%20SERTA%20POTENSI%20ANTAGONISMENYA%20TERHADAP%20Phytophthora%20infestans%20(Mont.)%20de%20Barry%20PENYEBAB%20PENYAKIT%20HAWAR%20DAUN%20SECARA%20IN%20VITRO)
- Zakaria, L., Yaakop, A. S., Salleh, B., Zakaria, M., & Pinang, P. (2010). Short communication endophytic fungi from paddy. *Tropical Life Sciences Research*, 21(1), 101–107.
- Zheng, Y. K., Miao, C. P., Chen, H. H., Huang, F. F., Xia, Y. M., Chen, Y. W., & Zhao, L. X. (2017). Endophytic fungi harbored in *Panax notoginseng*: Diversity and potential as biological control agents against host plant pathogens of root-rot disease. *Journal of Ginseng Research*, 41(3), 353–360. <http://doi.org/10.1016/j.jgr.2016.07.005>