

Research Article

Comparing seeds germination of some local plant species on two hydroseeding mulches for post mining revegetation

M.F. Anshari^{1,*}, A.A.R. Fernandes², E. Arisoesilaningsih¹

¹Biology Master Program, ²Department of Statistics, Faculty of Mathematics and Natural Sciences, Brawijaya University, Jl. Veteran, Malang, East Java, Indonesia

*corresponding author: delansh@gmail.com

Received 06 December 2017, Accepted 20 December 2017

Abstract: The aims of this study were to determine seed germination rate of some local plant species in two hydroseeding mulches containing different tackifier concentration, as well as to determine the optimal hydroseeding mulch media composition for germinating seeds. This study used seeds of 13 local plant species: two species of Cyperaceae (*Cyperus brevifolius*, *C. javanicus*), five species of Leguminosae (*Cajanus cajan*, *Crotalaria pallida*, *Sesbania grandiflora*, *S. sesban*, *Tephrosia purpurea*), and six species of Poaceae (*Eleusine indica*, *Paspalum conjugatum*, *Sorghum timorense*, *S. bicolor*, *Sporobolus indicus*, *Themeda arundinaceae*). Two hydroseeding mulch media with different tackifier composition were mixed with seeds of each species and then sowed in pots. Each treatment was repeated three times. Moistened cotton wool was used as control and comparative media for observing seed viability. Seed germination in mulch media was observed during 13 days. The results showed that only 8 of 13 species could be germinated: *S. indicus*, *S. timorense*, *T. arundinaceae*, *C. cajan*, *C. pallida*, *S. grandiflora*, *S. sesban*, and *T. purpurea*. The highest germination rate was shown by *S. sesban* (67%) in M₂ medium and the lowest one was shown by *T. arundinaceae* (2%) in both media. The fastest germination time was recorded for *C. pallida* and *S. sesban* seeds that germinated in 2 days after sowing (DAS) in both media, while *S. timorense* and *T. arundinaceae* seeds showed the lowest ones in 11 DAS. The fluid M₁ medium was optimal for seeds germination of *S. sesban* (50%) and *S. grandiflora* (35%), while the thicker M₂ medium was optimal for seeds germination of *S. sesban* (67%) and *S. timorense* (50%) in 13 DAS. The maximum germination rate was generally reached in 11 DAS.

Keywords: seed, hydroseeding, germination rate, mulch

To cite this article: Anshari, M.F., Fernandes, A.A.R. and Arisoesilaningsih, E. 2018. Comparing seeds germination of some local plant species on two hydroseeding mulches for post mining revegetation. J. Degrade. Min. Land Manage. 5(2): 1103-1110, DOI: 10.15243/jdmlm.2018.052.1103.

Introduction

Indonesia is one of the countries with significant role in producing global mines such as coal, gold, nickel, copper, and tin, and is among the largest thermal coal exporting countries in the world. High mineral and coal deposits placed Indonesia as a country with high mining prospects (Devi and Prayogo, 2013; PwC Indonesia, 2017). However, mining exploitation practiced in Indonesia, especially by open mining techniques, resulted in massive alteration and degradation of physical, chemical and biological aspects of the

environment. To restore the function of degraded environments, land reclamation effort needs to include several stages; one of those is revegetation (Kilowasid et al., 2011; Sudarmadji and Hartati, 2016). Revegetation promotes some post mining development, creates aesthetically landscaping, and increases post-mining productivity (Skousen and Zipper, 2010). Selection of appropriate plant species, planting methods, and maintenance of vegetation after program, are necessary in the revegetation stages (Sudarmadji and Hartati, 2016). A used method in revegetation activities on degraded lands is hydroseeding (Albaladejo-

Montoro et al., 2000; Martinez-Ruiz et al., 2007; Skousen and Zipper, 2010). Hydroseeding is a method of land revegetation for stabilizing overburden surfaces and preventing soil erosion through spraying of mixed mulch fibers, seeds, fertilizers, and water on the soil surface. Mulch applications can reduce soil evaporation, maintain soil moisture, suppress weed growth, control temperature and soil structure, and affect soil microorganisms. Such environmental conditions will allow seeds germination (Cerenó et al., 2011; Cadre et al., 2017).

The use of local plants as seed sources is highly recommended in post mining land revegetation due to its ability to adapt to local biotic and abiotic conditions (Oliveira et al., 2013). Several previous greenhouse studies have demonstrated the successful germination of local plant seeds from Cyperaceae, Poaceae, and Leguminosae families in some hydroseeding mulch media containing natural gum. The seeds of *C. brevifolius*, *C. eragrotis*, *C. odoratus*, *C. strigosus*, *E. indica*, *Eulalia amaura*, *P. conjugatum*, *S. indicus*, *C. cajan*, *C. pallida*, *Desmodium triflorum*, *Indigofera spicata*, and *S. grandiflora* were able to germinate and grow on these media (Yulianingsih and Arisoesilaningsih, 2015; Baiti and Arisoesilaningsih, 2015; Rahma and Arisoesilaningsih, 2015; Azalia et al., 2016). However, the previous mulches were not resistant under heavy rain, the slurry tended to be disintegrated and finally seeds germinated in low rate. Moreover, research in laboratory, greenhouse, and field conditions conducted by Oliveira et al. (2012) indicated that the success of local seed germination in the field has a low frequency. In order to improve the mulch performance, the natural gum could then be replaced by degradable tackifier. So far, the influence of the components of the hydroseeding mulch media on the performance of local plant seeds germination has not been widely reported. The main purpose of this study was to determine seed germination rate of some local plant species in two hydroseeding mulches containing different tackifier concentration. Therefore, we could determine the optimal hydroseeding mulch media composition for the seed germination in the laboratory scale before trial in the field.

Materials and Methods

This research was conducted in the Laboratory of Ecology, Biology Department, Faculty of Mathematics and Natural Sciences, Brawijaya University, Malang. The study used seeds from some local plant species, i.e. two species of Cyperaceae (*C. brevifolius*, *C. javanicus*), five

species of Leguminosae (*C. cajan*, *C. pallida*, *S. grandiflora*, *S. sesban*, *T. purpurea*), and six species of Poaceae (*E. indica*, *P. conjugatum*, *S. bicolor*, *S. timorensis*, *S. indicus*, *T. arundinaceae*). Available seeds were soaked in water for 12 hours. The floating seeds were discarded and the submerged seeds were removed to be soaked in GA3 solution for 1 hour. Furthermore, the seeds that have been soaked in GA3 solution were washed. Hydroseeding mulch as seeds germination media was made from tackifier agent, water, compost, and fiber composition. Two variations of hydroseeding mulch media composition were used, the first mulch media with less tackifier agent content (M₁) and the second with twice of tackifier agent content (M₂). For seed viability test, moistened cotton wool was also used as a control medium.

The prepared seeds of each species were sown by mixing them into 30 grams of mulch. The total weight of used seeds of each species was different and depended on the seed size. Seeds of Cyperaceae, Poaceae, *C. pallida*, *S. sesban*, and *T. purpurea* were respectively used 0.1 grams. While seeds of *C. cajan* and *S. grandiflora* used were 1 gram. Three replications were performed on each species. Seeds germination in M₁ and M₂ media were observed for 13 days under room temperature (20.9 ± 0.5 °C) with 77.5 ± 1.9% of relative humidity and 23.7 ± 7.5 lux of light intensity conditions. Data were tabulated and analyzed using Ms. Excel and R software.

Results and Discussion

Germination rate among local species seeds

Generally, based on the seed viability test on moistened cotton wool media, it was known that the seeds of *C. brevifolius*, *C. javanicus*, *E. indica*, *P. conjugatum*, *S. bicolor*, *S. indicus*, *T. arundinaceae*, *C. cajan*, *S. grandiflora*, and *T. purpurea* had a low viability rate, as indicated by its germination rate of ≤ 25%. While *S. sesban* seeds showed 73% germinating seeds and considered had a moderate seed viability, then *S. timorensis* and *C. pallida* seeds showed a higher viability. Its germinating seeds reached ≥ 75%. Seed germination is a controlled process by a number of mechanisms required for the growth and development of embryos, with the end result is new individuals (Miransari and Smith, 2014). The low rate of seed viability could be resulted by a drastic fluctuation of humidity and temperature after storage condition that influences the seed moisture content (Shaban, 2013). Patil and Krishna (2016) reported that the moisture content of *Calamus thwaitesii* and *Calamus nagabettai*

seeds affected the germination rate. The seeds with higher moisture content showed higher germination rate.

Moreover, it was also recorded that the seeds germination at first days in mulch media was longer than in the moistened cotton wool media (Table 1). Seeds germination of each species in mulch showed different germination rate in 13 DAS (Table 1). The seeds of Leguminosae showed a higher germination rate and a faster germination time compared to the seeds of Cyperaceae and Poaceae. This supported a finding of Azalia et al. (2016) that the germination rate of Leguminosae seeds was

higher and the germination time was faster than the rate and time of germination of Poaceae and Cyperaceae seeds.

Seeds of *C. brevifolius* and *C. javanicus* of Cyperaceae did not show any germination in both mulch media until 13 DAS. The mean of germination time of Cyperaceae seeds in some media was 19 DAS (Azalia et al., 2016). A factor influencing the time of sedge seed germination is temperature. Cyperaceae grown in open habitats requires a higher temperature than plants from the same family growing in the forest area. The optimum temperature for sedges seed growth ranges from 20°C to 25°C (Żukowski et al., 2010).

Table 1. The seeds germination rate of some local plant species in hydroseeding mulch media

No	Species	Germination rate*(%)			First Days of Germination (DAS)		
		M ₁	M ₂	C	M ₁	M ₂	C
1	<i>C. brevifolius</i>	0	0	0	-	-	-
2	<i>C. javanicus</i>	0	0	0	-	-	-
3	<i>E. indica</i>	0	0	0	-	-	-
4	<i>P. conjugatum</i>	0	0	0	-	-	-
5	<i>S. bicolor</i>	0	0	6	-	-	3
6	<i>S. indicus</i>	6	3	16	4	4	2
7	<i>S. timorensis</i>	6	50	97	11	7	1
8	<i>T. arundinaceae</i>	2	2	0	11	11	-
9	<i>C. cajan</i>	0	20	13	-	3	2
10	<i>C. pallida</i>	10	2	87	2	2	1
11	<i>S. grandiflora</i>	35	19	20	4	4	2
12	<i>S. sesban</i>	50	67	73	2	2	1
13	<i>T. purpurea</i>	13	20	0	4	3	-

Note: M₁ = Mulch 1, M₂ = Mulch 2, C = Control, DAS = days after sowing, * = until 13 DAS

The seeds of Poaceae showed variable germination rates among species and experiments. Sown seeds of *E. indica*, *P. conjugatum*, and *S. bicolor* did not germinate in two mulches. This was different with the results reported by Azalia et al. (2016) that germination rate of *E. indica* reached 13% and *P. conjugatum* reached 4% in 10 DAS. Furthermore, a study conducted by Rahma and Arisoesilaningih (2015) showed that the fastest germination time of *E. indica* and *P. conjugatum* was in 3 DAS with the highest germination rate of 19% and 44% until 15 DAS. The possible factor influencing these variations was seed dormancy. As reported by Hu et al. (2014), the rate of seedling emergence from fresh seeds, that had non-deep physiological dormancy, was significantly lower than the seeds stored for 1 year. Another study conducted by Gasque and García-Fayos (2003) showed that the seedling emergence of the recently collected seeds (2% ± 2) was lower than those stored for two months (3% ± 4) and four months (32% ± 3). Seeds of *S.*

indicus, *S. timorensis*, and *T. arundinaceae* germinated at different times. The seeds of *S. indicus* initiated to germinate in 4 DAS, both in M₁ and M₂ media. However, the highest germination rate until 13 DAS was present in M₁ medium (6%), while seed germination in M₂ medium only reached 3%. The mean of seed germination starting time of *S. indicus* was 3 DAS with the highest germination rate of 58% in 15 DAS (Rahma and Arisoesilaningih, 2015). The first germination of *S. timorensis* seeds occurred in 7 DAS in M₂ medium, with germination rate of 50% in 13 DAS. However, in M₁ medium, seed germination occurred a week later (11 DAS) with a rate of 6% in 13 DAS. *Sorghum* is a sensitive plant to low temperature. A low temperature can decrease the rate of seed germination (Razmi et al., 2013). Meanwhile, first seed germination of *T. arundinaceae* occurred in 11 DAS, in both media, with the same seed germination rate (2%) until 13 DAS. Some environmental factors such as temperature, light, and humidity greatly affect the

ability of germination of Cyperaceae and Poaceae seeds. In addition, variations of mass and seed size also affect germination. There is a positive correlation between seed mass and germination index, and between seed mass and germination rate. Seeds with larger size have higher germination rate than seeds with smaller size (Maas, 1989; Wu and Du, 2007; Żukowski et al., 2010; Khan et al., 2014).

C. cajan seeds of Leguminosae just germinated in M₂ medium, in 3 DAS, with 20% of germination rate until 13 DAS. The similar result was reported by Azalia et al. (2016) that *C. cajan* seeds were able to germinate in all mulch media with the fastest germination time in 3 DAS and the mean of seed germination starting time in all media occurred in 7 DAS. Baiti and Arisoesilaningsih (2015) reported that seeds of *C. cajan* sown in mulch media with 0.5 cm depth did not germinate at all. Seed germination of *C. cajan* is hypogeal. Germinating seeds are able to emerge from a depth of up to 5 cm under an optimal germination temperature ranging from 29°C to 36°C (Singh and Oswalt, 1992). Seeds of *C. cajan* show a high promising food source or seeds for revegetation program, because of its optimal nutritional content and high tolerance to environmental stress. Furthermore, *C. cajan* has high biomass productivity and moisture contributions to the soil (Odeny, 2007). Species selection and numbers will depend on the seed availability and test plot results.

C. pallida seeds were able to germinate in both media, but the highest germination rate was in M₁ medium (10-16%) at 7-13 DAS (Figure 1). Seed germination rate in M₂ medium only reached 2%. It seemed that M₂ medium containing with more tackifier agent had lower porosity and aeration rate than M₁ medium. According to Purwantoro (2016), media with low porosity and aeration decreased the germination rate. This low germination rate could also be increased by soaking *C. pallida* seeds in hot water with 60°C temperature for 30 minutes (Kak et al., 2007). The time required for seeds to germinate in both media was the same, in 2 DAS. These results indicated that the germination time of *C. pallida* seeds in M₁ and M₂ media was a day faster than those in previous study conducted by Azalia et al. (2016), as well as Baiti and Arisoesilaningsih (2015).

The seeds of *S. grandiflora* and *S. sesban* germinated in both media. Seeds of *S. grandiflora* initiated to germinate in 4 DAS in both media with the highest germination rate in M₁ medium (35%) until 13 DAS. In M₂ medium, seed germination rate reached 19%. Meanwhile, *S. sesban* seeds in both media were able to germinate in 2 DAS. The highest rate of seed

germination was shown by M₂ medium (67%) until 13 DAS, whereas in M₁ medium, the highest seed germination rate (58%) occurred in 5-9 DAS. However, this rate decreased to 50% in 13 DAS (Figure 1). The enhancement of seed germination rate of *Sesbania* could be conducted by soaking the seeds in distilled water for 24 hours or in a solution of GA3 (250 mg/L) for 24 hours (Shreelalitha et al., 2015). Although the light can affect seed germination in general, the treatment of dark and light conditions does not show the difference in seeds germination rate of *S. sesban* as reported by Dan and Brix (2007). Otherwise, the temperature is very influential on seed germination. The optimal temperature for seed germination of *S. sesban* is 30°C to 37°C, with seed germination rate reaches 85%. While in 5°C and 13°C, *S. sesban* seed did not germinate at all.

Seeds of *T. purpurea* showed the highest germination rate in M₂ medium (20%) in 9-13 DAS with first germination time in 3 DAS, while in M₁ medium, the highest germination rate (16%) was observed in 7-9 DAS and decreased to 13% in 13 DAS. Seeds of *T. purpurea* in M₁ media initiated to germinate in 4 DAS. The germination capacity of *T. purpurea* seeds was significantly greater in the temperature range 30°C-35°C (Sy et al., 2001). Seeds germination rate of *T. purpurea* could be increased by soaking seeds in boiling water or in IAA and GA3 solutions. Seeds soaking treatment in boiling water (100°C) for 90 seconds increased the seeds germination rate compared to treatment in hot water (60°C) and normal water (Naikawadi, 2016).

Selecting optimal mulch medium

Comparing the germination rate of seeds in M₁ and M₂ media, based on the Mixed Model Analysis, specifically, both media were suitable for germination of *S. sesban* seeds with the highest rate among species. While *S. grandiflora* seeds germinated better in M₁ medium than in M₂ medium. Otherwise, germination of *S. timorensis* seeds was better in M₂ medium. Overall, M₂ medium was better than M₁ medium for seed germination (Figure 2). Although M₂ medium had lower porosity and aeration rate than M₁ medium, its more tackifier agent content was expected to decrease the moisture content by absorbing the water.

Matsushima and Nakagami (2013) reported that germination media with moderate moisture content (8%-11%) gave faster seed germination time than those the higher. The medium with moderate water holding capacity was better for seed germination and the most suitable range of medium water content for the process of seedling emergence and growth was about 12%. Either low

or excessive media water content would reduce the ability of seedling emergence and growth (Gairola et al., 2011; Tang et al., 2016). The analysis of all seeds germination rate in both media using Cubic Equation (Table 2) showed equation as follow,

$y = -0.0217x^3 + 0.2869x^2 + 1.6164x - 2.5073$
 from this equation, it was known that the maximum germination rate of all seeds in both media occurred in 11 DAS (Figure 3).




Figure 1. Germination rate of some local plant species until 13 DAS in both media. Descriptions: A) *S. indicus*, B) *C. cajan*, C) *S. grandiflora*, D) *S. sesban*, E) *T. arundinaceae*, F) *C. pallida*, G) *T. purpurea*, H) *S. timorensis*, GR = germination rate, DAS = days after sowing, M₁ = Mulch 1, M₂ = Mulch 2.

Table 2. Model summary and parameter estimates

No	Equation	Model Summary					Parameter Estimates			
		R Square	F	df1	df2	Sig.	Constant	b1	b2	b3
1	Logarithmic	0.921	128.1	1	11	0.000	-4.0	9.8		
2	Inverse	0.677	23.0	1	11	0.001	19.2	-25.1		
3	Quadratic	0.975	197.5	2	10	0.000	-6.1	4.3	-0.2	
4	Cubic	0.988	256.3	3	9	0.000	-2.5	1.6	0.3	0.0
5	Compound	0.397	7.2	1	11	0.021	0.1	1.7		
6	Power	0.685	23.9	1	11	0.000	0.0	3.6		
7	S	0.936	159.7	1	11	0.000	4.6	-12.5		
8	Growth	0.397	7.2	1	11	0.021	-2.2	0.5		
9	Exponential	0.397	7.2	1	11	0.021	0.1	0.5		
10	Logistic	0.397	7.2	1	11	0.021	9.3	0.6		

Note: Dependent variable = germination rate, Independent variable = days after sowing




Figure 2. Mixed model of seeds germination.

Descriptions: A. Seeds germination in M₁ medium, B. Seeds germination in M₂ medium, C. Comparison between seeds germination in M₁ and M₂ media, Cp = *C. pallida*, Sg = *S. grandiflora*, Si = *S. indicus*, Ss = *S. sesban*, St = *S. timorensis*, Ta = *T. arundinaceae*, Tp = *T. purpurea*, DAS = days after sowing, M₁ = Mulch 1, M₂ = Mulch 2.




Figure 3. Model of all species seeds germination rate in both mulch media
 Descriptions: GR = germination rate, DAS = days after sowing.

Conclusion

Seeds of *S. indicus*, *S. timorensis*, *T. arundinaceae*, *C. pallida*, *S. grandiflora*, *S. sesban*, and *T. purpurea* were able to germinate in M₁ and M₂ media, while *C. cajan* only germinated in M₂ medium. Seeds of *C. brevifolius*, *C. javanicus*, *E. indica*, *P. conjugatum*, and *S. bicolor* did not show any germination. The highest seed germination rate was observed on *S. sesban* (67%) in M₂ medium and the lowest seed germination rate was shown by *T. arundinaceae* (2%) in both media. M₁ medium was optimal for seed germination of *S. sesban* and *S. grandiflora*, while M₂ medium was optimal for seed germination of *S. sesban* and *S. timorensis*. The maximum germination rate occurred in 11 DAS.

Acknowledgements

This study was carried out under valuable support of Dr. Catur Retnaningdyah as the Head of Ecology and Animal Diversity Laboratory, as well as the reviewers of Research Group of Biotechnology for Conserving Sustainably Tropical Ecosystem. Our sincere gratitude goes to these amazing individuals for their support throughout the study. We also acknowledge Mr. Purnomo as Laboratory Technical Assistant and Miss Tiara Putri Anggini for the contributions.

References

Albaladejo-Montoro, J., Alvarez-Rogel, J., Querejeta, J., Diaz, E. and Castillo, V. 2000. Three hydroseeding revegetation techniques for soil erosion control on anthropic steep slopes. *Land Degradation and Development* 11:315-325.

- Azalia, D., Retnaningdyah, C. and Arisoelaningsih, E. 2016. Germination of seeds of some local pioneer plant species in different hydroseeding mulches for revegetation of post-coal mining soil. *Journal of Degraded and Mining Lands Management* 3(4):609-615.
- Baiti, R.N. and Arisoelaningsih, E. 2015. Reclamation of post coal mining using hydroseeding involving seeds of some local Papilionaceae. *Jurnal Biotropika* 3(1):36-41.
- Cereno, M.M., Tan, F.J. and Uy, F.A.A. 2011. Combined hydroseeding and coconut reinforcement for soil erosion control. *Soil Erosion Studies*. InTech.
- Dan, T.H. and Brix, H. 2007. The influence of temperature, light, salinity and seed pre-treatment on the germination of *Sesbania sesban* seeds. *African Journal of Biotechnology* 6(19):2231-2235.
- Devi, B. and Prayogo, D. 2013. Mining and development in Indonesia: an overview of the regulatory framework and policies. Action Research Report. International Mining for Development Center, Brisbane.
- Gairola, K.C., Nautiyal, A.R. and Dwivedi, A.K. 2011. Effect of temperature and germination media on seed germination of *Jatropha curcas* Linn. *Advances in Bioresearch* 2(2):66-71.
- Gasque, M. and García-Fayos, G. 2003. Seed dormancy and longevity in *Stipa tenacissima* L. (Poaceae). *Plant Ecology* 168:279-290.
- Hu, X.W., Wu, Y.P., Ding, X.Y., Zhang, R., Wang, Y.R., Baskin, J.M. and Baskin, C.C. 2014. Seed dormancy, seedling establishment and dynamics of the soil seed bank of *Stipa bungeana* (Poaceae) on the Loess Plateau of Northwestern China. *PLoS One* 9(11): e112579.
- Kader, M.A., Senge, M., Mojid, M.A. and Ito, K. 2017. Recent advances in mulching materials and methods for modifying soil environment. *Soil and Tillage Research* 168:155-166.

- Kak, A., Devi, L.C., Gupta, V. and Singh, N. 2007. Response of seed treatments on seed germination in wild *Crotalaria* species. *Acta Horticulturae* 752:261-265.
- Khan, A.R., Khan, M.J., Yousaf, A., Baloch, N.U., Fatima, M. and Shaheen, G. 2014. Seed size variation and its effect on seed germination and growth rate in Poaceae. *International Journal of Pharmaceutical Sciences Review and Research* 26(1):6-10.
- Kilowasid, L.M.H., Herlina, Syaf, H., Safuan, L.O., Tufaila, M., Leomo, S. and Widiawan, B. 2015. Engineering of soil biological quality from nickel mining stockpile using two earth worm ecological groups. *Journal of Degraded Mining Lands Management* 2(3):361-367.
- Maas, D. 1989. Germination characteristics of some plant species from calcareous fens in southern Germany and their implications for the seed bank. *Holarctic Ecology* 12:337-344.
- Martinez-Ruiz, C., Fernandez-Santos, B., Putwain, P.D. and Fernandez-Gomez, M.J. 2007. Natural and man-induced revegetation on mining wastes: changes in the floristic composition during early succession. *Ecological and Engineering* 30:286-294.
- Matsushima, K. and Sakagami, J. 2013. Effects of seed hydropriming on germination and seedling vigor during emergence of rice under different soil moisture conditions. *American Journal of Plant Species* 4:1584-1593.
- Miransari, M. and Smith, D.L. 2014. Plant hormones and seed germination. *Environmental and Experimental Botany* 99:110-121.
- Naikawadi, V.B. 2016. Seed germination of important medicinal plant *Tephrosia purpurea* (Linn.), Pers. *International Research Journal of Natural and Applied Sciences* 3(8):186-207.
- Odeny, D.A. 2007. The potential of pigeonpea (*Cajanus cajan* (L.) Millsp.) in Africa. *Natural Resources Forum* 31:297-305.
- Oliveira, G., Clemente, A., Nunes, A. and Correia, O. 2013. Limitations to recruitment of native species in hydroseeding mixtures. *Ecological Engineering* 57:18-26.
- Oliveira, G., Nunes, A., Clemente, A. and Correia, O. 2012. Testing germination of species for hydroseeding degraded Mediterranean areas. *Restoration Ecology* 20(5):623-630.
- Patil, S.S. and Krishna, A. 2016. Influence of seed moisture content on seed germination and quality in Canes. *Journal of Plant Science and Research* 3(2):156.
- Purwanto, R.S. 2016. Short communication: effect of growing media on seed germination and seedling growth of *Aganope heptaphylla* (Leguminosae). *Nusantara Bioscience* 8(2):150-154.
- PwC Indonesia, 2017. Mining in Indonesia: investment and taxation guide. PricewaterhouseCoopers Indonesia. <https://www.pwc.com/id/en/energy-utilities-mining/assets/May%202016/PwC%20Indonesia-mining-in-Indonesia-survey-2016.pdf>. Accessed on 18th October 2017.
- Rahma, A.F. and Arisoelaningsih, E. 2015. Growth of pioneer grasses in monoculture and polyculture systems of hydroseeding applied in a coal mining tailing from the South Kalimantan. *Jurnal Biotropika* 3(2):33-36.
- Razmi, Z., Hamidi, R. and Pirasteh-Anosheh, H. 2013. Seed germination and seedling growth of three sorghum (*Sorghum bicolor* L.) genotypes as affected by low temperatures. *International Journal of Farming and Allied Sciences* 2(20):851-856.
- Shaban, M. 2013. Study on some aspects of seed viability and vigor. *International Journal of Advanced Biological and Biomedical Research* 1(12):1692-1697.
- Shreelalitha, S.J., Sridhar, K.R. and Suresh, S. 2015. Seed dormancy and germination in two wild genotypes of *Sesbania* of the southwest mangroves in India. *International Journal of Agricultural Technology* 11(4):895-902.
- Singh, F. and Oswald, D.L. 1992. Pigeonpea botany and production practices. International Crops Research Institute for the Semi-Arid Tropics. Patancheru, Andhra Pradesh, India.
- Skousen, J. and C.E. Zipper. 2010. Revegetation species and practices. Virginia Cooperative Extension 460-122.
- Sudarmadji, T. and Hartati, W. 2016. The process of rehabilitation of mined forest lands toward degraded forest ecosystem recovery in Kalimantan, Indonesia. *Biodiversitas* 17(1):185-191.
- Sy, A., Grouzis, M. and Danthu, P. 2001. Seed germination of seven Sahelian legume species. *Journal of Arid Environments* 49(4):875-882.
- Tang, J., Busso, C.A., Jiang, D., Wang, Y., Wu, D., Musa, A., Miao, R. and Miao, C. 2016. Seed burial depth and soil water content affect seedling emergence and growth of *Ulmus pumila* var. *sabulosa* in the Horqin Sandy Land. *Sustainability* 8(68):1-10.
- Wu, G. and Du, G. 2007. Germination is related to seed mass in grasses (Poaceae) of the eastern Qinghai-Tibetan Plateau, China. *Nordic Journal of Botany* 25:361-365.
- Yulianingsih, D. and Arisoelaningsih, E. 2015. Application of some hydroseeding mulches for pioneer sedge seeds germination in post mining soil from South Kalimantan. *Jurnal Biotropika* 3(1):27-31.
- Żukowski, W., Bogdanowicz, A.M. and Lembicz, M. 2010. Seed germination in sedges: a short review. *Biodiversity Research and Conservation* 19:15-22.