

BULLETIN OF THE MARINE GEOLOGY

VOLUME 31 • NUMBER 1 • JUNE 2016



BULL. MGI

VOL. 31

NO. 1

PAGE 1 - 53

BANDUNG, JUNE 2016

ISSN 1410-6175

Accredited : LIPI No. 665/AU2/P2MI-LIPI/07/2015



MARINE GEOLOGICAL INSTITUTE
RESEARCH AND DEVELOPMENT AGENCIES FOR ENERGY AND MINERAL RESOURCES
MINISTRY OF ENERGY AND MINERAL RESOURCES

PUSAT PENELITIAN DAN PENGEMBANGAN GEOLOGI KELAUTAN
BADAN PENELITIAN DAN PENGEMBANGAN ENERGI DAN SUMBERDAYA MINERAL
KEMENTERIAN ENERGI DAN SUMBERDAYA MINERAL

BULLETIN OF THE MARINE GEOLOGY

Vol. 31, No. 1, June 2016

INSURED EDITOR

Director of Marine Geological Institute

VICE CHIEF OF INSURED EDITOR

Head of Afiliation Division

CHIEF OF EDITORIAL BOARD

Dr. Ir. Noor Cahyo D. Aryanto, MT.

VICE CHIEF OF EDITORIAL BOARD

Ir. Yudi Darlan, M.Sc.

EDITORIAL BOARDS

Dr. Ir. Ediar Usman, MT.

Dr. Ir. Hananto Kurnio, M.Sc.

Ir. I Wayan Luga

Kris Budiono, M.Sc.

Dr. Priatin Hadi Widjaya, ST., MT.

Dr. Ir. Dicky Muslim, M.Sc.

SCIENTIFIC REVIEWERS

Prof. (Ris) Dra. Mimin Karmini

Dr. Ir. Haryadi Permana

Lili Sarmili, M.Sc.

PUBLISHER BOARDS

Drs. Dwi Agus, HS. (*Chief*)

Franto Novico, ST, MSc.

Luli Gustiantini, ST, MT.

Hendro Dwi Bayu Abadi, S.Sos.

Sutisna

For communication of this publications, please contact :

MARINE GEOLOGICAL INSTITUTE

Dr. Junjunan 236, Bandung-40174, Indonesia

Telephone : +62-22-6032020, 6032201, Fax : +62-22- 6017887

E-mail : bulletin@mgi.esdm.go.id

Cover Figure : Crystalline limestone outcrops around of Baginda Cape, Tuboali Coast, Bangka Belitung Province
(By: Noor CD Aryanto)

PREFACE

Marine Geological Institute of Indonesia (MGI's) responsibilities are to provide marine geoscientific map, research and information to support sustainable development of Indonesian's mineral and petroleum industries, mapping of Indonesian Coastal and Ocean Territory, identification of marine and coastal geological hazards, and to provide marine and coastal geological and geophysical data base for marine and coastal landscape.

In this first edition of year 2016, the number of important information are highlighted involving: Concentration and Distribution of Polycyclic Aromatic Hydrocarbons (PAHS) During Bioremediation Processes of Oil-contaminated Beach Sediments in Karang Song Beach, Indramayu; Shallow Gas Features Based on Interpretation of Bottom Profiling Records at Topang Delta, Meranti Regency, Riau Province; The Mechanism of Sediment Depositional Environment of Core Drilling of Gilimanuk Coast, Bali and Ketapang, East Java, Based on Sediment Textures; Interpretation of Paleo-Channel Based on Shallow Seismic Reflection Record in Banten Bay, Banten Province; The Content of Placer Heavy Mineral and Characteristics of REE at Toboali Coast and Its Surrounding Area, Bangka Belitung Province. From the desk of editors, thank to the authors who contribute their valuable papers for the readers.

Editors

BULLETIN OF THE MARINE GEOLOGY

Vol. 31, No. 1, June 2016

CONTENTS

<i>Concentration and Distribution of Polycyclic Aromatic Hydrocarbons (PAHS) During Bioremediation Processes of Oil-contaminated Beach Sediments in Karang Song Beach, Indramayu</i> Khozanah and Dede Falahudin	1-10
<i>Shallow Gas Features Based on Interpretation of Bottom Profilling Records at Topang Delta, Meranti Regency, Riau Province</i> Purnomo Raharjo, Andrian Willyan Djaja and Ediar Usman.....	11-20
<i>The Mechanism of Sediment Depositional Environment of Core Drilling of Gilimanuk Coast, Bali and Ketapang, East Java, Based on Sediment Textures</i> Ediar Usman	21-33
<i>Interpretation of Paleo-Channel Based on Shallow Seismic Reflection Record in Banten Bay, Banten Province</i> Yogi Noviadi	35-43
<i>The Content of Placer Heavy Mineral and Characteristics of REE at Toboali Coast and Its Surrounding Area, Bangka Belitung Province</i> Noor Cahyo D. Aryanto and Udaya Kamiludin	45-53

Interpretation of Paleo-Channel Based on Shallow Seismic Reflection Record in Banten Bay, Banten Province

Penafsiran Alur Purba Berdasarkan Rekaman Seismik Pantul Dangkal di Teluk Banten, Provinsi Banten

Yogi Noviadi

Marine Geological Institute, Ministry of Energy and Mineral Resources, Jl. Dr. Djunjunan No. 236 Bandung 40174

Corresponding author : yoginoviadi@yahoo.com

(Received 08 December 2015; in revised from 21 December 2015; accepted 10 May 2016)

ABSTRACT: The objective of this study is to find out the pattern of paleo channel which was formed in Banten Bay and its surrounding. The aims are to find out the paleo-channel pattern at study area. The study methods are including vessel positioning, and shallow seismic reflection work. Vessel positioning method is to locate the exact position of seismic work when recording the data from single channel of shallow seismic reflection. Seismic line orientations are determined by regional geological setting of the area. Trend of seismic lines are dominantly north – south. In order to get the seismic data which could give geological setting configuration, seismic lines should be perpendicular to the strikes of the sediments.

Based on the calculation of velocity of seismic refraction in sea water 1,500 meters/second, while within sediment 1,600 meters/second, it could be concluded that the paleo channels were more or less in 32 meters below sea floor depth.

This layer was the system that occur during the process of an interglacial on the Sunda Shelf when it was still a part of land that connects the Java, Sumatra and Kalimantan Islands. Paleo-channel deposits are characterized by subparallel - chaotic reflection character with a thickness between 5-35 meters.

Keywords: Paleo-channels, seismic records and Banten Bay

ABSTRAK: Maksud dari penelitian ini adalah untuk mengetahui pola sungai purba yang terdapat di Teluk Banten dan sekitarnya, yang tujuannya adalah untuk mengetahui pola penyebaran alur sungai purba di daerah penelitian. Metode penelitian terdiri dari penentuan posisi kapal dan penelitian seismik pantul dangkal. Penentuan posisi kapal berguna untuk menemukan posisi yang tepat saat merekam data oleh peralatan seismik saluran tunggal dangkal. Lokasi lintasan seismik disesuaikan dengan kondisi geologi daerah penelitian. Arah lintasan seismik pada umumnya berarah utara – selatan. Untuk mendapatkan data seismik yang bisa memberikan konfigurasi kondisi geologi, lintasan seismik harus tegak lurus terhadap kedudukan lapisan batuan.

Berdasarkan cepat rambat gelombang seismik di air laut 1.500 meter/detik, dan sedimen 1.600 meter/ detik, dapat disimpulkan bahwa alur purba kurang lebih berada pada kedalaman 32 meter di bawah dasar laut.

Lapisan ini merupakan sistem pengendapan yang terjadi selama proses interglasial di Paparan Sunda yang pada saat itu masih merupakan bagian dari daratan yang menghubungkan P. Jawa, Sumatera dan P. Kalimantan. Endapan alur purba dicirikan dengan pola refleksi subparalel sampai tidak beraturan dengan ketebalan antara 5-35 meter.

Kata kunci: Alur purba, rekaman seismik dan teluk Banten

INTRODUCTION

The objective of this study is to find out of Paleo-Channel which were formed in Teluk Banten Waters. The aims are to conclude the development of Paleo Channel, hopefully the result of this study would be useful as a database for various needs such as for study and other development in the future.

Administratively, Banten Bay is part of Serang Regency, Banten Province, and geographically is

situated at 106°00'–106°25' E and 05°45' – 06°05' S. The study area is about 1,700 Km² (Figure 1).

In the land area are usually intermontane basin rivers flowing in the valleys of these rivers. The rivers to supply of sediment so that it is possible to sedimentation in these valleys. In addition to the supply of sediment from the rivers that time, sedimentation occurs when the sea level rises relatively quickly over a period of 18,000 years sea levels rose about 140 meters high, the valleys will first be inundated and also




Figure 1. Study area location

experienced sedimentation. It can be seen on a map the thickness of sediment that is the areas that have a large sediment thickness (WU Chen., 1991).

To determine the sub surface configuration form of Paleo-channel is required seismic methods to explore the ancient morphological. The seismic reflection is one method of exploration which based on the measurement of the response sound wave propagates in a layer boundary and then reflected and refracted from all the difference sedimentary rocks.

Needs of marine geophysical data shows an increasing trend due to the more widespread exploration of mineral resources and energy in the ocean. One method that is powerful enough to meet the needs is seismic reflection. This method has a high accuracy to know the characteristics of the sub-marine, such as the thickness and volume of sediment deposition, sea level, the structure of the seabed, and the depth of the waters (Susilawati, 2004). Basic skills in presentation of high-resolution information with a relatively simple operation, so that this method is often used in geological study.

Based on Santosa, et al (1982), the general geological condition of study area consist of Marikangan Volcanic Rock, Product of Gede Volcano, Banten Tuff, Gede Volcanic Rock, Pinang Mt. Basalt and Alluvium Deposits (Figure 2).

Concerning to the location of seismic reflection survey, the rock unit that will influent the result of seismic survey are Banten Tuff and Alluvium Deposits.

Banten Tuff is divided into lower Banten Tuff and upper Banten Tuff. Lower Banten Tuff consist of tuff breccias, agglomerate, pumiceous tuff, lapilli tuff and sandy tuff.

Tuff breccias, is composed by the clasts are made up of sand to bomb size, subangular to subrounded; composed of basalt, andesite, pumice, obsidian with very fine pyroclastic groundmass, thickness is several meters. The bomb clasts are scattered and at limited numbers (Santosa, et al, 1982).

Agglomerate, clast are made up of lapilli to bomb size, well rounded to subrounded; composed of basalt, andesite, pumice with sandy or fine clastic tuff matrix; as small intercalations in the volcanic breccias, the thickness is about several meters. The exposes are found at the upper course of the Anyer River and on the northeastern escarpment of Danau Caldera (Santosa, et al, 1982)

Pumiceous tuff, dirty white to gray, clasts make up gravel to sand size, sub rounded; composes of dominantly pumice, basalt, andesite and obsidian; loose and weathered; the thickness of the layer is about several centimeters.

METHODS

The methods are including vessel positioning, sounding and shallow seismic reflection. Trackline positioning method is to locate the exact position of the survey vessel when recording the data from single




Figure 2. The geological map of Anyer Quadrangle (modified from Santosa et all, 1982)

channel of shallow seismic reflection by using GPS (Global Positioning System) devices.

In practice however, the reflection seismic technique is mostly - complex because the echoes (reflected energy or seismic events) of interest are noised by both coherent and random. To compensate, sophisticated acquisition and processing methods have been developed to enhance the relative amplitudes of the reflected seismic events of interest. Many of these methodologies are site and target dependent. The interpretation of reflection seismic data is also complex, and as much an art as a science. Interpreted velocity/depth models can be unreliable because of either inaccurate velocity control or incorrect seismic event identification. Similarly, seismic amplitudes can be misinterpreted because of attenuation and improperly applied gain control. (Anderson N and Akingbade A, 1995)

The success of continuous marine seismic profiling methods are sitedependent but have the potential to produce high resolutionrecords in shallow water (Haeni, 1986, 1988)

The interpretation is based on the seismic stratigraphy interpretation (Mitchum et al, 1977 a and 1977b). Its objective is to define the genetic reflection

packages by the surfaces that envelope seismic sequence and system tracts. These bounding discontinuities are identified on the basis of reflection termination patterns and their continuity.

Boundaries are defined on a seismic line by identifying the termination of seismic reflectors at the discontinuity surfaces.

Seismic lines and location is determined by regional geological setting of the area. Trend of seismic lines are dominantly north – south. In order to get the seismic data which could give geological setting configuration, sismic lines should be perpendicular to the strikes of the sediments (Figure 3).

The drawing of seismic horizon was based on the criteria proposed by Ringis (1986). The assumption wave velocity has proposed that all horizons of seismic is 1,500 meters / seconds.

Data from the seafloor depth measurements of analog data, namely tide correction to get the value of the actual sea depth. Contouring process bathymetric data is done using Surfer software version 8.0 which then produce bathymetric contours. Furthermore, bathymetric map-making is done by using the Mapinfo program. While data analog recording is used to look at the cross section of seabed morphology more clearly, a




Figure 3. The location of seismic line (modified from Hadikusumo, S et.al, 1988)

good cross section perpendicular to the shoreline as well as a cross section parallel to the shoreline.

RESULTS

More than 200 km seismic reflection line was carried out and based on the analog recording data, the sub-surface geological condition of study area can be explained as follows:

Sequence A is the most upper layer which is characterized by strong reflector, parallel to sub parallel, high amplitude and continuous. This sequence can be seen at depth about 20 – 70 m below sea level (Figure 4, 5, 6 and 7). Based on the reflector character, the upper most layers can be interpreted as a fine grained sediment (clay, silt and mud) which was deposited as a near shore deposit. The morphology where this unit was deposited is characterized by flat to submarine hill undulation. Below sequence A is sequence B which were dominated by concave bedding form which is characterized by wavy sub-parallel to parallel and most of them have tranparent reflector, low amplitude, weak and uncontinuous reflector. Based on the characteristic of reflector, this unit probably is characterized by undifferentiated sediments and interpreted as sub-marine paleo-river environment. The biggest channel which can be found in this area is about 4 – 5 km (Figure 4, 5, 6 and 7). Below sequence B is sequence B1. This sub sequence is characterized by medium to strong reflector, high amplitude, sub parallel

– parallel and wavy reflector. This sequence shows as a bottom part of a big submarine channel and probably have a differences lithology character with the upper part channel. Sequence C is overlain by sequence A and B1. This sequence is dominated by strong reflector, high amplitude, sub parallel – parallel and wavy. Based on the regional geology condition, this sequence is assumed as coastal or fluatile deposit. With the ages about Upper Plistocene to Holocene. The lower most sequence is sequence D (seismic basement) which is characterized by strong reflector, high amplitude, subparallel – parallel, wavy and continuous. The upper boundary of this sequence is dominated by wavy undulated morphology and is catagoried as erosional truncation.

Based on the regional geological condition of the study area, sequence D can be classified as Plistocene volcanic product.

DISCUSSIONS

The seismic sections clearly show the characteristic curved geometries of classic cut-and-fill or channel features (Figure 4). From the morphology feature can be identified at least two channel features, each approximately 2 to 4 km wide with one slightly offset yet superimposed on part of the other. According to seismic interpretation shows that the paleochannel geomorphology with respect to interbed sequences and its characteristic variability.




Figure 4. Original and interpretation of seismic reflection record of line B-1




Figure 5. Original and interpretation of seismic reflection record of line B-2 (Upper original and bellow interpreted record)




Figure 6. Original and interpretation of seismic reflection record of line B-3.




Figure 7. Original and interpretation of seismic reflection record of line B-4




Figure 8. Bathymetric map of study area.

From the bathymetric mapping survey results, shows the seabed relatively flat with an average a depth of 20 m, where also found the shape of the sea floor that resembles with some valleys groove formed (closures) and forming a lineament trending northeast - southwest. The valleys as a small notches constitute grooves strait within Kopo Cp. and Porong Cp. The layout of this valley groove near the mouth of the river towards Porong Cp. that assumed, flow channel river valley which has been inundated by sea level rises (Figure 8).

Based on the interpretation of shallow seismic reflection which is taken in the North Serang waters, that found a layer of sea sand deposited in paleo channel morphology. This layer is the system that occur during the an interglacial process on the Sunda Shelf is still a part of land that connecting among the Java, Sumatra and Kalimantan Islands. Paleo-channel deposits are characterized by reflection character subparallel - chaotic with a thickness of between 5-35 meters.

Based on the fence diagram of several seismic line, shows that the direction of the channel is east-west (Figure 9).

CONCLUSION

The water depths in study area range from 5 to 20 meters in the southern part of Tunda Island. Furthermore, to the north of the study area to Tunda Island the depths reach 50 meters.

The study area is a part of the Sunda Shelf which connecting Java, Sumatra and Kalimantan Islands that influenced by inter-glacial process. Sand deposits are characterized by reflection character subparallel - chaotic with thicknesses between 5-35 meters.

Sequence A is interpreted dominated by clay which in some areas containing lenses of fine sands, mollusc shells, and carbonate material while Sequence B is interpreted as sedimentary layers of sand. Contact between A and B sequences is erosional truncation and downlap. Distribution of paleochannel which be indicated containing sand is occurred on sequence B.

ACKNOWLEDGEMENT

With the completions of this paper, the writer gives a gratitude to the Director of Marine Geological Institute. Thanks alot also to my colleagues for supporting to finalize this paper.

REFERENCES

- Anderson N and Akingbade A., 1995, Overview of the Shallow Seismic Reflection Technique Geophysical Atlas for Kansas: Kansas Geological Survey bulletin 237
- Haeni, F.P., 1986, Use of continuous seismic reflection methods in a hydrologic study in Massachusetts, a case study, in National Water Well Association Conference on




Figure 9. Fence diagram of seismic record

Surface and Borehole Geophysical Methods and Ground Water Instrumentation, Denver, Colorado, October 15- 17, 1986, Proceedings: Worthington, Ohio, National Water Well Association, p. 381-395.

Haeni, F.P., 1988, Evaluation of the continuous seismic refraction method for determining the thickness and lithology of stratified drift in the glaciated northeast, in A.D. Randall and A.I. Johnson, eds., Regional aquifer systems of the United States-The northeast glacial aquifers: American Water Resources Association monograph 11, p. 63-82.

Hadikusumo, S., Sarmili, L., Silitonga, F., Kurnio H., Hakim S., 1988. Laporan Penyelidikan Geologi dan Geofisika Kelautan Banten dan Sekitarnya, Pusat Penelitian dan Pengembangan Geologi Kelautan (Unpublished report)

Mitchum, R.M., Vail, P. R., Jr., and Sangree, J.B., 1977a, Seismic stratigraphy and global changes of sea level; Part 6, Seismic interpretation of seismic reflection patterns in depositional sequences, in Payton, C.E., ed., Seismic Stratigraphy—Applications to Hydrocarbon Exploration: American

Association of Petroleum Geologists Memoir 26, p.117-133.

Mitchum, R.M., Vail, P.R., Jr., and Thompson, S., III, 1977b, Seismic stratigraphy and global changes of sea level; Part 2, The depositional sequence as a basic unit for stratigraphic analysis, in Payton, C.E., ed., Seismic Stratigraphy—Applications to Hydrocarbon Exploration: American Association of Petroleum Geologists Memoir 26, p. 53-62.

Ringis, J., 1986. *Seismic Stratigraphy In Very High Resolution Shallow Marine Seismic Data*. Proceedings of the Joint ASCOPE/CCOP Workshop I, 119 - 128

Susilawati, 2004. Seismik refraksi (dasar teori dan akuisisi data). Fakultas Matematika dan Ilmu Pengetahuan Alam, Jurusan Fisika, Universitas Sumatera Utara. Medan. 50 hlm.

WU Chen, 1991. *Study of Paleochannels on The North China Plain* [M]. Beijing: China Science and Technology Press, 172. (in Chinese)

Guide for Authors - Geoscience Publications

Bulletin of the Marine Geology

MANUSCRIPTS

1. Manuscripts should be written in English, occasional contribution in Bahasa Indonesia will be considered for publication. This bulletin is a medium for the publication of original of marine geology, geophysics, geochemistry and related subjects.
2. The original and two copies of the manuscripts should be submitted for review purposes.
3. Manuscripts should be typewritten and doubled space. First lines of each new paragraph should be indented.
4. The entire manuscript should be paginated starting with the title page.
5. Only words to be set in italics should be underlined.
6. Manuscripts should in general be organized in the following order :
 - a. Title
 - b. Name(s) and affiliation(s) of author(s)
 - c. Abstract (not more than 500 words) in two languages English and Bahasa Indonesia.
Articles in English should start with an abstract in English followed an abstract by Bahasa Indonesia.
The abstract must be an informative statement of the content of the paper, explaining what the problem is, the method used, a statement of the results and the main conclusions. Constructions using phrases such as "this paper discusses", "are described" and "is reported" have no place in an abstract : by reading the abstract, the reader should be able to understand the essential qualities of the paper without referring to the paper itself.
 - d. Introduction
 - e. Methods, techniques, material studies. and area descriptions.
 - f. Results
 - g. Discussion
 - h. Conclusions
 - i. Acknowledgements
 - j. References : the following system should be used for arranging references.
 - *For periodicals*
Katili. J.A., 1978, Past and present geotectonics position of Sulawesi, Indonesia. *Tectonophysics*, 45 : 289-322.
 - *For edited symposia, special issues, etc., published in periodical*
Silitonga. P.H., Pudjowaluyo, H. and Molat, H., 1981, Geological Reconnaissance and mineral prospecting on Bacan Island (Mollucas, Indonesia). In: A.J. Barber and S. Wiryosujono (Editors), *The Geology and Tectonic of Eastern Indonesia*, Pergamon Press : 373-381.
 - *For Books*
Bemmelen. R.W van, 1949, *The Geology of Indonesia*. Netherlands Govt. Printing Office. The Hague, 997p.

Submission of electronic text


In order to publish the paper, authors are encouraged to submit the manuscript on CDROM or by email.



LEMBAGA
ILMU PENGETAHUAN
INDONESIA

**P2
MI**

Panitia
Penilai
Majalah
Ilmiah



SERTIFIKAT

Nomor: 665/AU2/P2MI-LIPI/07/2015

Akreditasi Majalah Ilmiah

Kutipan Keputusan Kepala Lembaga Ilmu Pengetahuan Indonesia
Nomor 818/E/2015 Tanggal 15 Juli 2015

Nama Majalah : Bulletin of the Marine Geology
ISSN : 1410-6175
Redaksi : Pusat Penelitian dan Pengembangan Geologi Kelautan,
Badan Litbang ESDM, Kementerian ESDM,
Jl. Dr. Junjuran 236 - Bandung 40174

Ditetapkan sebagai Majalah Ilmiah

TERAKREDITASI

Akreditasi berlaku mulai Juli 2015 - Juli 2018

Cibinong, 15 Juli 2015
Lembaga Ilmu Pengetahuan Indonesia
Ketua Panitia Penilai Majalah Ilmiah-LIPI

Prof. Dr. Rochadi,
NIP 195007281978031001



MARINE GEOLOGICAL INSTITUTE

RESEARCH AND DEVELOPMENT AGENCIES FOR ENERGY AND MINERAL RESOURCES

MINISTRY OF ENERGY AND MINERAL RESOURCES

Jalan Dr. Junjunan No. 236, Bandung-40174, Indonesia

<http://www.mgi.esdm.go.id>, E-mail : bulletin@mgi.esdm.go.id; bulletin_mgi@yahoo.com

