

Insecticidal effects of eudesmanes from *Pluchea sagittalis* (Asteraceae) on *Spodoptera frugiperda* and *Ceratitis capitata*

A. Sosa¹, M. Costa¹, A. Salvatore⁴, A. Bardon^{1,3}, S. Borkosky¹, N. Vera^{2*}

¹Instituto de Química Orgánica, ²Instituto de Estudios Farmacológicos, ³INQUINOA-CONICET, Facultad de Bioquímica, Química y Farmacia, UNT, Phone: 54-381-4247752, Ayacucho 471 (4000), Tucumán, Argentina

⁴Facultad de Agronomía y Zootecnia, UNT, Phone: 54-0381-4364147, Florentino Ameghino s/n. B° Mercantil. Campo Experimental (4105), El Manantial, Tucumán, Argentina

*Correspondence: N. Vera. Facultad de Bioquímica, Química y Farmacia. Universidad Nacional de Tucumán. Ayacucho 471. Tucumán (4000).

Abstract— *Eudesmanoids* play an important role in the protection of plants against herbivores. *Pluchea sagittalis* (Lamarck) Cabrera (Asteraceae) is widespread in tropical South America and contains compounds that provide protection against phytophagous insects. In the present work we isolated seven sesquiterpenoids with eudesmane skeletons that were evaluated for their insecticidal activities against *Spodoptera frugiperda* and *Ceratitis capitata*, pests that cause serious damage to crops in the Argentine northwest. The Eudesmanes were incorporated at different concentrations to the diet of *Spodoptera frugiperda*. In the choice test, larval feeding behavior was altered. The eudesmanes **1**, **5** and **7** showed the highest activity with feeding election indexes (FEI) of 50, 50, and 72 %, respectively at 200 µg/g of diet. When tested for insecticidal activity using neonate larvae with the nochoice artificial diet bioassays, eudesmane **1** was the most toxic in the larval stage (LD₅₀ 177.80 mg/g of diet). Compounds **5** lowered the percentage of adult emergence and produced the most malformations (72%) compared with control. Drastic effects were observed in the oviposition deterrence activity against *C. capitata*. The maximum oviposition deterrence (87 %) was recorded with eudesmane **5** at dose 30 µg/cm² of artificial fruit. Finally, eudesmanes **6** and **7** showed significant larval and pupal mortality against the first generation larvae of viable eggs oviposited by females fed with the treated diet (100 µg / g artificial diet).

Keywords— sesquiterpenoids, antifeedant, insect growth regulation, oviposition deterrence.

I. INTRODUCTION

It is well accepted that plant natural products may constitute new sources of insect pest control. The species belonging to the genus *Pluchea* (Asteraceae) consist of approximately 90

herbaceous species that grow in several countries of South America, *i.e.* Paraguay, Argentina and Brazil (Bremer & Anderberg 1994). Previous chemical investigations of the genus have shown the presence of eudesmane-type sesquiterpenoids (Ahmad *et al.* 1990; 1991; Guilhon & Müller 1996; 1998 a,b; Mahmoud 1997) as well as monoterpenes, lignan glycosides, triterpenoids, (Chakravarty & Mukhopadhyay 1994) and flavonoids (Ahmed *et al.* 1987; Scholz *et al.* 1994). *Pluchea sagittalis* (Lam.) Cabrera is popularly known as “Lucera,” “Yerba Lucero” or “Quitoc. Pharmacological studies demonstrated that aqueous and dichloromethane extracts obtained from *P. sagittalis* have a wide spectrum of anti-inflammatory, antioxidant (Pérez-García *et al.* 1996; 2001), antinociceptive and gastroprotective activities (Figueredo *et al.* 2011). Nevertheless, there is little information about their action on insects. Insecticidal and antifeeding effects have been reported for other sesquiterpenes against different insect species (Wang *et al.* 1991; Gonzalez *et al.* 1993; Dadang *et al.* 1996). The results of previous works by Céspedes *et al.* (2001) and Alarcon *et al.* (2013) reported feeding deterrent and insecticidal activities of certain eudesmane-type sesquiterpenoids against *S. frugiperda* (Lepidoptera: Noctuidae) larvae and *Drosophila melanogaster*, respectively. In a previous paper, we reported the isolation and identification of twelve eudesmane-type sesquiterpenoids from a Bolivian collection of *P. sagittalis*, together with their antifeedant activity at 100 µg / g of diet against *S. frugiperda*, under chosen conditions (Vera *et al.* 2008).

Objectives of the present study were to evaluate the insecticidal activity and sublethal effects of eudesmanes isolated from *P. sagittalis* against fall armyworm *S. frugiperda* and *Ceratitis capitata* (Wiedemann) (Diptera

Tephritidae), known as mediterranean fruit fly. For both insect pests, the bioassays were conducted under laboratory conditions.

II. MATERIALS AND METHODS

Plants and compounds.

P. sagittalis (Lamarck) Cabrera was collected at the flowering stage in December of 2013 in Tarija, Bolivia. Structures of tested compounds are shown in Figure 1. Compounds **1-7** were isolated from aerial parts of the plant and were purified in sufficient amount to be used in the bioassays. Identification of the eudesmanes **1-7** was accomplished by spectroscopic methods (IR, ¹H-NMR, ¹³C-NMR, and MS) and direct comparison with authentic samples (Vera *et al.* 2008). Previous to the bioassays, purity of eudesmanes was checked by HPLC (Gilson with a differential refractometer, Middleton, USA) in Beckman C18 and C8 columns (250 x 10 mm i.d., 5 µm particle size), using MeOH : H₂O mixtures as eluents.

Bioassays

Insect rearing

S. frugiperda larvae were obtained from our laboratory population. The larval diet consisted of a mixture of yeast (3 g), boiled and milled bean (250 g), wheat germ (12.5 g), agar agar (12.5 g), ascorbic acid (1.5 g), methyl p-hydroxybenzoate (1.5 g), formaldehyde (4 ml of a 38% water solution), and water (500 ml).

The colony of *C. capitata* used in the bioassays was derived from the laboratory of the Estación Experimental Agroindustrial Obispo Colombres (EEAOC). It was initiated with pupae of infested fruits obtained from northwestern Argentina cultivating oranges. Adults were fed on a diet prepared with an aqueous solution of a mixture of sucrose and hydrolyzed protein (3: 1 ratio). The brood chamber was maintained at 24 ± 2 °C, 60 ± 10% relative humidity and a photoperiod of 12L: 12D.

Antifeedant test against S. frugiperda (Choice test)

Antifeedant tests against *S. frugiperda* larvae were carried out as follows (Vera *et al.* 2008). A portion of artificial diet was mixed with acetone and, after solvent removal in vacuum this portion was employed as control diet. Another portion was mixed with an acetone solution of each test compound, in order to reach 25, 50, 100 and 200 µg/g of treatment per g of diet. After solvent evaporation, control and treated diets were placed in test tubes (20 replicates) in which third instars larvae were placed between both portions of diet to be kept at 27°C and 60 ± 15% relative humidity. When 50% of control diet had been eaten, control and treated diets were removed from the tubes and weighted

accurately. To evaluate the feeding behavior a “Feeding election index” was calculated as $FEI = (1 - T/C) 100$, where C and T represent the amounts of control and treated diets eaten, respectively.

Antifeedant test against S. frugiperda (No choice Test)

A portion of artificial diet was mixed with acetone and, after solvent removal in vacuum; this portion was employed as control diet. Another portion was mixed with an acetone solution of each test compound (treatment), in order to reach 25, 50, 100 and 200 µg/g of treatment per g of diet. After solvent evaporation, control diet was placed in a test tube and a larva was introduced. The same amount of treated diet was placed in a different test tube with a larva inside. Larvae were allowed to eat an after the 50% of control diet had been eaten, control and treated diets were removed from the tubes and weighted accurately. The experiment was carried out in 20 replicates. To evaluate the feeding behavior under no choice conditions, the FEI was calculated (Vera *et al.* 2006).

Insecticidal bioassay against S. frugiperda

The insecticidal bioassay activity against larvae of *S. frugiperda* was carried out as follows (Vera *et al.* 2006). Control and treated diets were placed in different test tubes (20 replicates for treated and 20 for control experiments) in which 2nd instar larvae were placed to be kept at 27 °C and 60 ± 15% relative humidity until emergency of the 1st generation of adults. Larval developmental periods as well as mortality rates were recorded for treatments containing eudesmanes (25, 50, 100 and 200 µg/g of diet) and control experiments. The larval period duration (days), larval and pupal mortality percentage and the number of malformed adults were registered.

Food consumption and utilisation

Ten days after the beginning of the experiment, the larval weight and diet eaten were determined again, in order to record the relative consumption rate (RCR), relative growth rate (RGR), efficiency of conversion of ingested food (ECI), efficiency of conversion of digested food (ECD) and approximate digestibility (AD) during the early larval instars (Waldbauer 1968; Farrar *et al.* 1989). The developmental indices were calculated as follows:

$$RCR = I/BaT$$

$$RGR = DB/BaT$$

$$ECI = (DB/I)*100$$

$$AD = [(I F)/I]*100$$

$$ECD = [DB/(I F)] * 100$$

where

I = weight of food consumed;

Ba = arithmetic mean of insect weight during the experiment $\frac{1}{4}$

$[(PF-PI)/\log (PF/PI)]$;

PF = larvae final weight (mg);

PI = larvae starting weight (mg);

T = feeding period in days;

DB = change in body weight;

F = weight of faeces produced during the feeding period

Oviposition-Deterrent test

Artificial fruits (oviposition substrates) were prepared by boiling a mixture of peach juice (500 ml), agar (15 g), and sodium benzoate (one teaspoonful) as preservative. This agar solution was poured into cylindrical molds, allowed to gel, and sliced. The agar cylinders were then wrapped in PVC film to avoid dehydration. The surface of the wrapped cylinder was pricked with a needle and treated with an acetone or methanol solution of the sample to be tested. An amount of 30 $\mu\text{g}/\text{cm}^2$ of the test compound was deposited. Control cylinders were impregnated only with the solvent that was then removed in vacuum. Three groups of *C. capitata* adults were selected from the EEAOC laboratory colony. Each group, consisting of seven male–female pairs, was placed in a small cage and covered with voile (a light, almost transparent cloth made of silk). Two agar cylinders (sample and control) were placed over the voile, and females oviposited on one or the other according to their preference (Fig. 2). After 4 days, eggs were gently rinsed from the agar and counted.

The oviposition index was defined as $\text{IO}\% = (1-T/C) \times 100$, where T is the number of eggs laid in the treated artificial fruit, and C is the number of eggs deposited in the control fruit (Socolsky *et al.* 2008).

Insecticidal bioassay with *C. capitata* adult flies

Assays were performed by adding compounds to artificial diet of the adult flies at a concentration of 100 $\mu\text{g} / \text{g}$ (sample weight / weight of artificial diet). This diet is offered to adult flies (F1) of *C. capitata* (300 flies per trial). They are kept in cages until seen 2 days after the beginning of intercourse and then are allowed to oviposit.

In this assay were evaluated:

- 1- The volume of oviposited eggs.
- 2- The viability of eggs.
- 3- The mortality of adult flies.
- 4- The larval mortality F2 (1st generation of neonates from parents subjected to treatment).
- 5- The pupal mortality F2.

The experiment was performed in triplicate, using an identical but without the addition of substances, as a target system. The experience takes place at a controlled

www.ijeab.com

temperature (24 ± 2 °C) and r.h. (60 ± 10) and photoperiod L12:D12 (Salvatore *et al.* 2004).

Statistical analysis

The results are reported as mean \pm SEM. The differences in the mean values were evaluated by analysis of variance (ANOVA). The Tukey test was used for all pair wise multiple comparisons of groups. In all statistical analysis, $P > 0.05$ was considered not significant. The LD_{50} values for each activity were calculated using a probit analysis software program based on percentage of mortality obtained at each concentration of the samples (MINITAB Release 14 for Windows).

III. RESULTS

During our screening program on biological activities of plants from Argentine and Bolivia in a preliminary trial (Vera *et al.* 2008), the eudesmanes of *P. sagittalis* displayed antifeedant activity at 100 $\mu\text{g}/\text{g}$ of diet on *S. frugiperda*. Based on this information, we carried out several studies with eudesmanes **1 - 7** (Figure 1), in a concentration range 25-200 $\mu\text{g}/\text{g}$ of diet.

Antifeedant activity

The antifeedant activities of eudesmanes **1-7** against *S. frugiperda* are shown in Table 1. *S. frugiperda* larvae ate significantly less when fed a diet treated with compounds **1**, **5** and **7** at 200 $\mu\text{g}/\text{g}$ of diet (Feeding election indexes FEI, of 50, 50, and 72%, respectively). Eudesmane **5**, was the only one with an inhibition percentage higher than 25% at the lowest dose tested 25 $\mu\text{g}/\text{g}$ diet. All eudesmanes have a concentration dependent deterrent activity. In the no-choice feeding assay none of the tested compounds displayed antifeedant effects against *S. frugiperda* (Table 1).

Insecticidal bioassay against *S. frugiperda*

A significant growth reduction was observed in the larvae fed on the treated diet containing eudesmanes **1-3**, **5** and **7**, even at the lowest concentration. Growth inhibition of about 60% relative to control, was observed in the treatment with compound **7** at all the concentrations tested (Table 3). A marked decrease in the daily consumption of diet was observed in treatments with eudesmanes **1**, **3**, **5** and **7** at the four concentrations tested. At doses of 100 μg and 200 $\mu\text{g}/\text{g}$, diet intake inhibition percentage reached 85% relative to control. A significant reduction ($P < 0.01$) in ECI and ECD was also observed in the compounds at 100 and 200 $\mu\text{g}/\text{g}$ except for eudesmanes **4** and **6** that showed no significant differences compared to control. No significant differences ($P < 0.01$) on DA were observed in any of the eudesmanes at 100 $\mu\text{g}/\text{g}$ of diet (Table 2).

The larvae were fed during the course of the experiment. Eudesmanes 1-7 were incorporated to the larval diet at 25-200 $\mu\text{g/g}$ and larval development and mortality were recorded. Our results are outlined in Table 2. Eudesmane 1 was the most toxic in the larval stage with 57 % larval mortality at 200 $\mu\text{g/g}$ of diet. Products 1 and 5 also showed high percentages of pupal mortality with 33 and 30%, respectively, at the highest dose tested (200 mg/g diet). Adult insects that survived the treatment showed higher rates of malformation. Compound 5 was the most active at 200 $\mu\text{g/g}$ diet with a 50% of malformation in the wings which prevented them from reproducing. Also, the duration of larval and pupal cycle, compared to that of the control, were affected by the treatments at doses of 100 and 200 $\mu\text{g/g}$ of diet (Table 3).

Oviposition Deterrent test

The results are summarized in Table 4. All tested compounds (1-7) produced oviposition deterrence. At a dose 30 $\mu\text{g/cm}^2$ of artificial fruit the eudesmanes 1 and 5 were the most active with percentages of 84.18 and 87.52 of inhibition, respectively. (Table 4)

4.4 Insecticidal bioassay with *C. capitata* adult flies

Only the major eudesmanes (6 and 7) were evaluated. As shown in Table 5, the treatments did not produce mortality in adult insects that ingested the treated diet. The viability of oviposited eggs by females who ingested treatments was not affected. However, higher percentages of F2 neonate larvae mortality were observed (F2: the first generation larvae of viable eggs oviposited by females consuming the treated diet), 51% for eudesmane 6 and 54% for eudesmane 7 at 100 $\mu\text{g/g}$ artificial diet. F2 pupal mortality was 33 and 29% for 6 and 7 respectively (Table 5).

IV. DISCUSSION

Botanicals are a rich source of organic chemicals on earth. Discovery of novel toxins and/or antifeedants from plant extracts has been recently emphasized as a potential method for the development of ecologically safe pesticides (Wheeler *et al.* 2001). Antifeedants offer first line of crop protection against notorious insects. Any substance that reduces food consumption by an insect can be considered as an antifeedant or feeding deterrent (Isman 2002). In general, antifeedants have profound adverse effects on insect feeding behavior (Hummelbrunner LA & Isman MB). Antifeedants can be described as allomone substances which inhibit feeding and do not kill the insect pests directly, but rather limit its developmental potential considerably and act as a phagodeterrent or phagorepellent over test as well as permanent insect pests feeding on the plant (Lakshmanan *et al.* 2012). Plant substances acting as antifeedants are found

in all the compound groups of secondary plant metabolism. However, the most effective insect feeding inhibitors come from terpenoids, alkaloids, saponins and polyphenols (Koul 2005). In a previous study carried out by our group, we determined that sesquiterpenes eudesmane-type produced a significant deterrence of the intake in *Spodoptera frugiperda* larvae when the diet choice tests were performed at 100 $\mu\text{g} / \text{g}$ diet (Vera *et al.* 2008). The results of our bioassays showed that eudesmanes 1-7 produced inhibition of feeding in a concentration range between 25-200 $\mu\text{g} / \text{g}$ of diet and the effect is concentration dependent for all eudesmanes tested in the choice test. Compound 7 was the most active, reaching a value of inhibition at 200 $\mu\text{g} / \text{g}$ of diet seven times greater than the one at 25 $\mu\text{g} / \text{g}$ of diet. However, when evaluating the feeding behavior of the insect without giving it the choice, we observed that none of the eudesmanes inhibited the insect from feeding, even at the highest dose used (200 $\mu\text{g} / \text{g}$ diet). This difference in dietary behavior may be due to a suppressive effect of food where the reduction of food intake occurs after initial consumption. Similar results were presented by Macleod *et al.* (1990), who used two active ingredients isolated from *M. azedarach* (meliatoxin A2 and meliatoxin B1), which inhibited feeding when the larvae of *Spodoptera litura* were able to choose. However, in tests without this option Meliatoxin B1 did not reduce the intake. Other extracts also showed increased anti-food activity on *S. littoralis*, when the caterpillars had access to the control over non-choice trials (Sadek 2003).

The nutritional indices help us to be able to approach the mode of action of these compounds. The EUs 1, 3, 5 and 7 produced a significant reduction in larval growth (RGR) even at 25 $\mu\text{g} / \text{g}$ diet. In relation to consumption, there is a marked decrease in RCR for treatments with EUs 1, 3, 5 and 7 at all doses evaluated. A significant reduction in ECI and ECD were observed for all compounds at 100 and 200 $\mu\text{g} / \text{g}$, except for EUs 4 and 6 that did not show significant differences with the control (table 2). Energy diversion to other metabolic pathways, such as those involved in the detoxification of allelochemicals, may be the cause of the decrease in efficiencies (Koul & Isman 1991; Hernández & Vendramim 1997). This decrease causes inhibition of larval growth and is considered as a chronic post-ingestive toxic effect (Wheeler & Isman 2001; Sadek 2003) by several authors. As the diet concentration of the EUs increases, feed intake, growth and feed conversion efficiency reduction were observed suggesting that these compounds have a toxic effect on *S. frugiperda*, producing larval, pupal mortality and high percentages of malformations in adults.

The EUs 18 and 21 produced the greatest *Ceratitis capitata* inhibition of oviposition with% IO of 84 and 88%, respectively. That is to say, these substances caused a reduction of more than 80% in the number of eggs deposited in the treated artificial fruit in comparison with the artificial fruit control. These compounds exhibit an activity similar to naringin, isolated from *Elaphoglossum spathulatum* (IO = 74%, 6 µg / cm²), one of the natural products evaluated in our work group that produced the greatest inhibition of oviposition in *C. capitata* (Socolsky 2003), which would be promising for the control of this pest. These results constitute the first antecedent of the study of EUs as inhibitors of oviposition in *C. capitata*.

Although the number of tested eudesmanes was reduced to establish a relationship between structure and biological activity, our results show how small structural changes may modify the mode of action of these compounds. Eudesmanes that had a hydroperoxide (**5**) or formoxi groups in their structures (**1** and **7**) were the most active, while the eudesmanes possessing an acetyl group in position 4 lost such activity noticeably. Additionally, an α,β -unsaturated system played an important role in antifeedant activity, as seen when comparing compound **2** with compound **3**, that showed a low activity (Table 1). These results are in agreement with those published in previous works by other authors and our research group (Srivastava *et al.* 1990; Faini *et al.* 1997; Vera *et al.* 2008; Alarcon *et al.* 2013). Eudesmanes highly oxygenated has a marked influence on biological activity and the individual biological activities of sesquiterpenoids depend on the different functional groups present and the pattern of oxygenation. Similar results were published by Gonzalez *et al.* (1993); Céspedes *et al.* (2001); Guan *et al.* (2005); and Alarcon *et al.* (2013). All eudesmanes assayed have trans-fused decalin architecture differently from the sesquiterpenes used by Miyazawa *et al.* (2000) and Alarcon *et al.* (2013). The sites and mode of action of these compounds are being investigated and probably correspond to a combination of antifeedant action, as well as insecticidal and insect growth regulation activities. Although some natural insecticides are found on the market, the search for new compounds with activity against a variety of insects are always necessary either to prevent the emergence of resistance in insects or to guarantee the ready availability of natural insecticides through more widely distributed sources. This study suggests that eudesmanes and plants containing it might be used as new tools for protecting from harmful insects, especially in organic agriculture. Additionally, these compounds could be promising precursors to generate a

series of more active derivatives that acted as a chemical defense against predation by certain phytophagous insects.

Conflict of interest

The authors declare that there are no conflicts of interest and they have no actual or potential competing financial interests

ACKNOWLEDGMENTS

Work in Argentina was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (CONICET) and Consejo de Investigaciones de la Universidad Nacional de Tucumán (CIUNT).

REFERENCES

- [1] ALARCON, J., LAMILLA, C. & CESPEDES, C.L. (2013). Insecticidal activity of sesquiterpenes skeleton synthesized by the conventional Robinson annulations reaction on *Drosophila melanogaster*. *Industrial Crops and Products* **42**, 268-272.
- [2] AHMAD, V.U., SULTANA, A. & FIZZA, K.Z. (1990). Two new terpenoids from *Pluchea arguta*. *Zeitschrift für Naturforschung* **45B**, 385-388.
- [3] AHMAD, V.U., FIZZA, K.Z., KHAN, M.A. & FAROOQUI, T.A. (1991). Sesquiterpenes from *Pluchea arguta*. *Phytochemistry* **30**, 689-691.
- [4] AHMED AA, MELEK FR & MABRY TJ (1987). Sulfated and non-sulfated flavonoids from *Pluchea dioscoridis*. *Journal of Natural products* **50**, 311.
- [5] BREMER, K. & ANDERBERG, A. (1994). Asteraceae: cladistics & classification. Portland, Or.: Timber Press.
- [6] CÉSPEDES, C.L., ALARCÓN, J., ARANDA, E., BECERRA, J. & SILVA, M. (2001). Insect growth regulator and insecticidal activity of β -dihydroagarofurans from *Maytenus* spp. (Celastraceae). *Zeitschrift für Naturforschung C* **56c**, 603-613.
- [7] CHAKRAVARTY, A.K. & MUKHOPADHYAY, S. (1994). New thiophene derivatives from *Pluchea indica*. *Indian Journal of Chemistry* **33B**, 978-980.
- [8] DADANG OHSAWA, K., KATO, S. & YAMAMOTO, I. (1996). Insecticidal compound in tuber of *Cyperus rotundus* L. against the diamondback moth larvae. *Journal of Pest Science* **21**, 444-446.
- [9] FAINI, F., LABBE, C., TORRES, R., DELLE MONACHE, G., DELLE MONACHE, F. & COLL, J. (1997). Eudesmane derivatives from *Flourensia*

- thurifera: structure and biological activity. *Natural Products Letters* **11**, 1-4.
- [10] FARRAR, R.R., BARBOUR, J.D. & KENEDY, G.G. (1989). Quantifying food consumption and growth in insects. *Annales of the Entomological Society of America* **82**, 593-598.
- [11] FIGUEREDO, S.M., PINTO DO NASCIMENTO, F., FREITAS, C.S., HATSUKO BAGGIO, C., SOLDI, C., PIZZOLATTI, M.G., CHELLION DE IBARROLA, M., DEGEN DE ARRUA, R.L. & SOARES SANTOS, A.R. (2011). Antinociceptive and gastroprotective actions of ethanolic extract from *Pluchea sagittalis* (Lam) Cabrera. *Journal of Ethnopharmacology* **135**, 603-609.
- [12] GONZALEZ, A., JIMENEZ, I.A., RAVELO, A.G. & BAZZOCHI, I.L. (1993). Minor sesquiterpenes from *Maytenus canariensis* with insecticidal and antifeedant activity. *Tetrahedron* **49**, 6637-6644.
- [13] GUAN, Y.K., LI, P., FANG, L.F. & LI, Y.L. (2005). The first total synthesis of 3-oxo-11,12,13-trihydroxyeudesm-4-ene. *Chinese Chemical Letters* **16**, 711-712.
- [14] GUILHON, G.M.S.P. & MÜLLER, A.H. (1996). Eudesmane derivatives from *Pluchea quitoc*. *Phytochemistry* **43**, 417-421.
- [15] GUILHON, G.M.S.P. & MÜLLER, A.H. (1998a). Eudesmane sesquiterpenoids from *Pluchea quitoc*. *Phytochemistry* **47**, 227-229.
- [16] GUILHON, G.M.S.P. & MÜLLER, A.H. (1998b). Eudesmanolides and epoxycauthemones from *Pluchea quitoc*. *Phytochemistry* **49**, 1347-1351.
- [17] HERNANDEZ, C. R. & VENDRAMIN, J.D. (1997). Bioactivity evaluation of aqueous extracts of Meliaceae to *Spodoptera frugiperda*. *Revista agrícola* **72**, 305-318.
- [18] HUMMELBRUNNER, L.A. & ISMAN, M.B. (2001). Acute, sublethal, antifeedant and synergistic effects of monoterpenoid essential oil compounds on the tobacco cutworm *Spodoptera litura* (Lepidoptera: Noctuidae). *Journal of Agricultural and Food Chemistry* **49**, 715-720.
- [19] ISMAN, M.B. (2002). Insect antifeedants. *Pesticide Outlook* **13**, 152-157.
- [20] KOUL, O. (2005). Insect Antifeedants. CRC Press LLC: Boca Raton, FL, USA.
- [21] KOUL, O. & ISMAN, M. B. (1991). Effects of azadirachtin on the dietary utilization and development of the variegated cutworm *Peridroma saucia*. *Journal of Insect Physiology* **37**, 591-598.
- [22] LAKSHMANAN, S., KRISHNAPPA, K. & ELUMALAI, K. (2012). Certain plant essential oils against antifeedant activity of *Spodoptera litura* (Fab.), *Helicoverpa armigera* (Hub.) and *Achaea janata* (Linn.) (Lepidoptera: Noctuidae). *International Journal of Current Life Sciences* **2**, 5-11.
- [23] MACLEOD, J. K., MOELLER, P. D., MOLINSKI, T. F. & KOUL, O. (1990). Antifeedant activity against *b* larvae and [13C] NMR spectral assignments of the meliatoxins. *Journal of Chemical Ecology* **16**, 2511-2518.
- [24] MAHMOUD, A. (1997). 7-*epi*-eudesmanes, eudesmanoic acids, eudesmanolides and other sesquiterpenes from *Pluchea dioscoridis*. *Phytochemistry* **45**, 1633-1688.
- [25] MIYAZAWA, M., NAKAMURA, Y. & ISHIKAWA, Y. (2000). Insecticidal sesquiterpene from *Alpinia oxyphylla* against *Drosophila melanogaster*. *Journal of Agricultural and Food Chemistry* **48**, 3639-3641.
- [26] PEREZ-GARCÍA, F., MARÍN, E., CAÑIGUERAL, S. & ADZET, T. (1996) Anti-inflammatory action of *Pluchea sagittalis*: involvement of an antioxidant mechanism. *Life Sciences* **59**, 2033-2040.
- [27] PEREZ-GARCÍA, F., MARÍN, E., PARELLA, T., ADZET, T. & CAÑIGUERAL, S. (2005). Activity of taraxasteryl acetate on inflammation and heat shock protein synthesis. *Phytomedicine* **12**, 278-284.
- [28] SADEK, M.M. (2003). Antifeedant and toxic activity of *Adhatoda vasica* leaf extract against *Spodoptera littoralis* (Lep. Noctuidae). *Journal of Applied Entomology* **127**, 396-404.
- [29] SALVATORE, A., BORKOSKY, S., WILLINK, E. & BARDÓN, A. (2004). Toxic effects of lemon peel constituents on *Ceratitis capitata*. *Journal of Chemical Ecology* **30**, 323-333.
- [30] SCHOLZ, E., HEINRICH, M. & HUNKLER, D. (1994). Caffeoylquinic acids and some biological activities of *Pluchea symphytifolia*. *Planta Medica* **60**, 360-364.
- [31] SOCOLSKY, C., SALVATORE, A., ASAKAWA, Y. & BARDÓN, A. (2003). Bioactive new bitter-tasting p-hydroxystyrene glycoside and other constituents from the fern *Elaphoglossum spathulatum*. *Arkivoc* **10**, 347-355.
- [32] SOCOLSKY, C., FASCIO, M.L., D'ACCORSO, N.B., SALVATORE, A., WILLINK, E., ASAKAWA, Y. & BARDON, A. (2008). Effects of p-Vinylphenyl Glycosides and other related compounds on the oviposition behavior of *Ceratitis capitata*. *Journal of Chemical Ecology* **34**, 539-548.
- [33] SRIVASTAVA, R.P., PROKSCH, P. & WRAY, V. (1990). Toxicity and antifeedant activity of a

- sesquiterpene lactone from *Encelia* against *Spodoptera littoralis*. *Phytochemistry* **29**, 3445-3448.
- [34] VERA, N., POPICH, S., LUNA, L., CRAVERO, R., GONZÁLEZ SIERRA, M. & BARDÓN, A. (2006). Toxicity and synergism in the feeding deterrence of some coumarins on *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae). *Chemistry & Biodiversity* **3**, 21-26.
- [35] VERA, N., MISICO, R., GONZALEZ SIERRA, M., ASAKAWA, Y. & BARDON, A. (2008). Eudesmanes from *Pluchea sagittalis*. Their antifeedant activity on *Spodoptera frugiperda*. *Phytochemistry* **69**, 1689-1694.
- [36] WANG, M., QIN, H., KONG, M. & LI, Y. (1991). Insecticidal sesquiterpene polyol ester from *Celastrus angulatus*. *Phytochemistry* **30**, 3931-3933.

- [37] WALDBAUER, G.P. (1968). The consumption and utilization of food by insects. *Advances Insect Physiology* **5**, 229-288.
- [38] WHEELER, D.A. & ISMAN, M.B. (2001). Antifeedant and toxic activity of *Trichilia americana* extract against the larvae of *Spodoptera litura*. *Entomologia Experimentalis et Applicata* **98**, 9-16.
- [39] WHEELER, D.A., ISMAN, M.B., SANCHEZ-VINDAS, P.E., ARNASON, J.T. (2001). Screening of Costa Rican *Trichilia* species for biological activity against the larvae of *Spodoptera litura* (Lepidoptera: Noctuidae). *Biochemical Systematic and Ecology* **29**, 347-358.




Fig.1: Eudesmanes isolated from *Pluchea sagittalis*

Table.1: Insect antifeedant activities of eudesmanes (1-7) on *S. frugiperda*.

Feeding deterrence index								
Doses ($\mu\text{g/g}$) Compounds	a) % FEI _{CH} Choice test				a) % FEI _{NCH} No Choice test			
	25	50	100	200	25	50	100	200
1	10.9 ± 1.0 a	21.0 ± 3.0 b	30.7 ± 5.2 d	50.4 ± 5.0 c	1.0 ± 0.2 a	1.0 ± 0.4 a	4.0 ± 0.4 c	5.4 ± 0.4 d
2	10.1 ± 1.7 a	12.7 ± 0.8 a	13.7 ± 2.0 a	17.9 ± 2.8 b	1.1 ± 0.2 a	2.0 ± 0.8 b	2.0 ± 0.3 b	4.9 ± 0.8 d
3	10.8 ± 2.9 a	13.1 ± 2.1 a	29.8 ± 3.8 b	34.6 ± 4.3 d	0.8 ± 0.1 a	3.1 ± 0.2 bc	3.0 ± 0.4 b	4.6 ± 0.3 d
4	12.4 ± 2.5 a	19.1 ± 2.6 b	28.0 ± 3.6 b	37.3 ± 3.4 d	2.4 ± 0.5 b	3.1 ± 0.3 bc	7.0 ± 0.7 e	7.3 ± 0.8 de
5	27.2 ± 4.2 b	51.8 ± 3.6 c	50.1 ± 4.1 c	50.7 ± 5.3 c	2.2 ± 0.2 b	1.8 ± 0.6 ab	7.0 ± 1.0 e	6.7 ± 1.6 de
6	10.9 ± 2.0 a	11.0 ± 3.0 a	19.2 ± 2.2 b	19.4 ± 5.0 b	2.9 ± 0.8 b	3.5 ± 0.5 c	6.0 ± 1.0 de	6.4 ± 1.6 de
7	10.5 ± 2.0 a	31.8 ± 4.2 d	32.3 ± 4.2 d	72.2 ± 6.7 e	2.5 ± 0.2 b	3.8 ± 0.6 c	5.0 ± 0.9 d	6.2 ± 1.5 de

The values followed by the same letter are not significantly different. The significance level $P < 0.05$. * FEI (%) = Feeding election index = $[(1 - T/C) \times 100]$.

Table.2: Effect of eudesmanes (1-7) incorporated into larval diet on food consumption and utilization by *S. frugiperda* larvae.

Nutritional indices						
Compounds	Doses ($\mu\text{g/g}$)	a) RCR = I/(B _a T)	a) RGR = $\Delta\text{B}/(\text{B}_a\text{T})$	a) ECI = $(\Delta\text{B}/\text{I} * 100)$	a) AD = $(\text{I} - \text{F}/\text{I}) * 100$	a) ECD = $(\Delta\text{B}/\text{I} - \text{F}) * 100$
Control		0.51 ± 0.06 a	0.25 ± 0.03 a	21.12 ± 2.9 a	73.88 ± 3.5 a	23.45 ± 1.5 a
1	25	0.39 ± 0.06 b	0.12 ± 0.03 b	23.13 ± 1.8 a	73.20 ± 8.0 a	22.63 ± 2.2 a
	50	0.25 ± 0.05 c	0.11 ± 0.03 b	18.43 ± 2.2 b	68.21 ± 2.6 a	18.44 ± 2.8 b
	100	0.14 ± 0.03 d	0.09 ± 0.03 b	18.34 ± 3.8 b	69.26 ± 1.5 a	17.62 ± 2.5 b
	200	0.13 ± 0.02 d	0.06 ± 0.06 c	14.38 ± 2.6 c	67.51 ± 2.4 a	14.13 ± 1.8 c
2	25	0.45 ± 0.03 a	0.09 ± 0.02 b	20.09 ± 2.8 a	74.22 ± 2.3 a	20.09 ± 2.8 a
	50	0.44 ± 0.03 a	0.08 ± 0.008 d	22.32 ± 2.5 a	75.64 ± 3.3 a	22.32 ± 2.5 a
	100	0.43 ± 0.02 a	0.07 ± 0.01 d	17.78 ± 2.8 b	71.16 ± 3.1 a	17.78 ± 2.8 b
	200	0.34 ± 0.04 b	0.08 ± 0.02 d	17.48 ± 1.5 b	76.01 ± 2.2 a	17.48 ± 1.5 b
3	25	0.29 ± 0.06 b	0.11 ± 0.03 b	14.78 ± 2.6 c	71.13 ± 6.3 a	13.07 ± 7.2 c
	50	0.29 ± 0.07 b	0.11 ± 0.02 b	13.88 ± 1.6 c	78.67 ± 8.3 a	15.52 ± 1.2 c
	100	0.15 ± 0.02 d	0.04 ± 0.008 f	12.96 ± 2.3 c	72.00 ± 3.1 a	10.12 ± 3.1 c
	200	0.15 ± 0.04 d	0.04 ± 0.02 c	12.45 ± 1.8 c	69.91 ± 4.2 a	10.45 ± 4.9 c
4	25	0.47 ± 0.05 a	0.24 ± 0.04 a	20.58 ± 1.2 a	69.21 ± 3.0 a	27.14 ± 7.2 a
	50	0.44 ± 0.06 a	0.23 ± 0.04 a	22.02 ± 1.8 a	73.78 ± 4.2 a	26.96 ± 3.2 a
	100	0.42 ± 0.06 a	0.21 ± 0.05 a	21.71 ± 0.7 a	74.16 ± 0.8 a	24.79 ± 4.8 a
	200	0.42 ± 0.06 a	0.26 ± 0.05 a	21.51 ± 1.3 a	73.84 ± 4.0 a	23.29 ± 5.9 a
5	25	0.36 ± 0.03 b	0.06 ± 0.02 c	21.32 ± 0.5 a	75.49 ± 5.5 a	20.12 ± 3.3 a
	50	0.34 ± 0.04 b	0.05 ± 0.02 c	18.45 ± 0.5 b	72.79 ± 3.1 a	18.53 ± 5.4 b
	100	0.13 ± 0.01 d	0.02 ± 0.007 f	17.67 ± 0.4 b	69.21 ± 2.5 a	18.28 ± 8.2 b
	200	0.13 ± 0.03 d	0.05 ± 0.01 dc	14.65 ± 0.6 c	71.20 ± 2.4 a	14.73 ± 7.1 c
6	25	0.48 ± 0.05 a	0.29 ± 0.04 a	20.58 ± 1.2 a	71.58 ± 8.0 a	20.27 ± 4.2 a
	50	0.42 ± 0.04 ab	0.24 ± 0.01 a	22.02 ± 0.8 a	74.16 ± 5.2 a	22.61 ± 4.2 a
	100	0.43 ± 0.05 ab	0.24 ± 0.02 a	20.71 ± 0.7 a	78.83 ± 7.2 a	23.52 ± 5.3 a
	200	0.41 ± 2.1 b	0.25 ± 0.02 a	21.31 ± 1.3 a	76.64 ± 2.0 a	19.80 ± 2.9 a

7	25	0.38 ± 0.05 b	0.14 ± 0.02 b	16.17 ± 1.4 b	77.01 ± 4.9 a	18.12 ± 4.3 ab
	50	0.39 ± 0.04 b	0.15 ± 0.01 b	15.70 ± 0.3 b	75.40 ± 6.1 a	18.65 ± 7.1 ab
	100	0.40 ± 0.03 b	0.15 ± 0.02 b	12.64 ± 0.8 c	75.55 ± 2.5 a	13.67 ± 5.4 b
	200	0.40 ± 0.04 b	0.14 ± 0.03 bc	11.93 ± 1.8 c	78.20 ± 4.4 a	12.78 ± 4.4 b

Values are mean ± SE (n = 20). The values followed by the same letter are not significantly different. The significance level P < 0.05.

Table.4: Effect of eudesmanes (1-7) on the oviposition-behavior of *C. capitata*

Compounds	Number of Eggs Laid on the Control Fruit	Number of Eggs Laid on the Treated Fruit	% IO = (1-T/C) *100 30 µg/cm ²
1*	810 ± 43a	256 ± 40 a	84.18 ± 13.1a
2*	741 ± 25a	254 ± 27b	59.96 ± 7.5b
3*	876 ± 12a	670 ± 14c	34.85 ± 1.1c
4	457 ± 31a	411 ± 34d	11.2 ± 2.0 d
5*	522 ± 51a	91 ± 4 ^a	87.52 ± 5.5a
6	868 ± 23a	753 ± 14d	13.2 ± 3.0 d
7	894 ± 11a	488 ± 13 ^a	45.4 ± 4.0 a

Numbers represent mean ± SEM, n=3. Means within a column followed by the same letter are not significantly different (P>0.05, paired t test)

Table.5: Insecticidal activity of eudesmanes (6 and 7) against adults of *Ceratitis capitata*.

	Mortality F1 (%)	Oviposition of F1 Volume of eggs (ml) ^a	Hatching eggs of F1 (%)	Larval Mortality of F2 (%)	Pupal Mortality of F2 (%)
Control	5 a	1.2 ± 0.1a	94 a	12a	11a
6	3 a	0.4 ± 0.05b	95 a	51b	33b
7	2 a	0.4 ± 0.1b	96 a	54b	39b

F1: Adult insects fed on the treated diet. F2: The first generation larvae of viable eggs oviposited by females consuming the treated diet.

Numbers represent mean ± SEM, n=3. Means within a column followed by the same letter are not significantly different (P>0.05, paired t test)