

# *In Vitro* Study on total Phenols, Flavonoids Content and DPPH Activity of *Withania* Species

Nishtha Raj<sup>1</sup>, Ayushi Agarwal<sup>2</sup>, Neelam Chaturvedi<sup>3</sup>

<sup>1,2</sup>Research Scholar, Department of Food Science and Nutrition, Banasthali University, Dist. Tonk, Rajasthan, India.

<sup>3</sup>Associate Professor, Department of Food Science and Nutrition, Banasthali University, Dist. Tonk, Rajasthan, India.

**Abstract**— The escalating interest in appraisal of antioxidant power of herbal plant as medicine, the current study was carried out to explore the antioxidant potential of aqueous extracts of *Withania somnifera* root and *Withania coagulans* fruit in-vitro. Antioxidant activity; total phenol, total flavonoids and DPPH free radical scavenging assay of *Withania somnifera* root and *Withania coagulans* fruit aqueous extracts were determined by using reference standards gallic acid, quercetin and ascorbic acid, respectively. The highest total phenols content (mgGAE/g) and total flavonoids content (mgQE/g) was found to be  $33.1 \pm 0.82$  and  $1.86 \pm 0.01$  respectively in aqueous *somnifera* root extracts as compared to *coagulans* fruit extract. The DPPH radical scavenging activity of the both extracts was increased with the increasing concentration and was observed high in aqueous extract *somnifera* root ( $IC_{50} = 54$ ) than *coagulans* fruit ( $69 \mu\text{g/ml}$ ) aqueous extract. Thus, *Withania somnifera* root has potent antioxidant activity and may serve as a good pharmacotherapeutic agent which could be explored to provide affordable medicines to masses.

**Keywords**— DPPH (2, 2-diphenyl-1-picrylhydrazyl), Total Flavonoids Content, Total Phenol Content, *Withania somnifera*, *Withania coagulans*.

## I. INTRODUCTION

The use of medicinal plants are increasing day by day in developing and developed countries due to their no side effects<sup>1</sup>. Traditional herbal medicines are new therapeutic contender because of their structural complexity, chemical diversity and wide variety of antimicrobial activity<sup>2</sup>. The genus *Withania* (Family: Solanaceae) is a highly-acclaimed remedial plants in the Indian Ayurvedic system of medicine because of its valuable pharmaceutical and nutraceutical properties. It is a small group of herbs distributed from the Northern Africa to the South-west of Asia. Among the twenty-three-known species of *Withania*, only two (*Withania somnifera* and *Withania coagulans*) are economically significant medicinal plant<sup>3</sup>.

*Withania somnifera* commonly known as “Ashwagandha” or “Indian Ginseng” and is an important plant in Indian traditional Ayurvedic system of medicines<sup>4</sup>. Ashwagandha improves energy and also memory by enhancing the brain and nervous function; shows anxiolytic effects, has hepatoprotective property, raises haemoglobin level and red blood cell count, improve energy level, improve the cell-mediated immunity; promotes vigour and vitality along with cheerful sexual life and reproductive equilibrium and act as powerful adaptogen<sup>5</sup>. *Withania coagulans* also known as ‘Indian cheese maker’ or ‘vegetable rennet’<sup>6</sup>. It is traditionally used as digestant, anti-flatulent, sedative, antihyperglycemic antimicrobial, anti-inflammatory, antitumor, hepatoprotective, cardiovascular, immune-suppressive, free radical scavenging and central nervous system depressant activities<sup>7</sup>.

There are several plants used in Ayurvedic medicinal systems origin with potential therapeutic activity, which are widely used as Ayurvedic medicine<sup>8</sup>. Therefore, an endeavour has been made to investigate the antioxidant properties of *Withania somnifera* root and *Withania coagulans* fruit.

## II. MATERIALS AND METHODS

### Sample Collection

*Withania somnifera* roots and *Withania coagulans* fruits were collected from local market of Delhi and authenticated by scientist of National Institute of Ayurveda, Jaipur, Rajasthan. Roots and fruits were washed and dried in open air for 2-3 weeks at 35-40° C and then dried material was pulverized in an electric grinder and stored in plastic containers in refrigerator (5° C), until further analysis.

### Aqueous Extract Preparation

20 g of powdered plant material was kept in 200ml conical flask and add 100 ml of distilled water. The mouth of the conical flask was covered with the aluminum foil and kept in a reciprocating shaker for 25 mins for continuous agitation at 150 rev/min for thorough mixing. Then extracts was filtered by using muslin cloth followed by Whatmann

filter paper No. 42 (125mm). The content was filtered by using rotatory vacuum evaporator with the water bath temperature of 65° C and finally the residue were collected and used for the analysis<sup>9</sup>.



*Withania coagulans*



*Withania somnifera*

#### Determination of Total Phenols Content:

Total phenols were determined by Folin-Ciocalteu Reagent. A dilute extract of root and fruit (0.5 ml of 1:10g/ml) or Gallic acid (standard phenol compound) was mixed with Folin-Ciocalteu reagent (5 ml, 1:10 diluted with distilled water) and 4 ml of aqueous sodium carbonate (1M). The mixtures were kept at dark ambient condition for 15 min and the total phenols were determined by spectrophotometer at 765 nm. The standard curve was prepared using 0, 50, 100, 150, 200, 250mg/l solutions of Gallic acid in methanol: water (50:50, v/v). Total phenol values are expressed in terms of Gallic acid equivalent (mg/g of dry mass), which is a common reference compound<sup>10</sup>.

#### Determination of total flavonoids content:

Aluminum chloride colorimetric method was used for flavonoids determination. Root and fruit extracts (0.5 ml of 1:10 g/ml) in aqueous were separately mixed with 1.5 ml of

methanol, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. It remained at room temperature for 30 min; the absorbance of the reaction mixture was measured at 415 nm with a UV-Visible spectrophotometer. The calibration curve was prepared by preparing quercetin solutions at concentrations 12.5 to 100 g/ml in aqueous<sup>11</sup>

#### DPPH radical scavenging activity:

The ability of the aqueous extracts to scavenge free radicals was determined against a very stable free radical DPPH (2, 2-diphenyl-1-picrylhydrazyl) determined Spectrometric method. Aliquots of the sample extract at different concentrations 20-200 µg/ml were added to 1 mm aqueous solutions of DPPH. Each mixture was vortexed vigorously and left for 30 min at room temperature in the dark. The absorbance was measured at 517 nm and activity was expressed as percentage. IC<sub>50</sub> value was also determined by graph<sup>12</sup>. DPPH scavenging relative to control using the following equation:

$$\text{DPPH scavenging activity (\%)} = \frac{(\text{Absorbance of control} - \text{Absorbance of sample})}{\text{Absorbance of control}} \times 100$$

#### Statistical Analysis:

The results obtained were expressed as mean ±SD and student t-test of three determinations and also statistically analyzed to ascertain its significance. The significance was estimated at (p≤0.05level).

### III. RESULTS AND DISCUSSION

Table.1: Antioxidant potential of *Withania somnifera* root and *Withania Coagulans* fruit aqueous extract

Antioxidants	<i>Withania Somnifera</i> root	<i>Withania Coagulans</i> fruit
TPC (mgGAE/ml)	33.1±0.82	14.5±0.78*
TFC (mgQE/ml)	1.86 ±0.01	1.08±1.7 <sup>ns</sup>

(n=3) Values are expressed as means±SD.\* significant, ns-non-significant when compared with *W.coagulans* aqueous extract at P ≤0.05. TPC-,Total Phenol Content TFC-Total Flavonoids Content

It has been documented that phenols as well as flavonoids show antioxidant activity and their effects on human nutrition and health through scavenging or chelating process<sup>13</sup>. In the present study, total phenols content of *Withania somnifera* and *Withania coagulans* aqueous

extract expressed in Gallic equivalents (GAE) were found to be  $33.1 \pm 0.82$  and  $14.53 \pm 0.78$  GAE mg/g respectively as shown in Table 1. The data showed that *somnifera* root powder was significantly increased by 56.19% from *coagulans* fruit aqueous extract at  $P \leq 0.05$  levels. The flavonoids content was measured by aluminum chloride technique in terms of quercetin equivalent and found low value in both extracts of *Withania somnifera* and *Withania coagulan* had  $1.86 \pm 0.01$  and  $1.08 \pm 1.7$  (mgQE/ml) respectively. *W. somnifera* aqueous extract was insignificantly increased by at 22.54% at  $p \leq 0.05$  when compared to *W. coagulan*. According to study based on phytochemical analysis showed that *W. somnifera* had

$180.80 \pm 0.01$  mg/100mg gallic acid equivalent phenolic content whereas flavonoid content was  $136.97 \pm 0.01$  mg/100 mg quercetin equivalent in *Withania somnifera* root methanolic extract<sup>14</sup>.

The medicinal effects described in the ayurveda that traditional plants have excellent phenolic acids and flavonoids content that are important ingredients to prevent against oxidative stress related disorders<sup>15</sup>. The results acquired in this study thus suggest the identified bioactive constituents in both *Withania* species is proving to be a valuable reservoir of bioactive compounds of substantial medicinal merit.




Fig.1: Scavenging effects of the aqueous extract of *Withania somnifera* and *Withania Coagulans* on DPPH at Different concentration.

DPPH free radical scavenging method is a sensitive way to determine the antioxidant activity of plant extracts. The reduction capability of free radicals was determined by decrease in its absorbance at 517nm<sup>16</sup>. Several concentrations ranging from 20–200 µg/ml of the aqueous extract of *Withania somnifera* and *Withania coagulan* were tested for their antioxidant activity in different in vitro models. IC<sub>50</sub> value is defined as the concentration of substrate that causes 50% loss of the DPPH activity and was calculated by linear regression mentioned of plots of the percentage of antiradical activity against the concentration of the tested compounds<sup>17</sup>. The percentage scavenging at

IC<sub>50</sub> values were intended for both extracts were illustrated in Fig 1. It shows that there was a decrease in the concentration of DPPH radicals due to the scavenging ability of the soluble constituents in the aqueous extract of *W.somnifera* and *W.coagulans* and the standard ascorbic acid as a reference compound. As the concentration increases the free radical scavenging activity of plant extract also increases. The data depicts that IC<sub>50</sub> value was 54µg/ml for *W. somnifera* and 69µg/ml for *W. coagulans* which was lower value according to data shown by Shariar<sup>18</sup> that methanol extract of *Withania somnifera* root showed IC<sub>50</sub> of 267.818 µg/ml. Similar data stated by Nadia

Alam<sup>19</sup> that the *Withania somnifera* extracts was 101.73-801.93 µg/ml at IC<sub>50</sub> which indicated higher value from present data.

#### IV. CONCLUSION

Although both *Withania* species had potent antioxidant activity but from the study data indicated that *Withania somnifera* aqueous root extract showed higher bioactive compounds as well as DPPH radical scavenging activity when compared to *Withania coagulans* fruit aqueous extract. Thus, *W. somnifera* root will definitely serve as a high-quality phytotherapeutic agent in various metabolic and degenerative diseases.

#### V. FUTURE PROSPECTIVE

The phytochemistry and pharmacology characteristics of both *Withania* species have been widely investigated but the studies on toxicology of the extracts in different solvents are very few. So, further exploration is needed for identification and isolation of the particular compound responsible for the specific activity.

#### CONFLICT OF INTEREST STATEMENT

We declare that we have no conflict of interest.

#### ACKNOWLEDGEMENT

Authors are thankful to Professor, Aditya Shastri (Vice Chancellor) for providing all the required facilities to Food Science and Nutrition Department, Banasthali University that helped us for the successful completion of the project work.

#### REFERENCES

- [1] M. Modak, P. Dixit, J.Londhe, S. Ghaskadbi, and A.D.T Paul. "Indian herbs and herbal drugs used for the treatment of diabetes," *Journal of Clinical Biochemical Nutrition*. 3<sup>rd</sup>ed, vol 40, pp.163-173, 2007.
- [2] S.M Mukhtar, L. Deslandes, M.C Auriac, Y. Marco, and I.E Somssich. "The Arabidopsis transcription factor WRKY27 influences wilt disease symptom development caused by *Ralstoniasolanacearum*," *Plant Journal*. vol56, pp.935-947, 2008.
- [3] S.S Negi. "Determining functionally important amino acid residues of the E1 protein of Venezuelan equine encephalitis virus," *J Mol Model*. vol12, pp.921-929, 2006.
- [4] S.A Bhat, S.A Lone, and S.S Ahmad. "Biocontrol of Damping off in *Withania Somnifera* (L) Dunal," *International Research Journal of Microbiology*. 4<sup>th</sup>ed, vol 5, pp.73-79, 2014.
- [5] A. Bano, N. Sharma, H.S. Dhaliwal, and V Sharma. "A Systematic and Comprehensive Review on *Withania somnifera*(L.) Dunal- An Indian Ginseng," *British Journal of Pharmaceutical Research*. 2<sup>nd</sup>ed, vol. 7, pp.63-75, 2015.
- [6] J. Dabheliya, S.A Khan, M.Joshiyura, M.Vasoys, S. Patel, and Vijaya S. "Diuretic potential of aqueous extract of fruits of *Withania coagulans*Dunal in experimental rats," *International Journal of Pharmacy of Pharmaceutical Science*. pp.51-53, 2010.
- [7] R. Maurya. "Chemistry and pharmacology of *Withania coagulans*: an Ayurvedic remedy," *Journal of Pharmacy and pharmacology*. pp 153-160, 2012.
- [8] R. Jain, S. Kachhwaha, and S.L Kothari. "Phytochemistry, pharmacology, and biotechnology of *Withania somnifera* and *Withania coagulans*: A review *Journal of medicinal plants research*," vol 6, pp.5388-5399, 2012.
- [9] R. Nagappan. "Evaluation of aqueous and metanol extracts of bioactive medicinal plant, *Cassia didymobotrya*(Fresenius) Irwin & barneby against immature stages of filarial vector, *Culex quinquesfasciatus* say (Diptera: Culiidae)". *Asian Pac J Trop Biomed*. 9<sup>th</sup>ed, vol 2, pp.707-711, 2012.
- [10] V.L Singleton, R.Orthofer, and R.M.L Ravento. "Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymol*," 9<sup>th</sup>ed, vol 29, pp.152-178, 1999
- [11] C.C Chang, M.H. Yang, H.M Wen, J.C Chern. "Estimation of total flavonoids content in propolis by two complementary colorimetric methods." *J. Food Drug Anal*. 3<sup>rd</sup>ed, vol 10, pp.178-182, 2002.
- [12] M.S Blois. "Antioxidant determinations by the use of a stable free radical." *Nature*. vol26, pp.1199-1200, 1958.
- [13] M. Kessler, G. Ubeaud and L. Jung. "Anti- and pro-oxidant activity of rutin and quercetin derivatives". *J Pharm and Pharmacol*. vol 55, pp.131-142, 2003.
- [14] D. Chaudhuri, N.B. Ghate, R. Sarkar, N. Mandal. "Phytochemical analysis and evaluation of antioxidant and free radical scavenging activity of *Withania somnifera* root." *Asian Journal of Pharmaceutical and Clinical Research*. 4<sup>th</sup>ed, vol 5, pp.193-199, 2012.
- [15] Franova S, Nosalova G, Mokry J, Mahmud THK, Ather A. Phytotherapy of cough. In Iwu M (ed.)

Advances in phytomedicine, U.S.A. Elsevier; pp.111-131, 2006.

- [16] G.H Katsube, and D. Woodend. "Effect of glycemic carbohydrates on short-term satiety and food intake," *Nutrition Review*. Vol61, pp.S17-S26, 2004.
- [17] L. Hemlatha. "Encyclopedia of Chemical processing, Lee," CRC Press.vol 21, pp.153-155, 2010.
- [18] M. Shahriar, Md.I Hossain, F.A. Sharmin, S. Akhter, Md.A. Haque, Md.A. Bhuiyan. In Vitro antioxidant and free radical scavenging activity of *Withania Somnifera* root. *Iosr Journal of Pharmacy*. 2<sup>nd</sup>ed, vol 3, pp.38-47, 2013.
- [19] N. Alam, M. Hossain, Md.A. Mottalib, S.A Sulaiman, S.H. Gan, Md.I. Khalil. "Methanolic extracts of *Withania somnifera* leaves, fruits and roots possess antioxidant properties and antibacterial activities," *Complementary and Alternative Medicine*. Vol 12, pp.175-183, 2012.