

Records of Arthropod Species Sampled from Avocado Plant (*Persea americana* Mill) in Small-scale Agro-ecosystems at Taita Hills and Mount Kilimanjaro

Odanga J. James^{3,2,1*}, Florence Olubayo², Richard Nyankanga², Sizah Mwalusepo^{1, 4}, Tino Johansson^{1, 5}

¹ICIPE-African Insect Science for Food and Health, P. O. Box 30772 - 00100, Nairobi, Kenya

²Department of Plant Science and Crop Protection, University of Nairobi, Kenya, P.O. Box 30197-00100, Nairobi, Kenya

³Invertebrate Zoology Section, National Museums of Kenya, P. O. Box 40658 - 00100, Nairobi, Kenya

⁴Department of General Studies, Dar es Salaam Institute of Technology, P. O. Box 2958, Dar es Salaam, Tanzania

⁵Department of Geosciences and Geography, University of Helsinki, P. O. Box 68, FI-00014, Finland

Abstract— Avocado, *Persea americana* Mill, plays a central role in distribution of both beneficial and detrimental arthropods thereby influencing local species diversity in agro-ecosystems adjacent to Afromontane forests at Mount Kilimanjaro in North-eastern Tanzania and Taita Hills in South-eastern Kenya. However, little is known about arthropod species that inhabit avocado trees in the two study areas despite the fact that the crop forms the major part of agro-ecosystem in the East African highlands. A novel survey was, therefore, carried out for two years between August 2012 and July 2014 to establish arthropod species in avocado orchards along South-eastern slopes of both Mount Kilimanjaro and Taita Hills. A total of sixty one species of arthropods were recorded from the avocado crop through fruit observation and canopy sampling. The present arthropod checklist provides baseline knowledge for scientists in evaluating beneficial and pest status of each species inhabiting avocado plant in the East African agro-ecosystems.

Keywords—Avocado, arthropods, East Africa, Mount Kilimanjaro, Taita Hills.

I. INTRODUCTION

Avocado, *Persea americana* Mill (Lauraceae), is an important crop in the world as it enhances both agro-forestry conservation concept and nutritional security (Griesbach, 2005; Bergh, 1992). The avocado trees thrives well in agro-ecosystems with relatively high altitude between 1000m a.s.l and 2600m a.s.l that receive average annual precipitation ranging from 120mm to 160mm and

average temperature of 21°C (Griesbach, 2005; Wasilwa *et al.*, 2004; Whiley, 2002). Such ecosystems are located near indigenous forests with favourable agricultural conditions as exemplified by Afrotropical highlands at Mount Kilimanjaro and Taita Hills where avocado is the dominant fruit crop (Griesbach, 2005; Wasilwa *et al.*, 2004). However, the potential land area available for avocado farming along slopes of Mount Kilimanjaro in North-eastern Tanzania and Taita Hills in South-eastern Kenya is shrinking as a result of ecological degradation (Conte, 2010) and human activities. The envisaged reduction of avocado orchards in these East African highlands will not only affect distribution of arthropod species but also livelihood of local farmers that depend on avocado fruits as a source of cash and nutritious food (Hemp, 2009).

Eight five percent of avocado production in Kenya and Tanzania is at small scale level with number of trees per farm varying from three to twelve where the crop is grown mainly for subsistence and local markets (Griesbach, 2005; Wasilwa *et al.*, 2004). Unlike commercial plantations, the small-holder avocado cropping systems do not utilized modern-day agricultural technology leading to poor farming practices, possible increase in arthropod pests and reduction of related natural enemies (Ware *et al.*, 2016; Ware *et al.*, 2012; Mwatawala *et al.*, 2009, Griesbach, 2005; Bale *et al.*, 2002; Bergh, 1992). Moreover, there is limited information on arthropods inhabiting avocado crop in Kenya and Tanzania. This paper was, therefore, initiated to provide a checklist of arthropods sampled from different parts of avocado crop in farmlands at Mount Kilimanjaro and Taita

Hills. Establishing checklist of arthropods inhabiting avocado plant in East Africa can furnish important baseline information on pest and beneficial status of each species

II. MATERIALS AND METHODS

2.1. Study areas

The study was carried out in avocado farmlands adjacent to East African montane forests at Mount Kilimanjaro in North-eastern Tanzania and Taita Hills in South-eastern Kenya (Fig 1a and b, respectively). Avocado crop is grown in the two study transects along altitudinal gradient from 900 to 2000m a.s.l. and form major part of agro-ecosystem in the region. Mount Kilimanjaro and Taita Hills are the first uppermost elevated montane forms inland from the Indian Ocean and these highlands are important catchment areas for surrounding lowland areas of Moshi and Voi in

Tanzania and Kenya, respectively (Hemp 2006a; Hemp 2006b; Bytebier, 2001; Bennun & Njoroge, 1999). The two study areas; Taita Hills and Mount Kilimanjaro, are situated 90km apart and both are about 150km from Indian Ocean. Mount Kilimanjaro study area is located between 03° 378'S, 37° 450'E and 03° 481'S, 37° 456'E in North-eastern Tanzania (Fig 1a). Mean elevation of Mount Kilimanjaro study area is 1372.69m a.s.l, average annual rainfall was 107.83mm, an average annual temperature of 20.14°C with a mean annual humidity of 78.97%. Taita Hills study area is located in South-eastern Kenya, 25km west of Voi town in the Taita-Taveta County between 03° 481'S, 38° 378'E and 03° 402'S, 38° 296'E (Fig 1b). Mean elevation of Taita Hills study area is 1397.02m a.s.l, average annual rainfall was 135.19mm; mean annual temperature was 19.56°C with a mean annual humidity of 81.46%.




Fig.1a: Map of Mount Kilimanjaro study area in North-eastern Tanzania.




Fig.1b: Map of Taita Hills study area in South-eastern Kenya.

2.2. Sampling design

Species of arthropods were sampled randomly from avocado plants for two consecutive years between August 2012 and July 2014 along each study transect; Mount Kilimanjaro and Taita Hills. A transect comprised of fifteen blocks with each block consisting of at least a hundred avocado trees. During each survey, five avocado trees from the hundred sampling unit of the plants at each block were randomly examined for arthropods using protocol described by Ekesi *et al* (2006), Stibick (2006), Palmer (1990) and Moritz *et al* (2013). Leaves and flowers were gently shaken on a tray to sample species that inhabits plant parts as described by Palmer (1990). However, avocado fruits were observed for other arthropod species as described by Ekesi *et al* (2006). Some species were handpicked from the avocado the plant using fine forceps and aspirator following protocols described by Millar *et al* (2000). The collected

specimens were preserved in vials containing 60% ethyl alcohol and later taxonomically identified at the National Museums of Kenya (NMK) entomology laboratory in Nairobi.

2.3. Statistical analysis

Rank abundance test was used to categorize arthropod species based on their population using Biodiversity-R software (R Development Core Team, 2012). Species accumulation curves were used to compare if the observed species richness at two study areas along slopes of Mount Kilimanjaro and Taita Hills reach saturation point (R Development Core Team, 2012). In order to check for the completeness of sampling, observed numbers of species were compared with projected ones (R Development Core Team, 2012). The estimated species richness was constructed for each study area using non-parametric

estimators; Chao, Jackknife 1 and Bootstrap (R Development Core Team, 2012; Crawley, 2007; Crawley, 2005).

III. RESULTS

A total of sixty one species of arthropods were recorded inhabiting avocado plants in the farmlands adjacent to Afromontane forests within Mount Kilimanjaro in North-eastern Tanzania and Taita Hills in South-eastern Kenya.

Further analysis using Rank abundance test revealed that most abundant arthropod species in avocado cropping systems at Mount Kilimanjaro and Taita Hills were; *Bactrocera (invadens) dorsalis* (Hendel) (Diptera: Tephritidae), *Thaumatotibia leucotreta* Meyrick (Lepidoptera: Tortricidae), *Frankliniella schultzei* Trybom (Thysanoptera: Thripidae) and *Heliothrips haemorrhoidalis* Bouche (Thysanoptera: Thripidae) (Table 1).

Table.1: Arthropod species recorded on avocado crop in the two study areas of Taita Hills and Mount Kilimanjaro for two years between August 2012 and July 2014. T = Taita Hills, K = Mount Kilimanjaro study area where the species were sampled. * = equals or less than 0.1 log abundance.

Rank	Scientific name	Common name	Plant part sampled	Log abundance	Order	Habitat sampled
1	<i>Bactrocera (invadens) dorsalis</i>	Asian invasive fruit fly	Ground collected ripened fruits	5.9	Diptera	T and K
2	<i>Thaumatotibia leucotreta</i>	False codling moth	Immature fruits	4.9	Lepidoptera	T and K
3	<i>Frankliniella schultzei</i>	common blossom thrips	Flowers	3.2	Thysanoptera	T and K
4	<i>Heliothrips haemorrhoidalis</i>	Greenhouse thrips	Leaves and young fruits	2.9	Thysanoptera	T and K
5	<i>Megalothrips sjostedti</i>	Cowpea flower thrips	Flowers and leaves	2.8	Thysanoptera	T and K
6	<i>Thrips austarlis</i>	Western flower thrips	Flowers and leaves	2.6	Thysanoptera	T and K
7	<i>Thrips pusillus</i>	Thrips	Flowers and leaves	2.5	Thysanoptera	T and K
8	<i>Aleyrodicus dispersus</i>	Spiralling whitefly	Leaves	2.4	Hemiptera	T and K
9	<i>Haplothrips gowdeyi</i>	Thrips	Flowers and leaves	2.4	Thysanoptera	T and K
10	<i>Cheilomenes sulphurea</i>	Ladybird beetle	Flowers and leaves	2.3	Coleoptera	T and K
11	<i>Pheidole megacephala</i>	Sugar ant	Flowers and leaves	2.2	Hymenoptera	T and K
12	<i>Rhinocoris</i> sp.	Assassin bug	Flowers and leaves	2.1	Hemiptera	T and K
13	<i>Trialeurodes vaporariorum</i>	Greenhouse whitefly	Leaves	2	Hemiptera	T and K
14	<i>Tetranychus</i> sp.	Red spider mite	Flowers and leaves	1.8	Trombidiformes	T and K
15	<i>Bactrothrips</i> sp.	Thrips	Flowers and leaves	1.7	Thysanoptera	T and K
16	<i>Nezara viridula</i>	southern green stink	Flowers and leaves	1.7	Hemiptera	T and K

		bug				
17	<i>Helopeltis schoutedeni</i>	Mirid (plant bugs)	Flowers and leaves	1.7	Hemiptera	T and K
18	<i>Cheilomenes lunata</i>	Ladybird beetle	Flowers and leaves	1.7	Coleoptera	T and K
19	<i>Thrips abyssiniae</i>	Thrips	Flowers and leaves	1.7	Thysanoptera	T and K
20	<i>Franklinothrips</i> sp.	Thrips	Flowers and leaves	1.6	Thysanoptera	T and K
21	<i>Probosciodoris fuliginosus</i>	Bugs	Flowers and leaves	1.6	Hemiptera	T and K
22	<i>Franklinothrips megalops</i>	Predatory thrips and mimics ant	Flowers and leaves	1.5	Thysanoptera	T and K
23	<i>Dolicholepta jeanneli</i>	Thrips	Flowers and leaves	1.4	Thysanoptera	T and K
24	<i>Gynaikothrips</i> sp.	Thrips	Flowers and leaves	1.3	Thysanoptera	T and K
25	<i>Chilothrips frontalis</i>	Thrips	Flowers and leaves	1.3	Thysanoptera	T and K
26	<i>Dendrothrips</i> sp.	Thrips	Flowers and leaves	1.3	Thysanoptera	T and K
27	<i>Scirtothrips dorsalis</i>	Chilli thrips or yellow tea thrips	Flowers and leaves	1.3	Thysanoptera	T and K
28	<i>Apterygothrips</i> sp	Thrips	Flowers and leaves	1.2	Thysanoptera	T and K
29	<i>Scirtothrips</i> sp1	Thrips	Flowers and leaves	1.2	Thysanoptera	T and K
30	<i>Frankliniella occidentalis</i>	Thrips	Flowers and leaves	1.2	Thysanoptera	T and K
31	<i>Haplothrips</i> sp.	Thrips	Flowers and leaves	1.0	Thysanoptera	K
32	<i>Gigantothrips</i> sp.	Thrips	Flowers and leaves	1.0	Thysanoptera	T and K
33	<i>Microcephalothrips abdominalis</i>	Thrips	Flowers and leaves	1.0	Thysanoptera	T and K
34	<i>Vuilletia houardi</i>	Thrips	Flowers and leaves	0.9	Thysanoptera	T and K
35	<i>Ceratothripoides brunnens</i>	Tomato thrips	Flowers and leaves	0.8	Thysanoptera	T and K
36	<i>Scirtothrips</i> sp.2	Thrips	Flowers and leaves	0.8	Thysanoptera	T and K
37	<i>Ecacanthothrips tibialis</i>	Thrips	Flowers and leaves	0.7	Thysanoptera	T
38	<i>Thrips revelatus</i>	Thrips	Flowers and leaves	0.7	Thysanoptera	T
39	<i>Sericothrips</i> sp.	Thrips	Flowers and leaves	0.6	Thysanoptera	T

40	<i>Neosmerinthothrips sp</i>	Thrips	Flowers and leaves	0.5	Thysanoptera	T
41	<i>Diarthrothrips sp.</i>	Thrips	Flowers and leaves	0.3	Thysanoptera	K
42	<i>Frankliniella williamsi</i>	Thrips	Flowers and leaves	0.3	Thysanoptera	T
43	<i>Rhipiprothrips sp.</i>	Thrips	Flowers and leaves	0.3	Thysanoptera	T and K
44	<i>Stenchaetothripssp.</i>	Thrips	Flowers and leaves	0.3	Thysanoptera	T
45	<i>Elaphrothrips sp.</i>	Thrips	Flowers and leaves	*	Thysanoptera	T
46	<i>Pselaphothrips pomeroyi</i>	Thrips	Flowers and leaves	*	Thysanoptera	T
47	<i>Stephanothrips sp.</i>	Thrips	Flowers and leaves	*	Thysanoptera	T
48	<i>Urothripine sp.</i>	Thrips	Flowers and leaves	*	Thysanoptera	T
49	<i>Craspedothrips sp.</i>	Thrips	Flowers and leaves	*	Thysanoptera	T
50	<i>Apis mellifera</i>	Honey bee	Flowers	*	Hymenoptera	T and K
51	<i>Camponotus maculatus</i>	Ant	Flowers and leaves	*	Hymenoptera	T and K
52	<i>Oecophylla longinoda</i>	Weave ant	Leaves	*	Hymenoptera	T and K
53	<i>Crematogaster sp.</i>	Ant	Flowers and leaves	*	Hymenoptera	T and K
54	<i>Brachypeplus sp.</i>	Sap-feeding beetles	Flowers and leaves	*	Coleoptera	T and K
55	<i>Corynasp.</i>	Pollen beetles	Flowers and leaves	*	Coleoptera	T and K
56	<i>Epitrix silvacola</i>	Leaf beetles	Flowers and leaves	*	Coleoptera	T and K
57	<i>Formicomus sp.</i>	Ant-like flower beetles	Flowers and leaves	*	Coleoptera	T and K
58	<i>Nematocerus sp.</i>	Weevils	Flowers and leaves	*	Coleoptera	T and K
59	<i>Paederus sabaceus</i>	Short-winged beetles	Flowers and leaves	*	Coleoptera	T and K
60	<i>Scymnus sp.</i>	Ladybugs	Flowers and leaves	*	Coleoptera	T and K
61	<i>Camponotus rufoglaucus</i>	Ant	Flowers and leaves	*	Hymenoptera	T and K

Species accumulation curve revealed that 59 species of arthropods were recorded at Taita Hills and 50 arthropod species at Mount Kilimanjaro (Fig 2; Table 2).




Fig.2: Species accumulation curve of arthropods recorded on avocado plants for (a) Taita Hills and (b) Mount Kilimanjaro study areas. The bars are standard errors. Sampling sites are study blocks at with each comprising of at least a hundred avocado trees.

Richness estimators (Boot, Jackknife and Chao) predicted a number of between 71 ± 3 (mean $\pm$ se) and 54 ± 2 species for Taita Hills whereas between 56 ± 2 and 43 ± 1 for Mount Kilimanjaro (Table 2).

Table.2: Observed species richness and non-parametric species richness estimators for Taita Hills and Mount Kilimanjaro transects.

Habitat	Observed species	Chao	Jackknife 1	Boot	Sampling blocks (n)
Taita Hills	59	71.51 ± 3.77	56.62 ± 2.23	54.43 ± 2.09	15
Mount Kilimanjaro	50	56.14 ± 2.81	50.20 ± 2.01	43.12 ± 1.36	15

IV. DISCUSSION

Our results revealed that the Asian invasive fruit fly (*Bactrocera dorsalis*), False codling moths (*Thaumatotibia leucotreta*), common blossom thrips (*Frankliniella schultzei*) and the Greenhouse thrips (*Heliothrips haemorrhoidalis*) are the most abundant arthropod species inhabiting avocado plant parts at Mount Kilimanjaro in Tanzania and Taita Hills in Kenya. *Bactrocera dorsalis* was recorded from ground collected ripened avocado fruits whereas *Thaumatotibia leucotreta* was sampled from immature avocado fruits confirming them as fruit a scenario that has been reported by Mwatawala *et al* (2009) and Prinsloo and Uys (2015), respectively. *Frankliniella schultzei* was recorded exclusively feeding on flower resources, however, *Heliothrips haemorrhoidalis* was recorded from leaves and young fruits proving to be pests of avocado crop as reported in Palmer (1990) and Prinsloo and Uys (2015). Beneficial or pest status of each of the species sampled was not evaluated in this novel paper since our study focused on generating the checklist of arthropods that inhabits avocado plant. However, Ware *et al* (2016), Prinsloo and Uys (2015), Ware *et al* (2012) and Mwatawala *et al* (2009) reported that *Bactrocera dorsalis* and

Thaumatotibia leucotreta are economically important pests of avocado fruits. Moreover, beneficial insects such as honey bees (*Apis mellifera*) was exhaustively described pollinating avocado flowers at Taita Hills in South-eastern Kenya by Luvanga (2015).

The failure of the species accumulation curves to reach saturation point indicated that more species were likely to be found if additional sampling effort continued at Taita Hills transect than at Mount Kilimanjaro study area. This hypothesizes that if surveys are conducted over multiple years, the species accumulation curves for arthropods are expected to reach saturation point probably due to a larger proportion of rare species which are often recorded in single encounters (Novotny and Basset, 2000). Most of the rare species in this study were Thysanopteran non-pest thrip species that contributed to the high projection of species richness for Taita Hills using Chao; a non-parametric estimator (R Development Core Team, 2012).

V. CONCLUSIONS

Relatively high species richness of arthropods sampled from a single crop confirmed that the avocado plant also plays vital ecosystem functioning at both Mount Kilimanjaro and

Taita Hills study areas. The present checklist provides baseline data for scientists to identify species of economic importance which can assist in designing strategies for avocado pest control and pollination services in East Africa. In order to enhance avocado farming strategies at the two study areas, spatiotemporal data is required for each species recorded. This study, therefore, recommends research on how the most abundant arthropods (*Bactrocera dorsalis*, *Thaumatotibia leucotreta*, *Frankliniella schultzei* and *Heliothrips haemorrhoidalis*) impacts on avocado farming at Mount Kilimanjaro and Taita Hills. The study, further, recommends research on possible beneficial insects such as *Apis mellifera* (Hymenoptera: Apidae), *Oecophylla longinoda* (Hymenoptera: Formicidae) and *Camponotus maculatus* (Hymenoptera: Formicidae) in the avocado orchards along slopes of Mount Kilimanjaro and Taita Hills.

ACKNOWLEDGMENT

We gratefully acknowledge the financial support for this research by the Ministry for Foreign Affairs of Finland to the African Insect Science for Food and Health (*icipe*). University of Dar es Salaam, Taita - Taveta county government, the late Joseph R. Mugambi, Granton Mwadime, Filbert Mlingi, Calistus Omondi, Peterson Nderitu and Martin Gathendu provided technical advice and support during fieldwork period in Kenya and Tanzania. The first author of this study is a doctoral research fellow working under “Climate change impacts on ecosystem services and food security in eastern Africa” (CHIESA) project/ avocado cluster.

REFERENCE

- [1] Bale, J., Masters, G., Hodkinson, I., Awmack, C., Bezemer, T., Brown, V., Butterfield, J., Buse, A., Coulson, J., Farrar, J. (2002). Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology* 8: 1 - 16.
- [2] Bennun, L., Njoroge, P. (1999). Important bird areas in Kenya. Nature Kenya, Nairobi.
- [3] Bergh, B.O. (1992). The origin, nature and genetic improvement of avocado *California Avocado Society Yearbook* 76: 61 - 75.
- [4] Bytebier, B. (2001). Taita Hills Biodiversity Project Report. 121 pp. National Museums of Kenya, Nairobi.
- [5] Conte, C.A. (2010). Forest history in East Africa's Eastern Arc Mountains: Biological science and the uses of history. *BioScience* 60(4): 309 - 313.
- [6] Crawley, M. (2005). Statistics, an introduction to R. John Wiley & Sons Ltd, West Sussex, England.
- [7] Crawley, M.J. (2007). The R book. Chichester, UK: John Wiley and Sons, Ltd.
- [8] Ekesi, S., Billah, M.K. (2006). A Field guide to the management of economically important tephritid fruit flies in Africa. ICIPE Science Press, Nairobi, Kenya.
- [9] Griesbach, J. (2005). Avocado growing in Kenya. World Agroforestry Centre (ICRAF). Kul Graphics Ltd. Nairobi, Kenya.
- [10] Hemp, A. (2006a). Continuum or zonation? Altitudinal gradients in the forest vegetation of Mt. Kilimanjaro. *Plant Ecology* 184: 27 - 42.
- [11] Hemp, A. (2006b). Vegetation of Kilimanjaro: hidden endemics and missing bamboo. *African Journal of Ecology* 44: 305 - 328.
- [12] Hemp, A. (2009). Climate change and its impact on the forests of Kilimanjaro. *African Journal of Ecology* 47(1): 3 - 10.
- [13] Luvonga, E.B. (2015). Diversity and pollination activity of flower visiting insects associated with avocado along the slopes of Taita hills in Kenya. Masters thesis. Masinde Muliro University of Science and Technology, Kenya.
- [14] Millar, I.M., Uys, V.M., Urban, R.P. (2000). Collecting and preserving insects and arachnids: a manual for entomology and arachnology. ARC – Plant Protection Research Institute Pretoria, South Africa.
- [15] Moritz, G., Brandt, S., Triapitsyn, S., Subramanian S. (2013). Pest thrips in East Africa - Identification and information tools (CD-ROM). QBIT, QAAFI Biological Information Technology, The University of Queensland, Australia., ISBN: 978-1-74272-067-8.
- [16] Mwatawala, M.W., De Meyer, M., Makundi, R.H., Maerere, A.P. (2009b). Host range and distribution of fruit-infesting pestiferous fruit flies (Diptera, Tephritidae) in selected areas of Central Tanzania. *Bulletin of Entomological Research* 99(6): 629 - 641.
- [17] Novotny, V., Basset Y. (2000). Rare species in communities of tropical insect herbivores: pondering the mystery of singletons. *Oikos* 89: 564 - 572.
- [18] Palmer, J.M. (1990). Identification of common thrips of tropical Africa (Thysanoptera:Insecta). *Tropical Pest Management* 36(1): 27 - 49.
- [19] Prinsloo, G.L., Uys, V.M. (2015). Insects of cultivated plants and natural pastures in Southern Africa. Entomological Society of Southern Africa, Hatfield, South Africa.
- [20] R Development Core Team. (2012). R: A language and environment for statistical computing. R

-
- Foundation for Statistical Computing, Vienna. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- [21] Stibick, J. (2006). New Pest Response Guidelines: False Codling Moth *Thaumatotibia leucotreta*. USDA-APHIS-PPQ-Emergency and Domestic Programs, Riverdale, Maryland [http://www.aphis.usda.gov/import_export/plants/ppq_manuals.shtml].
- [22] Ware, A.B., du Toit, C.L.N, du Toit, E., Collins, R., Clowes, R., Ekesi, S., Mohamed S. (2016). Host suitability of three avocado cultivars (*Persea americana* Miller: Lauraceae) to oriental fruit fly (*Bactrocera (invadens) dorsalis* (Hendel) (Diptera: Tephritidae). *Crop Protection* 90: 84 - 89.
- [23] Ware, A.R., Du Toit, C.L.N., Mohamed, S.A., Nderitu, P.W., Ekesi, S. (2012). Cold Tolerance and Disinfestation of *Bactrocera invadens* (Diptera:Tephritidae) in 'Hass' Avocado. *Journal of Economic Entomology*, 105(6): 1963 - 1970.
- [24] Wasilwa, L.A., Njuguna, J.K., Okoko, E.N., Watani, G.W. (2004). Status of Avocado Production in Kenya. Kenya Agricultural Research Institute, Nairobi, Kenya.
- [25] Whiley, A.W. (2002). Crop management. In: Whiley AW, Schaffer B, Wolstenholme BN (eds) Avocado: Botany, Production and Uses. CABI Publishing, Oxon UK, pp 231-258.