

SHORT COMMUNICATION

Spatial characterization of common blossom thrips (*Frankliniella schultzei*) in smallholder avocado orchards along slopes of Taita Hills and Mount Kilimanjaro

Odanga J. James^{1, 2a*, 6}, Samira Mohamed¹, Florence Olubayo⁶, Richard Nyankanga⁶, Irine A. Otieno³, Sizah Mwalusepo^{1, 4}, Geoffrey Mwachala^{2b}, Tino Johansson^{1, 5}

¹ICIPE-African Insect Science for Food and Health, P. O. Box 30772 - 00100, Nairobi, Kenya

^{2a}Invertebrate Zoology Section, National Museums of Kenya, P. O. Box 40658 - 00100, Nairobi, Kenya

^{2b}East African Herbarium, National Museums of Kenya, P.O. Box 40658 - 00100, Nairobi, Kenya

³Faculty of Geosciences, Utrecht University, Netherlands

⁴Department of General Studies, Dar es Salaam Institute of Technology, P. O. Box 2958, Dar es Salaam, Tanzania

⁵Department of Geosciences and Geography, University of Helsinki, P. O. Box 68, FI-00014, Finland

⁶Department of Plant Science and Crop Protection, University of Nairobi, Kenya, P.O. Box 30197-00100, Nairobi, Kenya

*Corresponding author: jkodss@yahoo.com

Abstract – *Frankliniella schultzei* Trybom (Thysanoptera: Thripidae) is an important flower pest of avocado crop (*Persea americana* Mill) at Taita Hills in South-eastern Kenya and Mount Kilimanjaro in North-eastern Tanzania. However, its geographical distribution is not known in the East African avocado cropping systems. In order to generate the spatial data of the common blossom thrips (*Frankliniella schultzei*), a survey was carried out in smallholder avocado orchards along altitudinal gradient (900 - 1800m.a.s.l.) of Taita Hills and Mount Kilimanjaro using a white coloured beating tray and camel brush. Once the specimens of thrips were taxonomically verified, the abundance data was tabulated into three altitudinal zones, namely; lowland (900-1199m.a.s.l.), sub-montane (1200-1499m.a.s.l.) and montane (1500-1799m.a.s.l.). *Frankliniella schultzei* was recorded in all altitudinal zones of both transects with mean abundance being highest at Taita Hills (5.4) compared to Mount Kilimanjaro (0.9). However, abundance of the pest was greater in cooler highlands (>1200m.a.s.l.) than warmer lowland areas (<1200m.a.s.l.) of both transects. The findings of this study contributes significantly towards spatial mapping of *Frankliniella schultzei* in East Africa and this information is important in developing strategies aimed at controlling infestation of avocado flowers by the insect pest at the two study transects.

Keywords— *Avocado, East Africa, Frankliniella schultzei, Mount Kilimanjaro, Taita Hills.*

I. INTRODUCTION

Common blossom thrips (*Frankliniella schultzei* Trybom) is an anthophilous pest species (Milne *et al.*, 1996; Odanga *et al.*, 2017b) whose diet is predominantly pollen and floral tissues (Kakkar *et al.*, 2012). Being a polyphagous insect pest, it feeds on flowers of various ornamental, vegetable and fruit crop hosts in different parts of the world (Milne *et al.*, 1996; Kakkar *et al.*, 2012). Palmer (1990), Palmer (1992) and Milne & Walter (2000) reported *Frankliniella schultzei* on 83 species of plants from 35 families with important hosts being cotton (*Gossypium* spp.), groundnut (*Arachis hypogaea*), beans (*Phaseolus vulgaris*) and pigeon pea (*Cajanus cajan*) and avocado (*Persea americana*). In Afrotropical highlands of Taita Hills and Mount Kilimanjaro, *Frankliniella schultzei* is a pest of avocado crop that feeds on floral resources (Odanga *et al.*, 2017a; Odanga *et al.*, 2017b) thereby contributing to flower abortion and subsequent low fruit-set. This pest, therefore, impacts negatively on livelihood of small-scale farmers as it contributes to low yield of avocado fruits which the local growers depend on as a source of cash and nutritious food. Although, Odanga *et al* (2017b) described in detail temporal fluctuations of the pest abundance at Taita Hills and Mount

Kilimanjaro, limited information is available on geographical distribution of *Frankliniella schultzei* in the East African highlands. This study was, therefore, initiated to provide first-ever spatial distribution data of *Frankliniella schultzei* in avocado orchards along altitudinal gradient of Taita Hills in South-eastern Kenya and Mount Kilimanjaro in North-eastern Tanzania.

II. MATERIALS AND METHODS

2.1. Study areas

This study was carried out in farmlands at Taita Hills in South-eastern Kenya and Mount Kilimanjaro in Northern-eastern Tanzania as described by Mwalusepo *et al* (2015) and Odanga *et al* (2017b). The study regions were selected because the avocado plant (*Persea americana* Mill) is the major fruit crop cultivated by the small-holder farmers at South-eastern slopes of Taita Hills and Mount Kilimanjaro. Furthermore, farming along the two study transects is rainfed and the small-holder growers do not use chemicals to control insect pests or diseases of avocado crop.

2.2. Study design

The survey was carried out along altitudinal gradient from 900 to 1800 m.a.s.l. in small-scale avocado orchards at the two study areas between August 2012 and July 2014. The common blossom thrips, *Frankliniella schultzei* Trybom, was sampled from avocado trees using a white coloured beating tray and camel brush as described by Palmer (1990). Six hundred avocado trees were sampled at each altitudinal zone, namely; lowland region (900-1199m a.s.l.), sub-montane (1200-1499m a.s.l.) and montane (1500-1799m a.s.l.). Geographical coordinates and elevation of every study site was verified using a hand-held Garmin GPS model eTrex 30. Collected specimens of the thrips were mounted and identified at the National Museums of Kenya entomology laboratory in Nairobi using taxonomic manuals; Palmer (1990), Palmer *et al* (1992), Moritz *et al* (2001) and Mound (2010). The fully identified and

confirmed thrip species were deposited in the entomology collection at the National Museums of Kenya.

2.3. Data analysis

Wilcoxon signed rank test was employed to test differences between paired datasets sampled at Taita Hills and Mount Kilimanjaro (R Development Core Team, 2012; Crawley, 2007). Sets of variables were normalized for further analysis using Tukey's HSD (Honestly Significant Difference) post hoc test to pinpoint what exact sub-sets within a data that had significant differences from each other (R Development Core Team, 2012). Spatial mapping was generated using kriging method by interpolating mean abundance of *Frankliniella schultzei* along altitudinal gradient of the two transects using QGIS version 1.8.0.

III. RESULTS

Mean abundance of *Frankliniella schultzei* differed significantly between Taita Hills (5.4 ± 0.8) and Mount Kilimanjaro (0.9 ± 0.1) ($V=1726.4$, $P<0.0001$). For both transects, mean abundance of *Frankliniella schultzei* was smaller at lowland zone (900-1199m a.s.l.) than the highlands. However, the abundance of *Frankliniella schultzei* at Taita Hills study area was highest at mid-altitudinal range (1200-1499m a.s.l.) (Figure 1a). For Mount Kilimanjaro, the abundance was highest at montane zone (1500-1799m a.s.l.) (Figure 1b). Tukey's HSD pair wise comparison of mean abundance of *Frankliniella schultzei* between agro-ecological zones at Taita Hills revealed a significant difference in two pairs (lowland and sub-montane, $P<0.0001$; sub-montane and montane, $P<0.0001$) except between montane and lowland ($P=0.344$) (Figure 1a). For Mount Kilimanjaro transect, the mean abundance of *Frankliniella schultzei* between agro-ecological zones revealed a significant difference in only one pair; lowland and montane ($P=0.011$), however, the rest did not show a significant difference (lowland and sub-montane, $P=0.312$; sub-montane and montane, $P=0.317$) (Figure 1b).




Fig.1: Distribution of *Frankliniella schultzei* within altitudinal zones of (a) Taita Hills and (b) Mount Kilimanjaro transects. Similarly, spatial distribution pattern for *Frankliniella schultzei* using kriging method revealed that the pest is highly abundant in highlands above 1200m a.s.l than lowlands of both Taita Hills and Mount Kilimanjaro study areas (Figure 2a & b).




Fig.2: Geographical distribution of *Frankliniella schultzei* based on its mean abundance along altitudinal gradient of (a) Taita Hills in South-eastern Kenya and (b) Mount Kilimanjaro in North-eastern Tanzania. Red pattern shows highest whereas green reveals least mean abundance of *Frankliniella schultzei*. Dotted red points are the sampling sites along each study transect.

IV. DISCUSSION

Mean abundance of common blossom thrips (*Frankliniella schultzei*) was greater in highlands above 1200m.a.s.l because these zones had plentiful host avocado trees with abundant floral resources at both study transects. Availability of productive host trees at highlands provided the pest with enough food and habitat (Palmer *et al.*, 1992). Consequently, the abundance of *Frankliniella schultzei* was very low at the lowland zone (1200m.a.s.l.) because the region is warmer compared to ever cold highlands.

Generally, the common blossom thrips thrives well in areas with mild temperatures (Milne *et al.* 1996; Kakkar *et al.* 2012; Palmer *et al.*, 1992) which were available in sub-montane (1200m – 1600m a.s.l) and montane (1500-1799m a.s.l) zones at both Taita Hills and Mount Kilimanjaro agro-ecosystems. However, distribution patterns of *Frankliniella schultzei* along altitudinal gradient of Taita Hills and Mount Kilimanjaro may shift drastically in future with changing climate and associated agricultural activities.

V. CONCLUSION

The common blossom thrips (*Frankliniella schultzei*) is major flower pest of avocado crop and the insect species is well established along altitudinal gradient of Taita Hills and Mount Kilimanjaro. In order to enhance productivity of the avocado trees, control measures for the pest should be focused in all altitudinal zones at both transects. However, applied research is required to develop a universal protocol for indexing floral infestation levels by *Frankliniella schultzei* in East African avocado orchards. Regional studies that integrate biogeographical approaches to predict shifts in distribution and exact pest status of *Frankliniella schultzei* as a function of future climate change are highly recommended.

ACKNOWLEDGMENTS

We gratefully acknowledge the financial support for this research by the Ministry for Foreign Affairs of Finland to the African Insect Science for Food and Health (*icipe*). Dar es Salaam University, Taita - Taveta county government, Martin Gathendu, Granton Mwadime, Filbert Mlingi, Mathew Mrema, Calistus Omondi, Peterson Nderitu and the late Joseph R. Mugambi provided technical advice and support during fieldwork period in Kenya and Tanzania. The authors would like to thank an anonymous Thysanopteran expert who taxonomically verified our specimens. The first author of this study is a research fellow working under “Climate change impacts on ecosystem services and food security in eastern Africa” (CHIESA) project/ avocado cluster.

REFERENCES

- [1] Crawley, M.J. (2007). The R book. Chichester, UK: John Wiley and Sons, Ltd.
- [2] Kakkar, G., Seal, D.R., Kumar, V. (2012). Assessing abundance and distribution of an invasive thrips *Frankliniella schultzei* (Thysanoptera: Thripidae) in south Florida. *Bulletin of entomological research* 102(3): 249 - 259.
- [3] Mound, L.A. (2010). Species of the Genus Thrips (Thysanoptera: Thripidae) from Afro-tropical region. *Zootaxa* 2423:1-24
- [4] Milne, J.R., Jhumlekhasing M., Walter, G.H. (1996). Understanding host plant relationships of polyphagous flower thrips, a case study of *Frankliniella schultzei* (Trybom). In Goodwin S, Gillespie P. (eds), Proceedings of the 1995 Australia and New Zealand Thrips Workshop: Methods, Biology, Ecology and Management, NSW Agriculture, Gosford. 8 - 14.
- [5] Milne, M., Walter, G.H. (2000). Feeding and breeding across host plants within a locality by the widespread thrips *Frankliniella schultzei*, and the invasive potential of polyphagous herbivores. *Divers.and Distri* 6: 243 - 257.
- [6] Moritz, G., Brandt, S., Triapitsyn, S., Subramanian S. (2013). Pest thrips in East Africa - Identification and information tools (CD-ROM). QBIT, QAAFI Biological Information Technology, The University of Queensland, Australia. ISBN: 978-1-74272-067-8.
- [7] Mwalusepo, S., Tonnang, H.E., Massawe, E.S., Okuku, G.O., Khadioli, N., Johansson, T., Le Ru, B.P. (2015). Predicting the impact of temperature change on the future distribution of maize stem borers and their natural enemies along East African mountain gradients using phenology models. *PloS one* 10(6).
- [8] Odanga, J.J., Olubayo, F., Nyankanga, R., Mwalusepo, S., Johansson, T. (2017a). Records of Arthropod Species Sampled from Avocado Plant (*Persea americana* Mill) in Small-scale Agro-ecosystems at Taita Hills and Mount Kilimanjaro. *International Journal of Environment, Agriculture and Biotechnology* (IJEAB) 2(5): 2457 - 2465
- [9] Odanga, J.J., Mohamed, S., Olubayo, F., Nyankanga, R., Mwalusepo, S., Subramanian, S., Johansson, T., Ekesi, S. (2017b). Datasets on abundance of common blossom thrips and weather variables in small-scale avocado orchards at Taita Hills and Mount Kilimanjaro. *Data-in Brief* (in press).
- [10] Palmer, J.M., Mound, L.A., Du Heamue, G.J. (1992). *CIE guides to insects of importance to man. 2. Thysanoptera*. Bretts, C. R. (editor). CAB international: Wallingford, UK.
- [11] Palmer, J.M. (1990). Identification of common thrips of tropical Africa (Thysanoptera:Insecta). *Tropical Pest Management* 36(1): 27 - 49.
- [12] R Development Core Team (2012). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.