

# Antifungal Activity of Petroleum and Ethanolic Extracts of *Moringa Oleifera* Leaves against *Penicillium Chrysogenum* and *Cryptococcus Neoformans*

Kale Bhagwat, Dr. N. S. Mali, Lonkar Amar, Jadhav Sourabh

Corresponding Author : Kale Bhagwat, Assistant Professor, Department of Biochemistry, Pandit Deendayal Upadhyay Dental College and Hospital, Kegaon, Dist : Solapur (Maharashtra).

**Abstract**— *Moringa oleifera* is well known medicinal plant. Its different parts are widely used for the treatment of different types of diseases since it has antibacterial and antifungal activity. The leaves are rich in iron, rhamnase and a unique group of compounds called glucosinolates and isothiocyanates. Other medical properties include antipyretic, antiepileptic, antiinflammatory, antiulcerative, antihypertensive, cholesterol lowering, antioxidant, anti diabetic. The current investigation was undertaken to evaluate the antifungal activities by petroleum ether and ethanolic extracts of *Moringa oleifera* leaves against *Penicillium chrysogenum* and *Cryptococcus neoformans*. From our study, it is found that as concentration of *Moringa* leaves extract increases the diameter of zone of inhibition found increased. The results were higher and effective for ethanolic extract than petroleum extract. The results were also found more effective against *Cryptococcus neoformans* than *Penicillium chrysogenum*.

**Keywords**— *Moringa oleifera*, *Cryptococcus neoformans*, Antifungal, Antibacterial, *Penicillium chrysogenum*.

## I. INTRODUCTION

*Moringa oleifera* (Fig :1) is well known, widely distributed natural species belongs to monogeneric family Moringaceae and order Violales.(Table:1)<sup>(1,2)</sup>

Table.1: Classification of *Moringa oleifera*

Botanical Classification	
Kingdom	Plantae
Sub-kingdom	Tracheobionta
Super-division	Spermatophyta
Division	Angiospermae/Magnoliophyta
Class	Magnoliopsida
Subclass	Dilleniidae

Order	Capparales
Family	Moringaceae
Genus	Moringa
Species	Oleifera



Fig.1: *Moringa oleifera* tree

*Moringa* tree can grow well in the humid tropic or hot dry land with average height that ranges from 5 to 10 m. It can survive in harsh climatic condition including destitute soil without being much affected by drought.<sup>(3)</sup> These are true vascular plants contains xylem and phloem for conduction of water and nutrients respectively (Sub-kingdom : Tracheobionta). Its trunk is soft, white corky and branches bearing a gummy bark. Each tripinnately compound leaves bear several small leaf legs.<sup>(4,5)</sup> *Moringa* is flowering plant(Division: Angiospermae/Magnoliophyta).The flowers

are fragrant, bisexual and pentapetalous. It contains five unequal, separate, thinly veined, yellowish-white petals.(Fig:2) Flower is syncarpous and posses united carpels(Subclass : Dilleniidae). They grow on slender, hairy stalks in spreading or drooping later flower clusters. Leaves of these plants are feathery, pale green, compound and tripinnate. They are about 30–60 cm long and contains many small leaflets.<sup>(6,7)</sup> Each leaflet is about 1 to 2 cm long and 0.5 to 1.0 cm wide. Upper surface of leaflets is smooth dark while lower is pale green. They are variable in size and shape.(Fig:3)



Fig.2: *Moringa oleifera* flowers



Fig.3: *Moringa oleifera* leaves

This plant is also known as drum stick tree or horse radish tree. It have many vernacular names such as kelor, marango, mlonge, moonga, mulangay, nebeday, saijhan, sajna or Ben oil tree<sup>(8,9)</sup> *Moringa oleifera* is well known medicinal plant. Its different parts are widely used for the treatment of different types of diseases since is have antibacterial and antifungal activity. The leaves are rich in iron, rhamnase and a unique group of compounds called glucosinolates and isothiocyanates. Other medical

properties include antipyretic, antiepileptic, antiinflammatory, antiulcerative, antihypertensive, cholesterol lowering, antioxidant, anti diabetic.<sup>(10,11)</sup> The current investigation was undertaken to evaluate the antifungal activities by petroleum ether and ethanolic extracts of *Moringa oleifera* leaves against *Penicillium crysogenum* and *Cryptococcus neoformans*.


## II. MATERIALS AND METHODS

- A) **Collection of leaves :** Naturally dried leaves were directly plucked from *Moringa* plant. The collection is carried out from different plants growing in area of Solapur city and surrounding villages. Plant leaves were further dried in an oven at 40°C for a total of three days and then finally ground to a fine powder by grinder. Extraction is carried out by using petroleum ether and ethyl alcohol.<sup>(12-16)</sup>
- B) **Preparation of leaf extract :** 20 gram of finely grinded powder of *Moringa* leaves were soaked in two different conical flasks containing petroleum ether and ethyl alcohol respectively. The sample were shaken on rotary shaker at 200 rpm for 24 hrs. The extract were filtered using Whatman filter paper. Then solvent were evaporated to obtain dry extract using a rotary evaporator and were stored in refrigerator for antifungal analysis. Before testing five different concentration of extract were prepared for both solvents separately. This was carried by dissolving 100 mg, 200 mg, 300 mg, 400mg and 500mg of extract in 1 ml solvent respectively.<sup>(12-16)</sup>
- C) **Antifungal acivity :** The antifungal activity of the *Moringa oleifera* leaf extracts was determined using agar well diffusion method by following the known procedure. Fungal strains were spread on the surface of agar plate aseptically by sterile cotton swab. When surface was little dried wells of 4 mm diameter were punched with the help of sterile stainless steel boarer. 20 µl of petroleum and ethanol extracts of *Moringa* leaves were loaded separately in wells in separate agar plates. Separate wells were used for different concentrations. Each plate further contained on well loaded with solvent act as negative control. While another well was loaded with 20 µl antibiotic Nystatin (100 mg/ml). The plates were incubated at 28°C for 72 hours and the antifungal activity was assessed by measuring the diameter of the zone of inhibition at the interval of 24 hrs.<sup>(12-16)</sup>


## III. RESULT AND DISCUSSION

From our study, it is found that as concentration of Moringa leaves extract increases the diameter of zone of inhibition found increased. Against *Cryptococcus neoformans* zone of inhibition due to ethanolic extract of Moringa leaves was found increased from 6.3 mm to 11.8 mm as concentration increases from 100 to 600 mg per ml. While zone of inhibition due to petroleum extract of Moringa leaves was found increased from 4.1 mm to 8.1 mm as concentration

increases from 100 to 600 mg per ml.(Graph-I) On other hand against *Penicillium crysogenum* zone of inhibition due to ethanolic extract of Moringa leaves was found increased from 5.3 mm to 11.7 mm as concentration increases from 100 to 600 mg per ml. While zone of inhibition due to petroleum extract of Moringa leaves was found increased from 3.2 mm to 7.3 mm as concentration increases from 100 to 600 mg per ml.(Graph-II)



Graph.1: Effect of Ethanolic extract (Red line) and Petroleum ether extract (Violet line) of *Moringa oleifera* leaves against *Cryptococcus neoformans*.



Graph.2: Effect of Ethanolic extract (Red line) and Petroleum ether extract (Violet line) of *Moringa oleifera* leaves against *Penicillium crysogenum*.

Nystatin is a polyene antibiotic. Polyene antibiotic is a class of antimicrobial polyene compounds that target fungi. It is most commonly used for the treatment of fungal infections. It was used as a positive control against both *Cryptococcus neoformans* and *Penicillium crysogenum*.

The results were higher and effective for ethanolic extract than petroleum extract. The result were also found more effective against *Cryptococcus neoformans* (Graph-I) than *Penicillium crysogenum*.(Graph-II)

#### IV. CONCLUSION

From our study it comes to known that there are some alkaloids present in extract of leaves of *Moringa oleifera* which can be effectively used to treat infection caused by *Cryptococcus neoformans* and *Penicillium crysogenum*.

#### REFERENCES

- [1] Nadkarni A. Indian Materia Medica. Popular Prakashan: Bombay, 1976, 810–6.
- [2] Anwar F and Bhangar M. Analytical characterization of *Moringa oleifera* seed oil grown in temperate regions of Pakistan. J. Agric. Food Chem. 2003;51:6558-63.
- [3] Anwar F et.al. *Moringa oleifera*: a food plant with multiple medicinal uses. *Phytother. Res.* 2007;21:17-25.
- [4] Morton JF. The horseradish tree, *Moringa pterygosperma* (Moringaceae)- A boon to arid lands? *Econ. Bot.* 1991, 45:318-333.
- [5] Gupta M. et.al. CNS activities of methanolic extract of *Moringa oleifera* root in mice. *Fitoterapia* 1999(3); 70: 244-50.
- [6] Shukla S et.al. Antifertility profile of the aqueous extract of *Moringa oleifera* roots. *Journal of Ethnopharmacology* 1998; 22(1): 51-62.
- [7] Lurling M and Beekman W. Anti-cyanobacterial activity of *Moringa oleifera* seeds. *Journal Appl Phycol* 2010;22(4):503-10.
- [8] Fahey J et.al. The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochemistry* 2001; 56: 5–51.
- [9] Bennett R et al. Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the multi-purpose trees *Moringa oleifera* L. (Horseradish tree) and *Moringa stenopetala* L. *J Agric Food Chem* 2003;51: 3546–53.
- [10] Pal S and Saha B. Studies on the antiulcer activity of *M. oleifera* leaf extract on gastric ulcer models in rats. *Phytother. Res.* 1995; 9: 463 – 5.
- [11] Bukar A and Oyeyi T. Antimicrobial profile of *Moringa oleifera* Lam. Extracts against some food-borne microorganisms. *Bayero Journal of Pure and Applied Sciences*, 2010; 3(1): 43-8.
- [12] Napolean, P et.al. Isolation, analysis and identification of phytochemicals of antimicrobial activity of *Moringa oleifera* Lam. *Current Biotica*, 2009; 3(1): 33 – 7.
- [13] Nwosu M and Okafor J Preliminary studies on the antifungal activities of some medicinal plants against *Basidiobolus* and some other pathogenic fungi. *Mycoses* 1995; 38(5-6): 191-5.
- [14] Chuang P.et.al. In vitro antifungal activity of ethanolic extract of the leaves of *Moringa oleifera* against dermatophytes. *Bioresources Technology* 2007; 98: 232–6.
- [15] Srinivasan, D et.al. Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *Journal of Ethnopharmacology.* 2001; 94: 217 - 22.
- [16] Himal Paudel Chhetri et al. Phytochemical and antimicrobial evaluations of some medicinal plants of Nepal. *Kathmandu university journal of science, engineering and technology* september 2008; 49-54.