

# Global structure of positive solutions for superlinear first-order periodic boundary value problems

Jiao Wang

**Abstract**—In this paper, we consider the nonlinear eigenvalue problems

$$\begin{cases} u'(t) + a(t)u(t) = \lambda h(t)f(t, u(t)), & t \in (0,1), \\ u(0) = u(1), \end{cases}$$

where  $\lambda > 0$ ,  $a \in C((0,1), [0, \infty))$ ,  $h \in C((0,1), [0, \infty))$ , and there exist  $t_0 \in (0,1)$ , such that  $h(t_0) > 0$ .  $f \in C([0, \infty), [0, \infty))$ ,  $f(0) = 0$ ,  $f(s) > 0$ ,  $s > 0$ .

**Index Terms**—Periodic boundary value problem; Eigenvalue Positive solutions; Existence MSC(2010):—39A10, 39A12

## I. INTRODUCTION

The first-order periodic boundary value problems play a very important role in many aspects. such as economy, finance, insurance, population structure and so on. Therefore, the existence of positive solutions is widely concerned by many scholars at home and abroad, and many rich and profound results have been obtained<sup>[1-10]</sup>. For example, the behavior of animal blood red blood cells, the survival competition between the two populations and the frequency of the circuit signals can be depicted by the first order periodic boundary value problem

$$\begin{cases} u'(t) = -a(t)g(u(t)) + \lambda b(t)f(u(t - \tau(t))), & t \in R, \\ u(0) = u(t + \omega), \end{cases} \quad (1.1)$$

with parameters.

In particular, Peng<sup>[1]</sup> uses fixed point theorems on cone to study the following questions

$$\begin{cases} u'(t) + f(t, u(t)) = 0, & t \in (0, T), \\ u(0) = u(T), \end{cases} \quad (1.2)$$

Where as  $T > 0$ , nonlinear term  $f \in C((0, T) \times R, R)$ ,

The main results are as follows:

**Theorem A.** Assume that there exists a positive number  $M > 0$ , such that  $Mx - f(t, x) \geq 0$  for  $x \geq 0$ ,  $t \in J$ . If

$$(A1) \liminf_{u \rightarrow 0^+} \min_{t \in (0, T)} \frac{f(t, u)}{u} > 0, \quad \limsup_{u \rightarrow +\infty} \max_{t \in (0, T)} \frac{f(t, u)}{u} < 0$$

$$(A2) \liminf_{u \rightarrow +\infty} \min_{t \in (0, T)} \frac{f(t, u)}{u} > 0, \quad \limsup_{u \rightarrow 0^+} \max_{t \in (0, T)} \frac{f(t, u)}{u} < 0$$

then, PBVP (1.2) has at least one positive solution.

Tisdell<sup>[2]</sup> uses Leray-Schauder degree and fixed point theory to discuss the first order periodic differential system

$$\begin{cases} u'(t) + a(t)u(t) = f(t, u(t)), & t \in (0,1), \\ u(0) = u(1), \end{cases} \quad (1.3)$$

Thus, we obtain some sufficient conditions for the existence of positive solutions of (1.3).

Inspired by the above literatures, we use the Dancer's bifurcation to study the global structure of positive solutions for following periodic boundary value problems

$$\begin{cases} u'(t) + a(t)u(t) = \lambda h(t)f(t, u(t)), & t \in (0,1), \\ u(0) = u(1), \end{cases} \quad (1.4)$$

We make the following assumptions:

(H1)  $h \in C((0,1), [0, \infty))$  is continuous, and there exist

$t_0 \in (0,1)$ , such that  $h(t_0) > 0$ ;

(H2)  $f \in C([0, \infty), [0, \infty))$ ,  $f(0) = 0$ ,  $f(s) > 0$ ,  $s > 0$ ;

(H3)  $a \in C((0,1), [0, +\infty))$ ;

(H4)  $f_0 = \infty$ , where  $f_0 = \lim_{s \rightarrow 0^+} \frac{f(s)}{s}$ ;

(H5)  $f_\infty \in [0, \infty]$ , where  $f_\infty = \lim_{s \rightarrow \infty} \frac{f(s)}{s}$

The main results of the present paper are as follows:

**Theorem 1.1.** Let (H1) - (H5) hold. Let

$E = \{u \in C[0,1] \mid u(0) = u(1)\}$ . Let  $\Sigma$  be the closure of the set of positive solutions for (1.4) in  $E$ .

(a) If  $f_\infty = 0$ , then there exists a sub-continuum  $\zeta$  of  $\Sigma$  with  $(0,0) \in \zeta$  and

$$\text{Pr } o_{j_R} \zeta = [0, \infty).$$

(b) If  $f_\infty \in (0, \infty)$ , then there exists a sub-continuum  $\zeta$  of  $\Sigma$  with  $(0,0) \in \zeta$  and

$$\text{Pr } o_{j_R} \zeta \supseteq [0, \frac{\lambda_1}{f_\infty}).$$

(c) If  $f_\infty = \infty$ , then there exists a sub-continuum  $\zeta$  of  $\Sigma$  with  $(0,0) \in \zeta$ ,  $\text{Pr } o_{j_R} \zeta$  is a bounded closed interval, and  $\zeta$  approaches  $(0, \infty)$  as  $\|u\| \rightarrow \infty$ .

**Theorem 1.2.** Let (H1) - (H5) hold.

(d) If  $f_\infty = 0$ , then (1.4) has at least one positive solution for  $\lambda \in (0, \infty)$ .

(e) If  $f_\infty \in (0, \infty)$ , then (1.4) has at least one positive

Jiao Wang, Department of Mathematics, Northwest Normal University, Lanzhou, Gansu, China, Mobile No18893703613.

solution for  $\lambda \in (0, \frac{\lambda_1}{f_\infty})$ .

(f) If  $f_\infty = \infty$ , then (1.4) has at least one positive solution for  $\lambda \in (0, \lambda_*)$ .

II. SUPERIOR LIMIT AND COMPONENT

$Y = C[0,1]$  is a Banach space,  $K = \{u \in Y \mid u(t) \geq 0, t \in [0,1]\}$ .

The norm in  $C[0,1]$  is defined as follows

$$\|u\|_0 = \max_{t \in [0,1]} |u(t)|.$$

Define an operator  $T : K \rightarrow Y$  by,

$$T u(t) = \int_0^1 H(t,s)h(s)f(u(s))ds, \quad t \in [0,1].$$

Where

$$H(t,s) = \begin{cases} \frac{e^{\int_s^t a(\theta)d\theta}}{e^{\int_0^t a(\theta)d\theta} - 1}, & 0 \leq t \leq s \leq 1, \\ \frac{e^{\int_s^t a(\theta)d\theta}}{1 - e^{-\int_0^t a(\theta)d\theta}}, & 0 \leq s \leq t \leq 1. \end{cases}$$

Denote  $\sigma = e^{-\int_0^1 a(\theta)d\theta}$ , then

$$\frac{\sigma}{\sigma - 1} \leq H(t,s) \leq \frac{1}{1 - \sigma}, \quad (t,s) \in (0,1) \times (0,1). \quad (2.1)$$

Denote the cone  $P$  in  $Y$  by

$$P = \{u \in Y \mid u(t) \geq \sigma \|u\|, t \in (0,1)\}.$$

Define an operator  $T_\lambda : P \rightarrow Y$  by,

$$T_\lambda u(t) = \lambda \int_0^1 H(t,s)h(s)f(u(s))ds, \quad t \in [0,1]. \quad (2.2)$$

**Definition 2.1.** Let  $Y$  be a Banach space and  $\{C_n \mid n = 1, 2, K\}$  be a family of subsets of  $Y$ . Then the superior limit  $D$  of  $\{C_n\}$  is defined by

$$D := \limsup_{n \rightarrow \infty} C_n = \{x \in Y \mid \exists \{n_i\} \subset N$$

and  $x_{n_i} \in C_{n_i}$ , such that  $x_{n_i} \rightarrow x\}$

**Definition 2.2.** A component of a set  $M$  is meant a maximal connected subset of  $M$ .

**Lemma 2.3.** Suppose that  $Y$  is a compact metric space,  $A$  and  $B$  are non-intersecting closed subsets of  $Y$ , and no component of intersects both  $A$  and  $B$ . Then there exist two disjoint compact subsets  $X_A$  and  $X_B$ , such that  $Y = X_A \cup X_B, A \subset X_A, B \subset X_B$ .

**Lemma 2.4.** Let  $Y$  be a Banach space, and let  $\{C_n\}$  be a family of connected subsets of  $Y$ , Assume that

(i) there exist  $z_n \in C_n, n = 1, 2, K$ , and  $z^* \in X$ , such that  $z_n \rightarrow z^*$ ;

(ii)  $\lim_{n \rightarrow \infty} r_n = \infty$ , where  $r_n = \sup\{\|x\| \mid x \in C_n\}$ ;

(iii) for every  $R > 0, (\bigcup_{n=1}^\infty C_n) \cap B_R$  is a relatively compact set of  $Y$ , where

$$B_R = \{x \in X \mid \|x\| \leq R\}.$$

Then there exists an unbounded component  $C$  in  $D$  and  $z^* \in C$ .

**Lemma 2.5.** Assume that (H1) hold. Then  $T_\lambda : P \rightarrow P$  is completely continuous.

**Lemma 2.6.** Let (H1) – (H2) hold. Let

$\Omega_r = \{u \in K \mid \|u\| < r, r > 0\}$ . If  $u \in \partial\Omega_r, r > 0$ , then

$$\|T_\lambda u\| \leq \lambda \hat{M}_r \int_0^1 G(s,s)h(s)ds. \quad (2.3)$$

Where  $\hat{M}_r = 1 + \max_{0 \leq s \leq r} \{f(s, u(s))\}$ .

**Proof:** since  $\forall t \in [0,1], f(u(t)) \leq \hat{M}_r$ , it follows that

$$\begin{aligned} \|T_\lambda u\| &\leq \lambda \int_0^1 G(s,s)h(s)f(s, u(s))ds \\ &\leq \lambda \hat{M}_r \int_0^1 G(s,s)h(s)ds. \end{aligned}$$

**Lemma 2.7.** Let (H1) hold, and let  $r(T)$  be the spectral radius of  $T$ . Then  $r(T) > 0$ , and  $r(T)$  is a simple eigenvalue with an eigenfunction  $\varphi \in \text{int } K_e$  and there is no other eigenvalue with a positive eigenfunction.

**Lemma 2.8.** Let (H1) hold, and let  $r(T)$  be the spectral radius of  $T$ . Then  $\lambda_1 = \frac{1}{r(T)}$  is a simple eigenvalue with an eigenfunction  $\varphi \in \text{int } K_e$  and there is the unique eigenvalue with an eigenfunction  $\varphi \in \text{int } K_e$  and there is no other eigenvalue with a positive eigenfunction.

III. EIGENVALUE WITH A POSITIVE EIGENFUNCTION

Denote  $e(t) = 1, t \in [0,1]$ , and let

$$Y_e = \bigcup_{\rho > 0} \rho[-1,1], |x|_e = \inf\{\rho \mid \rho > 0, x \in \rho[-1,1]\}.$$

Set

$$K_e = Y_e \cap K = \{x \in K \mid x \leq \rho e \text{ for some } \rho > 0\}.$$

Then

(a1)  $K_e$  is a normal cone of  $Y_e$  with nonempty interior;

(a2)  $(Y_e, |\cdot|_e)$  is a Banach space and continuously imbedding in  $(Y, \|\cdot\|)$ .

Notice also that an  $x \in Y_e$  is in  $\text{int } K_e$  the interior of  $K_e$  in  $Y_e$  if and only if  $x \geq \rho e$  for some  $\rho > 0$ .

IV. PROOF OF THE MAIN RESULT

To prove the main result, we define  $f^{[n]}(s): [0, \infty) \rightarrow [0, \infty)$  by

$$f^{[n]}(s) = \begin{cases} f(s), & s > (\frac{1}{n}, \infty), \\ nf(\frac{1}{n})s, & s \in [0, \frac{1}{n}]. \end{cases}$$

Then  $f^{[n]}(s) \in C([0, \infty), [0, \infty))$  with  $f^{[n]}(s) > 0$

for all  $s \in (0, \infty)$  and  $(f^{[n]})_0 = nf(\frac{1}{n}) > 0$ .

By (H3), it follows that  $\lim_{n \rightarrow \infty} (f^{[n]})_0 = \infty$ .

and accordingly, (b) hold. (c) can be deduced directly from the Arzela-Ascoli Theorem and the definition of  $g^{[n]}$ . Therefore, the superior limit of  $\{C_+^{[n]}\}$ , i.e.  $D$ , contains an unbounded connected component  $C$  with  $(0,0) \in C$ .

(a)  $f_\infty = 0$ . In the case, we show that  $Proj_R C = [0, \infty)$ .

Assume on the contrary that  $\sup\{\lambda \mid (\lambda, y) \in C\} < \infty$ , then there exists a sequence  $\{(\mu_k, y_k)\} \subset C$  such that

$$\lim_{k \rightarrow \infty} \|y_k\| = \infty, \mu_k \leq C_0, \quad (4.1)$$

for some positive constant  $C_0$  depending not on  $k$ . From Lemma 2.5, we have that  $\lim_{k \rightarrow \infty} \|y_k\| = \infty$ . This together with the fact

$$\min_{\sigma \leq t \leq 1-\sigma} y_k(t) \geq \sigma \|y_k\|, \text{ for all } 0 < \sigma < \min\{t_0, 1-t_0\} \quad (4.2)$$

implies that

$$\lim_{k \rightarrow \infty} y_k(t) = \infty, \text{ uniformly for } t \in [\sigma, 1-\sigma]. \quad (4.3)$$

Since  $(\mu_k, y_k) \in C$ , we have that

$$\begin{cases} y'_k(t) + a(t)y_k(t) = \mu_k h(t)g(y_k(t)), & t \in (0,1), \\ y_k(0) = y_k(1), \end{cases} \quad (4.4)$$

Set  $v_k(t) = y_k(t)/\|y_k\|$ . Then  $\|v_k\| = 1$ .

Now, choosing a subsequence and relabelling if necessary, it follows that there exists  $(\mu_*, v_*) \in [0, C_0] \times E$  with

$$\|v_*\| = 1$$

such that

$$\lim_{k \rightarrow \infty} (\mu_k, v_k) = (\mu_*, v_*), \text{ in } R \times E \quad (4.5)$$

Moreover, using (4.3)–(4.4) and the assumption  $f_\infty = 0$ , it follows that

$$\begin{cases} v'_*(t) + a(t)v_*(t) = \mu_* h(t) \cdot 0, & t \in (0,1) \\ v_*(0) = v_*(1). \end{cases}$$

And subsequently,  $v_*(t) \equiv 0$  for  $t \in [0,1]$ . This contradicts (4.5). Therefore

$$\sup\{\lambda \mid (\lambda, y) \in C\} = \infty.$$

(b)  $f_\infty \in (0, \infty)$ . In this case, we show that  $Proj_R C \subseteq [0, \frac{\lambda_1}{f_\infty})$ .

Let us rewrite (1.4) to the form

$$\begin{cases} u'(t) + a(t)u(t) = \lambda h(t)g_\infty u + \lambda h(t)\xi(u(t)), & t \in (0,1) \\ u(0) = u(1). \end{cases}$$

where  $\xi(s) = g(s) - g_\infty s$ . Obviously  $\lim_{|s| \rightarrow \infty} \xi(s)/s = 0$ .

Now by the same method used to prove [6, Theorem 5.1], we may prove that  $C$  joins  $(0,0)$  with  $(\frac{\lambda_1}{f_\infty}, \infty)$ .

(c)  $f_\infty = \infty$ . In this case, we show that  $C$  joins  $(0,0)$  with  $(0, \infty)$ .

Let  $\{(\mu_k, y_k)\} \subset C$  be such that  $\|\mu_k\| + \|y_k\| \rightarrow \infty$  as  $k \rightarrow \infty$ . then

$$\begin{cases} y'_k(t) + a(t)y_k(t) = \mu_k h(t)g(y_k(t)), & t \in (0,1) \\ y_k(0) = y_k(1). \end{cases}$$

If  $\{\|y_k\|\}$  is bounded, say,  $\|y_k\| \leq M_1$  for some  $M_1$  depending not on  $k$ , then we may assume that

$$\lim_{k \rightarrow \infty} \mu_k = \infty. \quad (4.6)$$

Note that

$$\frac{g(y_k(t))}{y_k(t)} \geq \inf\left\{\frac{g(s)}{s} \mid 0 < s \leq M_1\right\} > 0.$$

By condition (H1), there exist some  $0 < \alpha < \beta < 1$  such that  $h(t) > 0$  for  $t \in [\alpha, \beta]$ . So there exists a constant  $M_2 > 0$ , such that

$$h(t) \frac{g(y_k)}{y_k(t)} > M_2 > 0, \quad t \in [\alpha, \beta]. \quad (4.7)$$

Combining (4.6) and (4.7) with the relation

$$y'_k(t) + a(t)y_k(t) = \mu_k h(t) \frac{g(y_k(t))}{y_k(t)} y_k(t), \quad t \in (0,1) \quad (4.8)$$

From [3, Theorem 6.1], we deduced that must change its sign on  $[\alpha, \beta]$  if  $k$  is large enough. This is a contradiction. Hence  $\{\|y_k\|\}$  is unbounded.

Now, taking  $\{(\mu_k, y_k)\} \subset C$  be such that

$$\|y_k\| \rightarrow \infty \text{ as } k \rightarrow \infty \quad (4.9)$$

We show that  $\lim_{k \rightarrow \infty} \mu_k = 0$ .

Suppose on the contrary that, choosing a subsequence and relabelling if necessary,  $\mu_k \geq b_0$  for some constant  $b_0 > 0$ .

Then we have from (4.9)  $\|y_k\| \rightarrow \infty$

To apply the nonlinear Krein-Rutman Theorem, we extend  $f$  to an odd function  $g: R \rightarrow R$  by

$$g(s) = \begin{cases} f(s) & \text{if } s \geq 0, \\ -f(-s) & \text{if } s < 0. \end{cases}$$

Similarly we may extend  $f^{[n]}$  to an odd function

$$g^{[n]} : R \rightarrow R \text{ for each } n \in N.$$

Now let us consider the auxiliary family of the equations

$$\begin{cases} u'(t) + a(t)u(t) = \lambda h(t)g^{[n]}u, & t \in (0,1), \\ u(0) = u(1). \end{cases}$$

Let  $\zeta \in C(R)$  be such that

$$g^{[n]}(u) = (g^{[n]})_0 u + \zeta^{[n]}(u) = nf\left(\frac{1}{n}\right)u + \zeta^{[n]}(u).$$

Note that

$$\lim_{|s| \rightarrow 0} \frac{\zeta^{[n]}(s)}{s} = 0.$$

Let us consider

$$Lu - \lambda h(t)(g^{[n]})_0 u = \lambda h(t)\zeta^{[n]}(u) \quad (4.10)$$

As a bifurcation problem from the trivial solution  $u \equiv 0$ .

Equation (4.1) can be converted to the equation

$$\begin{aligned} u(t) &= \int_0^1 H(t,s)[\lambda h(s)(g^{[n]})_0 u(s) + \lambda h(s)\zeta^{[n]}(u(s))]ds \\ &:= (\lambda L^{-1}[h(\cdot)(g^{[n]})_0 u(\cdot)](t) + \lambda L^{-1}[h(\cdot)\zeta^{[n]}(u(\cdot))](t)) \end{aligned}$$

Further we note that  $\|L^{-1}[h(\cdot)\zeta^{[n]}(u(\cdot))]\| = o(\|u\|)$  for  $u$  near 0 in  $E$ .

By Lemma 2.7 and the fact  $(g^{[n]})_0 > 0$ , the results of nonlinear Krein-Rutman Theorem can be stated as follows: there exists a continuum  $C_+^{[n]}$  of positive solutions of (4.1) joining to infinity in  $\lambda$ . Moreover,  $(\lambda_1 / (g^{[n]})_0, 0)$  is the only positive bifurcation point of (4.1) lying on trivial solutions line  $u \equiv 0$ .

**Proof of Theorem 1.1** Let us verify that  $\{C_+^{[n]}\}$  satisfies all of the conditions of Lemma 2.4. Since

$$\lim_{n \rightarrow \infty} \frac{\lambda_1}{(g^{[n]})_0} = \lim_{n \rightarrow \infty} \frac{\lambda_1}{nf\left(\frac{1}{n}\right)} = 0,$$

Condition (a) in Lemma 2.4 is satisfied with  $z^* = (0,0)$ . Obviously

$$r_n = \sup\{\|\lambda\| + \|y\| \mid (\lambda, y) \in C_+^{[n]}\} = \infty,$$

as  $k \rightarrow \infty$ . This together with (4.3) and condition

(H1) imply that there exist constants  $\alpha_1, \beta_1$  with

$$\sigma < \alpha_1 < \beta_1 < 1 - \sigma, \text{ such that}$$

$$h(t) > 0, \lim_{k \rightarrow \infty} \mu_k \frac{g(y_k(t))}{y_k(t)} = \infty, \text{ for all } t \in [\alpha_1, \beta_1]$$

for every fixed constant  $0 < \sigma < \min\{t_0, 1 - t_0\}$ . Thus, we have from (4.8) and [3, Theorem 6.1] that  $y_k$  must change its sign on  $[\alpha_1, \beta_1]$  if  $k$  is large enough. This is a contradiction.

Therefore  $\lim_{k \leftarrow \infty} \mu_k = 0$ .

**Proof of Theorem 1.2** (a) and (b) are immediate

consequences of Theorem 1.1(a) and (b), respectively.

To prove (c), we rewrite (1.4) to

$$u = \lambda \int_0^1 H(t,s)h(s)f(u(s))ds =: T_\lambda u(t).$$

By Lemma 2.6, for every  $r > 0$  and  $u \in \partial\Omega_r$ ,

$$\|T_\lambda u\| \leq \lambda \hat{M}_r \int_0^1 G(s,s)h(s)ds,$$

where  $\hat{M}_r = 1 + \max_{0 \leq s \leq r} \{f(s)\}$ .

Let  $\lambda_r > 0$  be such that

$$\lambda_r \hat{M}_r \left(1 + \frac{\sum_{i=1}^{m-2} \alpha_i}{1 - \sum_{i=1}^{m-2} \alpha_i \eta_i}\right) \int_0^1 G(s,s)h(s)ds = r.$$

Then for  $\lambda \in (0, \lambda_r)$  and  $u \in \partial\Omega_r$ ,  $\|T_\lambda u\| \leq \|u\|$ . This means that

$$\Sigma \cap \{(\lambda, u) \in (0, \infty) \times K \mid 0 < \lambda < \lambda_r, u \in K, \|u\| = r\} = \emptyset \quad (4.11)$$

By Lemma 2.5 and Theorem 1.1, it follows that there is also an unbounded component joining  $(0,0)$  and  $[\infty, \infty)$  in  $[0, \infty) \times Y$ . Thus, (4.10) implies that for  $\lambda \in (0, \lambda_r)$ , (1.4) has at least two positive solutions.

#### REFERENCES

- [1] S. G. Peng, Positive solutions for first order periodic boundary value problem[J]. *Applied Mathematics and Computation*, 2004, 158: 345 - 351.
- [2] C. T. Christopher, Existence of solutions to first-order periodic boundary value problems[J]. *Journal of Mathematical Analysis and Applications*, 2006, 323(2): 1325-1332.
- [3] Z. Deng, Introduction to BVPs and Sturmian Comparison Theory for Ordinary Differential Equations[J]. *Central Chain Normal University Press*, 1987.
- [4] Wan, D. Jiang, Existence of positive periodic solutions for functional differential equations[J]. *Kyushu Journal of Mathematics*, 2002, 56(1): 193-202.
- [5] Y. Li, Positive solutions of second-order boundary value problems with sign-changing nonlinear terms[J]. *Journal of Mathematical Analysis and Applications*, 1994, 120(3): 743-748.
- [6] R. Ma, D.O'Regan, Nodal solutions for second-order m-point boundary value problems with nonlinearities across several eigenvalues[J]. *Nonlinear Anal.* 2006, 64: 1562- 1577.
- [7] J. R. Graef, L. J. Kong, Existence of multiple periodic solutions for first-order functional differential equations[J]. *Mathematical and Computer Modelling*, 2011, 54(11-12): 2962-2968.
- [8] H. Y. Wang, Positive periodic solutions of functional differential equations[J]. *Journal of Differential Equations*, 2004, 202: 354-366.
- [9] G. T. Whyburn, Topological Analysis, Princeton Math. Ser. vol. 23, Princeton University Press, Princeton, N. J. 1958.
- [10] R. Ma. Nonlinear perturbation of linear differential equation[M], Beijing: Science and Technology Press, 1994.
- [11] D. Guo. Nonlinear functional analysis[M]. Shandong: shandong Science and Technology Press, 1985

**Jiao Wang**, Collage of Mathematics and Statistics, Northwest Normal University, Lanzhou, China, Mobile 86- 18893703613.