

Existence of extremal periodic solutions for first-order functional differential equations

Jiao Wang

Abstract— In this paper, we study the existence of periodic extremal solutions for the first-order functional differential equations

$$x'(t) = a(t)g(x(t))x(t) - f(t, x(t - \tau(t)))$$

Where $a \in C(R, [0, \infty))$ is ω - periodic function and $\tau(t) \in C(R, R)$ is a continuous ω - periodic function. $\int_t^{t+\omega} a(t)dt > 0$, $f \in C(R^2, [0, \infty))$ and $f(t, x(t)) > 0$ for $x > 0$, $g \in C(R, [0, \infty))$ is non-decreasing function. The proof of the main results are based on the upper and lower solutions and monotone iteration method.

Index Terms— Upper and lower solutions, Extremal periodic solutions, Monotone iterative method, Existence.
 MSC(2010):—39A10, 39A12

I. INTRODUCTION

Functional differential equations with delays appear in many ecological, economic, control and physiological models. An important question is whether these equations can support periodic solutions. Many authors have studied it extensively and obtained some results^[1-15].

In particular, in 2004, Wang^[1] applied the fixed point index theory to study when the condition (A1) – (A2) is satisfied, the existence result of the solution of functional differential equations

$$x'(t) = a(t)g(x(t))x(t) - \lambda b(t)f(x(t - \tau(t))). \quad (1.1)$$

Where

(A1) $a, b \in C(R, [0, \infty))$ is ω - periodic function,

$\int_t^{t+\omega} a(t)dt > 0$, $\int_t^{t+\omega} b(t)dt > 0$, $\tau(t)$ is a continuous ω - periodic function;

(A2) $f, g \in C([0, \infty), [0, \infty))$ is continuous, $\forall u > 0$, $f(u) > 0$, $0 < l < g(u) < L < \infty$, where l, L is a constant.

Denote $f_0 = \lim_{u \rightarrow 0^+} \frac{f(u)}{u}$, $f_\infty = \lim_{u \rightarrow \infty} \frac{f(u)}{u}$, $\sigma = e^{-\int_0^\omega a(t)dt}$,

$$M(r) = \max_{t \in [0, r]} f(t), \quad m(r) = \min_{t \in [\frac{\sigma^L(1-\sigma^L)}{1-\sigma^L}r, r]} f(t).$$

Let $i_0 = \{ \text{the number of elements that valued is zero}$

in $f_0, f_\infty \}$,

$i_\infty = \{ \text{the number of elements that valued is infinity in } f_0, f_\infty \}$.

The main results are as follows:

Theorem A Assume that (A1) – (A2) hold, then

(a) if $i_0 = 1$ or $i_0 = 2$, then for $\lambda > \frac{1 - \sigma^L}{m(1)\sigma^L \int_0^\omega b(s)ds} > 0$,

problem (1.1) has i_0 positive ω - periodic function.

(b) if $i_\infty = 1$ or $i_\infty = 2$, then for $0 < \lambda < \frac{1 - \sigma^L}{M(1)\sigma^L \int_0^\omega b(s)ds}$,

problem (1.1) has i_∞ positive ω - periodic function.

(c) if $i_0 = 0$ or $i_\infty = 0$, then for λ large or small enough, problem (1.1) has no positive ω - periodic function.

In 2010, Kang^[5] apply the upper and lower solutions obtained the existence of periodic extremal solutions for the first-order functional differential equations

$$y'(t) = -a(t)y(t) - f(t, y(t - \tau(t))). \quad (1.2)$$

where $a(t) > 0$, f is T periodic function about t , and is a non decreasing continuous function about y .

The main results are as follows:

Theorem B If problem (1.2) has a lower solution $v_0(t)$ and a upper solution $\omega_0(t)$, for $\forall t_0 \in [t, t + \omega]$, $v_0(t_0) \leq \omega_0(t_0)$, problem (1.2) has a minimal solution $v^*(t)$ and a maximal solution $\omega^*(t)$. and it solution $y(t)$ are all in the $M = \{y \in C : v^*(t) \leq y(t) \leq \omega^*(t)\}$.

Inspired by the above works, we will discuss the existence of periodic extremal solutions for a broader class of first-order functional differential equations

$$x'(t) = a(t)g(x(t))x(t) - f(t, x(t - \tau(t))) \quad (1.3)$$

We make the following assumptions:

(H1) $a \in C(R, [0, \infty))$ is ω - periodic function, $\int_0^\omega a(t)dt > 0$, and $\tau(t) \in C(R, R)$ is a continuous ω - periodic function;

(H2) $f \in C(R^2, [0, \infty))$ and f is ω - periodic function about t for $\forall t \in R$, $g \in C(R, [0, \infty))$ is non-decreasing function.

The main results of the present paper are as follows:

Theorem 1.1. Let (H1) - (H2) hold. If problem (1.3) has a lower solution α and a upper solution β , such that

$$\alpha(t) \leq \beta(t), \quad t \in R, \quad (1.4)$$

and f satisfies

$$f(t, v) - f(t, w) \leq L(v - w), \quad v \leq w \quad (1.5)$$

where $L > 0, v, w \in D = \{x \in X : \alpha \leq x \leq \beta\}$, then problem (1.3) has a minimal solution x_{\min} and a maximal solution x_{\max} .

II. PRELIMINARIES

Let $X := \{x \in C(R, R), x(t + \omega) = x(t), t \in R\}$, which is the Banach space with the norm

$$\|x\| = \max_{t \in [0, \omega]} |x(t)|.$$

Define 2.1 If $\alpha \in X \cap C^1(R, R)$ satisfies:

$$\alpha'(t) \geq a(t)g(\alpha(t))\alpha(t) - f(t, \alpha(t - \tau(t))),$$

then α is the lower solution of problem (1.3).

Define 2.2 If $\beta \in X \cap C^1(R, R)$ satisfies:

$$\beta'(t) \leq a(t)g(\beta(t))\beta(t) - b(t)f(t, \beta(t - \tau(t))),$$

then β is the upper solution of problem (1.3).

For $\forall x \in X, \alpha, \beta$ are the upper and lower solutions of the problem (1.3). define

$$p_{\alpha, \beta}(x(t)) = \max\{\alpha(t), \min\{u(t), \beta(t)\}\},$$

$$g_{\alpha, \beta} = g(p_{\alpha, \beta}(x(t))), f_{\alpha, \beta} = f(t, p_{\alpha, \beta}(x(t))).$$

then

$$\begin{cases} p_{\alpha, \beta}(x(t)) = \alpha(t), & x(t) \leq \alpha(t), \\ p_{\alpha, \beta}(x(t)) = \beta(t), & x(t) \geq \beta(t), \\ \alpha(t) \leq p_{\alpha, \beta}(x(t)) \leq \beta(t), & x(t) \in X. \end{cases} \quad (2.1)$$

where $g_{\alpha, \beta}, f_{\alpha, \beta} : X \rightarrow X$, and $g_{\alpha, \beta}, f_{\alpha, \beta}$ is bounded.

Consider auxiliary problem

$$\begin{cases} x'(t) = a(t)g_{\alpha, \beta}(x(t))x(t) - f_{\alpha, \beta}(t, x(t - \tau(t))), \\ x(t) = x(t + \omega). \end{cases} \quad (2.2)$$

define the operator $T_{\alpha, \beta} : X \rightarrow X$ as follow

$$T_{\alpha, \beta} x(t) = \int_t^{t+\omega} G_{\alpha, \beta}(t, s, x) f_{\alpha, \beta}(t, x(t - \tau(t))) dt.$$

where

$$G_{\alpha, \beta}(t, s, x) = \frac{e^{-\int_t^s a(\theta)g_{\alpha, \beta}(x(\theta))x(\theta)d\theta}}{1 - e^{-\int_0^\omega a(\theta)g_{\alpha, \beta}(x(\theta))x(\theta)d\theta}}.$$

The method of [10] can verify $T_{\alpha, \beta} : X \rightarrow X$ is continuous and the solution of the problem (1.3) is equivalent to the fixed point of $T_{\alpha, \beta}$ in X .

Lemma 1 $\forall x_1, x_2 \in C^1(R, R)$, if

$$\begin{aligned} & x_1'(r) - a(r)g(x_1(r))x_1(r) \\ & \leq x_2'(r) - a(r)g(x_2(r))x_2(r), \quad r \in [t, t + \omega], \end{aligned} \quad (2.3)$$

and $x_1(t + \omega) \geq x_2(t + \omega)$, then $\forall r \in [t, t + \omega], x_1(r) \geq x_2(r)$.

Proof. Suppose on the contrary that $\exists r_0 \in [t, t + \omega]$, s.t.

$$\min_{r \in [t, t + \omega]} (x_1 - x_2) = x_1(r_0) - x_2(r_0) < 0.$$

if $r_0 \in (t, t + \omega)$, then $x_1'(r_0) = x_2'(r_0)$, by (2.2) and g is non-decreasing function, we can know

$$\begin{aligned} 0 &= x_1'(r_0) - x_2'(r_0) \\ &\leq a(r_0)g(x_1(r_0))x_1(r_0) - a(r_0)g(x_2(r_0))x_2(r_0) < 0. \end{aligned}$$

which is contradiction.

if $r_0 = 0$, then $\exists r_1 \in (t, t + \omega]$, s.t. for $\forall r \in [0, r_1]$, $x_1(r) - x_2(r) \leq 0$, and $x_1'(r) - x_2'(r) > 0$, has

$$\begin{aligned} 0 &< x_1'(r) - x_2'(r) \\ &\leq a(r)g(x_1(r))x_1(r) - a(r)g(x_2(r))x_2(r) < 0. \end{aligned}$$

which is contradiction. thus for $\forall r \in [t, t + \omega]$, has $x_1(r) \geq x_2(r)$. The conclusion is proved.

III. PROOF OF THE MAIN RESULT

Proof of Theorem 1 Through the monotone iterative method we construct x_{\min} , let $x_0 = \alpha$, define the solution

$$\begin{cases} x_{n+1}'(t) - a(t)g_{\alpha, \beta}(x_{n+1}(t))x_{n+1}(t) = -f_{\alpha, \beta}(t, x_n(t - \tau(t))), \\ x_{n+1}(t) = x_{n+1}(t + \omega). \end{cases} \quad (3.1)$$

We first prove:

$$\alpha(r) \leq x_n(r) \leq \beta(r) \quad (n = 0, 1, 2, K), \quad r \in [t, t + \omega]. \quad (3.2)$$

It is known from the inductive method, as $n = 0$,

$$\begin{aligned} & x_1'(r) - a(r)g_{\alpha, \beta}(x_1(r))x_1(r) \\ &= -f_{\alpha, \beta}(r, x_0(r - \tau(r))) \\ &= -f_{\alpha, \beta}(r, \alpha(r - \tau(r))) \\ &\leq \alpha'(r) - a(r)g_{\alpha, \beta}(\alpha(r))\alpha(r). \end{aligned}$$

By Lemma 1, we can know that $x_1(r) \geq \alpha(r)$. suppose $\alpha(r) \leq x_n(r) \leq \beta(r)$, next to prove $\alpha(r) \leq x_{n+1}(r)$. By g is a non-decreasing function and (3.1) we can obtain that

$$\begin{aligned} & x_{n+1}'(r) - a(r)g_{\alpha, \beta}(x_{n+1}(r))x_{n+1}(r) \\ &= -f_{\alpha, \beta}(r, x_n(r - \tau(r))) \\ &\leq L(\alpha((r - \tau(r)) - x_n((r - \tau(r)))) - f_{\alpha, \beta}(r, \alpha(r - \tau(r)))) \\ &\leq -f_{\alpha, \beta}(r, \alpha(r - \tau(r))) \\ &\leq \alpha'(r) - a(r)g_{\alpha, \beta}(\alpha(r))\alpha(r). \end{aligned}$$

by lemma 1 we can know $\forall r \in [t, t + \omega], \alpha(r) \leq x_{n+1}(r)$.

Similarly $\forall r \in [t, t + \omega], x_{n+1}(r) \leq \beta(r)$, so (3.2) has been proved.

Next to prove

$$x_n(r) \leq x_{n+1}(r) \quad (n = 0, 1, 2, K), \quad r \in [t, t + \omega]. \quad (3.3)$$

By the inductive method, when $n = 0$, (3.3) is clearly established. Suppose $x_{n-1} \leq x_n$. by (3.1)(3.2) we can obtain that

$$\begin{aligned} & x_{n+1}'(r) - a(r)g_{\alpha, \beta}(x_{n+1}(r))x_{n+1}(r) = -f_{\alpha, \beta}(r, x_n(r - \tau(r))) \\ & \leq L(x_{n-1}((t - \tau(r)) - x_n((r - \tau(r)))) - f_{\alpha, \beta}(r, x_{n-1}(r - \tau(r)))) \\ & \leq -f_{\alpha, \beta}(r, x_{n-1}(r - \tau(r))) \\ & = x_n'(r) - a(r)g_{\alpha, \beta}(x_n(r))x_n(r). \end{aligned}$$

By lemma 1, for any $r \in [t, t + \omega]$, (3.3) is established. By (3.2) and (3.3) we can obtain that

$$\begin{aligned} \alpha(r) &= x_0(r) \leq x_1(r) \leq \dots \leq x_n(r) \\ &\leq x_{n+1}(r) \leq \dots \leq \beta(r), r \in [t, t + \omega]. \end{aligned}$$

Thus, as $r \in [t, t + \omega]$,

$$x_{\min}(r) := \lim_{n \rightarrow \infty} x_n(r) \quad (3.4)$$

is existence. Because $\{x_n\}_{n \geq 1}$ is bounded in D , by Arzela-Ascoli theorem we can get $x_{\min} \in D$, and by (3.4) and Dini theorem, $\{x_n\}$ can be uniformly converged to x_{\min} in D .

Consider the following questions

$$\begin{cases} x'_{n+1}(t) - a(t)g_{\alpha,\beta}(x_{n+1}(t))x_{n+1}(t) = -f_{\alpha,\beta}(t, x_{n+1}(t - \tau(t))), \\ x_{n+1}(t) = x_{n+1}(t + \omega). \end{cases} \quad (3.5)$$

Define an operator as follows

$$T_{n,\alpha,\beta}(x(t)) := \int_t^{t+\omega} G(t, s, x) f_{\alpha,\beta}(s, x_n(s - \tau(s))) ds.$$

Because $T_{n,\alpha,\beta}(x_{n+1}) = x_{n+1}$, continuity of operator T and

$$f_{\alpha,\beta}(t, x_{n+1}(t - \tau(t))) \rightarrow f_{\alpha,\beta}(t, x_{\min}(t - \tau(t)))$$

we can get $T_{n,\alpha,\beta}(x_{n+1}) \rightarrow T_{\alpha,\beta}(x_{\min})$ in D . That is $T_{x_{\min}} = x_{\min}$, So $x_{\min} \in D$ is the solution of the problem (1.3).

Finally, suppose that another $x \in D$ is the solution of the problem (1.3), then

$$x_n(r) \leq x(r) \quad (n = 0, 1, 2, \dots, K), r \in [t, t + \omega], \quad (3.6)$$

obviously when $n = 0$, (3.6) was set up, from the inductive method, we can assume that $x_{n+1}(r) \leq x(r)$, and then prove

$x_{n+1}(r) \leq x(r)$, by (3.1), we can obtained that

$$\begin{aligned} x'_{n+1}(r) - a(r)g_{\alpha,\beta}(x_{n+1}(r))x_{n+1}(r) &= -f_{\alpha,\beta}(r, x_n(r, \tau(r))) \\ &\leq L(x((r - \tau(r)) - x_n((r - \tau(r)))) - f_{\alpha,\beta}(r, x(r - \tau(r)))) \\ &= x'(r) - a(r)g_{\alpha,\beta}(x(r))x(r), \end{aligned}$$

by the lemma 1, (3.6) is set up.

So from (3.6), we can get $x_{\min} \leq x$, that is $x_{\min} \in D$ is the minimal solution to the problem (1.3).

Similarly, let $y_0 = \beta$, by the existence of descending sequence $\{y_n\}$, we can obtain $x_{\max} \in D$ is the maximal solution to the problem (1.3). Theorem 1 is proved.

ACKNOWLEDGMENT

The author is very grateful to the anonymous referees for their valuable suggestions. Our research was supported by the NSFC(11626016).

REFERENCES

- [1] H. Wang. Positive periodic solutions of functional differential equations [J]. *J. Differential Equations*. 2004, 202: 354-366.
- [2] A. Wan, D. Jiang. Existence of positive periodic solutions for

- functional differential equations[J]. *Kyushu J Math*, 2002, 56: 193-202.
- [3] C. Fabry, J. Mawhin, N. Nkashamam. A multiplicity result for periodic solutions of forced nonlinear second order ordinary differential equations[J]. *Bull. London Math. Soc.* 1986, 18: 173-180.
- [4] C. Bereanu, J. Mawhin. Existence and multiplicity results for periodic solutions of nonlinear difference equations[J]. *J. Difference Equ. Appl.* 2006, 12: 677-695.
- [5] S. Kang, B. Shi, G. Wang. Existence of maximal and minimal periodic solutions for first-order functional differential equations[J]. *Appl. Math. Lett.* 2010, 23: 22 - 25.
- [6] L. Kong, Q. Kong. Uniqueness and parameter dependence of solutions of second-order boundary value problems[J]. *Appl. Math. Lett.* 2009, 22: 1633 - 1638.
- [7] C. Bereanu, J. Mawhin. Existence and multiplicity results for some nonlinear problems with singular ϕ -Laplacian[J]. *J. Differential Equations*. 2007, 243: 536-557.
- [8] R. Ma, C. Gao, J. Xu. Existence of positive solutions for first order discrete periodic boundary value problem with delay[J]. *Nonlinear Anal.* 2011, 74: 4186-4191.
- [9] I. Rachhkova. Upper and Lower solutions and multiplicity results[J]. *J. Math. Anal. Appl.* 2000, 246: 446 -464.
- [10] J. Graef, L. Kong. Periodic solutions of first-order functional differential equations[J]. *Appl. Math. Lett.* 2011, 24: 1981-1985.
- [11] R. Ma, B. Yang, Z. Wang. Positive periodic solutions of first-order delay differential equations with impulses[J]. *Appl. Math. Comput.* 2013, 219: 6074-6083.
- [12] J. R. Graef, L. Kong. Existence of multiple periodic solutions for first order functional differential equations[J]. *Math. Comput. Model.* 2011, 54: 2962-2968.
- [13] F. Kong, S. Lu, Z. Luo. Positive periodic solutions for singular higher order delay differential equations[J]. *Results Math.* 2017, 1: 71-86.
- [14] N. Wang. Positive Solutions of Three-Order Delayed Periodic Boundary Value Problems[J]. *Discrete Dyn. Nat. Soc.* 2017, 6:34-18.
- [15] A. Ardjouni, A. Djoudi. Existence of positive periodic solutions for second-order nonlinear neutral difference equations with variable delay[J]. *Nonlinear Stud.* 2017, 2: 367-375.

Jiao Wang, Collage of Mathematics and Statistics, Northwest Normal University, Lanzhou, China, Mobile 86- 18893703613.