# Statistical Method for Analysis of Responses in Control Critical Trials with Three Outcomes

# Oyeka I. C. A. and Nwankwo Chike H.

Abstract— This paper proposes a statistical method for the analysis of multiple responses or outcome data in case control studies including situations in which the observations are either continuous or frequency data. Test statistics are proposed for assessing the statistical significance of differences between case-control response score. The proposed methods are illustrated with some sample data. When there only three possible response options in which the proposed method and the Stuart-Maxwell test can be equally used to analyse the data, the proposed test statistic is show to be at least as powerful as the Stuart-Maxwell test statistic.

*Index Terms*— Multiple Response, Case Control, Scores, Test Statistic, Treatment, Prospective, Retrospective

#### I. INTRODUCTION

Often in controlled comparative prospective or retrospective studies involving matched samples of subjects or patients, the response of a subject to a predisposing factor in a retrospective study or to a condition or treatment in a prospective study may be dichotomous with only two possible naturally exclusive outcomes and appropriate for analysis using the McNemar Test (Gibbons1973). But the responses may be much finer than simply dichotomous, assuming several possible values. For example in a retrospective study where the predisposing factor may be a subject's employment status, a subject may be classified as unemployed, self employed, public servant, student, housewife etc. In a prospective study involving some conditions or tests, subjects or patients may be classified as recovered, much improved, improved, no change, worse or dead. A treatment or drug may be graded as very effective, effective, ineffective etc

If there are only three possible response options or categories, then the Stuart-Maxwell test (Fleiss, 1981; Robertson et al, 1974; Schlesselman, 1992; Zhao and Kolonel, 1992; Box and Cox, 1964; Maxwell, 1970; Stuart, 1955; Fleiss, 1981; Everitt, 1977) may be used to analyse the data. We here propose an alternative and easier to use method that is often more powerful than the usual Stuart/Maxwell test for three outcomes in a clinical trial and which is easily generalisable when there are more than three outcomes.

### II. THE PROPOSED METHOD

Suppose we have a random sample of n pairs of patients or subjects matched on a number of characteristics to be exposed to two experimental conditions, treatments, drugs or tests. Suppose further that the responses of these pairs of subjects are more than dichotomous but numbering  $c(c \ge 3)$ 

Manuscript received January 30, 2015.

**Oyeka I. C. A. and Nwankwo Chike H.,** Department of Statistics, Nnamdi Azikiwe University, Awka Nigeria.

possible response options. Suppose further that the  $i_{th}$  pair of patients is selected, for  $i=1,2,\ldots,n$  and one member of the pair is randomly assigned to one of the treatments  $T_I$  (standard drug; control), say, and the remaining member of the pair is assigned to the second treatment  $T_2$  (new drug; case) say, and the various c possible responses are recorded for each subject. If in particular the responses of each matched pair of subjects are classified into c=3 mutually exclusive categories or classes, the data presentation format is as in Table 1 below.

**Table 1:** Format For Presentation Of Data On 'C' = 3 Outcomes in a Clinical Trial of Matched Pairs

Outcomes in a Clinical Trial of Matched Pairs						
	Outcome Category for Control (Standard $T_I$ )					
Outcome Category for Cases (Experimental Condition $T_2$ )	1	2	3	Total (n <sub>i</sub> .)		
1	n <sub>11</sub>	n <sub>12</sub>	n <sub>13</sub>	n <sub>1.</sub>		
2	n <sub>21</sub>	n <sub>22</sub>	n <sub>23</sub>	n <sub>2.</sub>		
3	n <sub>31</sub>	n <sub>32</sub>	n <sub>33</sub>	n <sub>3.</sub>		
Total (n. <sub>j</sub> )	n <sub>.1</sub>	n <sub>.2</sub>	n <sub>.3</sub>	n(=n)		

Each entry in Table 1 consists of a matched pair of case and control subjects. For example  $n_{ij}$  is the number of pairs in which the case is in category 'i' response while the corresponding control subject is in outcome or response category j for i, j = 1, 2, 3.  $n_i$ . and  $n_{ij}$  are respectively the total number of pairs in which the case is in category 'i' response and the control is in category 'j' response for i, j = 1, 2, 3.

In all, there are a total of 
$$n=n$$
.  $=\sum_{i=1}^3\sum_{j=1}^3n_{ij}=\sum_{i=1}^3n_{i.}=\sum_{j=1}^3n_{.j}$ 

pairs of subjects studied. A null hypothesis usually tested using the Stuart-Maxwell test is that case and control subjects or patients do not differ in their response to the treatments. The corresponding star-Maxwell test statistic for this purpose is

$$x^{2} = \frac{\bar{n}_{23}.d_{1}^{2} + \bar{n}_{13}.d_{2}^{2} + \bar{n}_{12}.d_{3}^{2}}{2(\bar{n}_{12}.\bar{n}_{13} + \bar{n}_{12}.\bar{n}_{23} + \bar{n}_{13}.\bar{n}_{23})} \dots (1)$$

which under  $H_o$  has approximately the chi-square distribution with 2 degrees of freedom, for sufficiently large n = n. where

$$d_i = n_{i,-}n_{i}$$
 ... (2)  
and  
 $\bar{n}_{ij} = \frac{n_{ij} + n_{ji}}{2}$  ... (3)  
for  $i, j = 1, 2, 3; i \neq j$ 

Now to develop the proposed method let, as in Stuart-Maxwell method, the difference between the number of pairs of respondents in the  $i_{th}$  category of responses for case and  $j_{th}$  category of responses for control (Miettinen, 1969; Maxwell, 1970; Everitt, 1977; Stuart, 1955) be  $d_i$  (equation 2) which is independent of  $n_{ii}$ , i = 1, 2, 3 the number of pairs in which both case and control subjects have the same response or outcome. Also let

$$d_{ij} = n_{ij} - n_{ji} \qquad \dots (4)$$


which is the difference between the number of pairs in which the case is in the response category i and the control is in the response category j and the number of pairs in which the case is in response category j and the control is in the response category i; i, j = 1, 2, 3  $i \neq j$ .

Now having selected our random sample of n matched pairs, let  $x_{ij}$  be the response by a member of the randomly selected  $i_{th}$  pair of patients or subjects randomly assigned treatment  $T_1$ (control, standard drug) and  $x_{i2}$  be the response by the other member of the pair of patients or subjects assigned treatment  $T_2$  (case, new drug) for  $i = 1, 2, \dots, n$ . We here assume for ease of presentation, but without loss of generality, that the three mutually exclusive possible response categories have been ordered from the highest or most serious (lowest or least serious) level of response to the lowest or least serious (highest or most serious) level of response. For example, a patient's response to a treatment for an illness or disease may range from recovered, through no change to dead; a subject's response to a screening test may range variously from definitely positive, no change, to definitely negative. A candidate's or student's performance in a job interview or examination may range from very poor, good, to excellent.

We here assume that those responses have been appropriately arranged either in increasing or decreasing order of seriousness.

1, If  $x_{i2}$ , i.e the response by the member in the  $i_{th}$ 

Now let



This means that  $u_i$  assumes the value 1, if the response of the member of the  $i_{th}$  pair of patients administered treatment  $T_2$  (case) is a higher or more serious (lower or less serious) level or response than the response of the other member of the pair administered treatment  $T_i$  (control); 0, if the response of the two members of the pair are the same, and -1, if treatment  $T_2$  (case) the response of the pair administered is a lower or less serious (higher or more serious) level of response than the response of the other member of the pair administered treatment  $T_i$  (control) for all the 3 response categories. Now let

$$\pi^{+} = P(u_{i} = 1); \ \pi^{0} = P(u_{i} = 0); \ \pi^{-} = P(u_{i} = -1)$$

Where

 $\pi^{+} + \pi^{0} + \pi^{-} = 1$ 

Let

 $W = \sum_{i=1}^{n} u_{i}$ 

Now

 $E(u_{i}) = \pi^{+} - \pi^{-}$ 

And

 $Var(u_{i}) = \pi^{+} + \pi^{-} - (\pi^{+} - \pi^{-})^{2}$ 

Also

 $E(W) = \sum_{i=1}^{n} Eu_{i} = n(\pi^{+} - \pi^{-})$ 

... 11

Note that  $\pi^+ - \pi^-$  is the differential response rate between the sub-populations administered treatments  $T_2$  (case) and  $T_1$  (control) respectively in the paired population of patients or subjects for all the c = 3 response categories and is estimated by

$$\hat{\pi}^+ - \hat{\pi}^- = \frac{w}{\pi} \qquad \dots 12$$

Note also that  $\pi^+$ ,  $\pi^0$  and  $\pi^-$  which are respectively the probabilities that a randomly selected case is at a higher (or more serious) level, the same or lower (or less serious) level of response than the corresponding control subject in the pair for all the three response categories are estimated using the frequencies in Table 1 and following the specification in Equation 5 as

$$\hat{\pi}^{+} = P^{+} = \frac{f^{+}}{n} = \frac{\sum_{i=1}^{2} \sum_{j=2}^{3} n_{ij}}{n} = \frac{n_{12} + n_{13} + n_{23}}{n} \qquad ... 13$$

$$\hat{\pi}^{0} = P^{0} = \frac{f^{0}}{n} = \frac{\sum_{i=1}^{3} n_{ii}}{n} = \frac{n_{11} + n_{22} + n_{33}}{n} \qquad ... 14$$

$$\hat{\pi}^{-} = P^{-} = \frac{f^{-}}{n} = \frac{\sum_{i=1}^{3} \sum_{j=1}^{2} n_{ij}}{n} = \frac{n_{21} + n_{31} + n_{32}}{n} \qquad ... 15$$

Where  $f^+$ ,  $f^0$  and  $f^-$  are respectively the number of 1's, 0's and -1's in the frequency distribution of the n values of these numbers in  $u_i$  in accordance with Equation 5,  $i = 1, 2, \dots, n$  Hence using these results in equation 12, we have that

$$W = f^{+} - f^{-} = \frac{\sum_{i=1}^{2} \sum_{j=2}^{8} n_{ij}}{i < j} - \frac{\sum_{i=2}^{8} \sum_{j=1}^{2} n_{ij}}{i > j} = (n_{12} - n_{21}) + (n_{13} - n_{31}) + (n_{23} - n_{32})$$

Now from equations 8 and 10 we have that

$$Var(W) = \sum_{i=1}^{n} Var(u_i) = n((\pi^+ + \pi^-) - (\pi^+ - \pi^-)^2)$$

Whose sample estimate is from equations 13 and 15 as

$$Var(W) = n((\hat{\pi}^+ + \hat{\pi}^-) - (\hat{\pi}^+ - \hat{\pi}^-)^2)$$
 ... 18

As noted above  $\pi^+$  is the proportion of pairs of case and control subjects in which on the average the response rate by the sub-population of patients or subjects administered treatment  $T_2$  (experimental, case) is greater (less) than the rate by the sub-population of patients or subjects administered treatment  $T_I$  (standard, control); while  $\pi^-$  is the proportion of pairs in which on the average the response rate by the sub-population of patients or subjects administered treatment  $T_1$  (standard, control) is greater (less) than the response rate by the sub-population of patients administered treatment  $T_2$ (experimental, case) in the paired population of patients or subjects for all the three response categories. Hence the null hypothesis that there exists no difference between the response rates by the sub-population of patients administered treatment  $T_2$  (experimental, case) and the sub-population of patients administered treatment  $T_I$  (standard, control) in the paired population of patients for all response categories is equivalent to the null hypothesis

$$H_0: \pi^+ - \pi^- = 0 \ versus \ H_1: \pi^+ - \pi^- \neq 0$$

To test this null hypothesis, we may use the test statistic

$$x^2 = \frac{w^2}{Var(w)} = \frac{w^2}{n(\hat{\pi}^+ + \hat{\pi}^- - (\hat{\pi}^+ + \hat{\pi}^-)^2)} = \frac{(f^+ - f^-)^2}{f^+ + f^- - \frac{(f^+ - f^-)^2}{n}}$$
... 20
Which under  $H_0$  has approximately a chi-square distribution with 1 degree of freedom for sufficiently large n. Although strictly speaking, the test statistic in Equation 20 has a Chi-square distribution with 1 degree of freedom, however because its construction in equation 5 involves a combination of some  $c = 3$  response categories, to help increase its power

because its construction in equation 5 involves a combination of some c=3 response categories, to help increase its power and reduce the chances of erroneously accepting a false null hypothesis (Type II error), it is here recommended that all comparisons should be made against critical Chi-square values with 3-1=2 degrees of freedom, instead of 1 degree of freedom. Hence here  $H_0$  is rejected at the  $\alpha$  level of

significance if 
$$x^2 \ge x^2_{(1-\alpha;2)}$$
 ... 21

Otherwise  $H_0$  is accepted.

$$n(\hat{\pi}^+ + \hat{\pi}^-) = \frac{\sum_{i=1}^2 \sum_{j=2}^3 n_{ij}}{i < i} + \sum_{i=2}^3 \sum_{j=1}^2 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{21} + n_{31} + n_{32})) \dots 2n_{i=1}^2 \sum_{j=1}^3 n_{ij} = n((n_{12} + n_{13} + n_{23}) + (n_{12} + n_{23} + n_{23} + n_{23}) + (n_{12} + n_{23} + n_{23} + n_{23} + n_{23}) + (n_{12} + n_{23} + n_{23} + n_{23} + n_{23}) + (n_{12} + n_{23} + n_{23} + n_{23} + n_{23}) + (n_{12} + n_{23} + n_{23} + n_{23} + n_{23}) + (n_{12} + n_{23} + n_{23} + n_{23} + n_{23} + n_{23}) + (n_{12} + n_{23} + n_{23$$

Hence using equation 16 and 22 in equation 20 the test statistic becomes  $\chi^2$ 

$$\frac{\left((n_{12}-n_{21})+(n_{13}-n_{31})+(n_{23}-n_{32})\right)^2}{n_{12}+n_{13}+n_{23}+n_{21}+n_{31}+n_{32}-\left(\frac{(n_{12}-n_{21})+(n_{13}-n_{31})+(n_{23}-n_{32})}{n}\right)^2}$$

... 23

If there are only two possible outcomes or responses, that is c = 2, equation 20 under  $H_0$  reduces to a modified version of the McNemar test statistic which is

$$\chi^2 = \frac{(n_{12} - n_{21})^2}{n_{12} + n_{21} - \frac{(n_{12} - n_{21})^2}{n}} \dots 24$$

This has a chi-square distribution with 1 degree of freedom.

Note that equation 24 has smaller variance than the usual McNemar test because of its modification to provide for possible ties between case and control subject pairs in their responses.

If c = 3, equation 20, under  $H_0$  reduces to

$$x^2 =$$

$$\frac{\left((n_{12}-n_{21})+(n_{13}-n_{31})+(n_{23}-n_{32})\right)^2}{n_{12}+n_{13}+n_{21}+n_{23}+n_{31}+n_{32}-\frac{\left((n_{12}-n_{21})+(n_{13}-n_{31})+(n_{23}-n_{32})\right)^2}{n}}$$

And this has a chi-square distribution with c - 1 = 3 - 1 = 2degrees of freedom.

Finally, note that if we let

$$m_{ij} = n_{ij} + n_{ji}, \ i,j = 1,2,3 \; ; \ i \neq j$$

Then the test statistic of equation 20 can be written in an easier and more compact form using equations 4 and 26 as

$$x^{2} = \frac{\left(\sum_{i=1}^{8} \sum_{j=1}^{8} d_{ij}\right)^{2}}{\sum_{i=1}^{8} \sum_{j=1}^{8} m_{ij} - \frac{\left(\sum_{i=1}^{8} \sum_{j=1}^{8} d_{ij}\right)^{2}}{n}}$$

If equation 20 leads to a rejection of the null hypothesis of equal response rates then one may wish to proceed to identify the response categories or combination of categories that may have led to the rejection of H<sub>0</sub>. This is done by appropriately pooling or combining the response options into (2) groups and apply the McNemar test (McNemar 1983, Somes 1983, Sheskin 2000) to each of the groups. In all cases comparisons are made using critical chi-square values with 2 degrees of freedom to again avoid erroneous conclusions.

# A. Illustrative Example 1

We here use data on matched pairs of 151 patients from a controlled comparative clinical trial who manifest three possible responses to illustrate the proposed method. Suppose the data in Table 2 are obtained by assigning a standard treatment  $T_I$  (control) and a new treatment  $T_I$  (case) at random to members of each pair of a random sample of 151 pairs of HIV patients matched on age, gender and body weight used in a controlled clinical trial to compare the effectiveness of two HIV drugs.

Table 2: Data from Controlled Comparative Clinical Trial Using Matched Pairs with Three Responses

	Standard Treatment $T_I$ (control)				
New reatment	Improved	No	Dead	Total (n <sub>i</sub> .)	
$T_2$ (case)		Change			
Improved	60	31	4	95 (=n <sub>1.</sub> )	
No Change	16	24	6	46 (=n <sub>2.</sub> )	
Dead	3	4	3	10 (=n <sub>3.</sub> )	
Total (n <sub>.j</sub> )	79	59	13	151 (=n <sub></sub> )	
	(n1)	(n. <sub>2</sub> )	(n3)		

To test the null hypothesis that case and control do not differ in their response to the treatments (Equation 19,) we have from equation 13 that

$$\hat{\pi}^+ = \frac{31+4+6}{151} = \frac{41}{151} = 0.271$$

And from Equation 15, we have that 
$$\hat{\pi}^- = \frac{16+3+4}{151} = \frac{23}{151} = 0.152$$
Note that  $\hat{\pi}^0 = 1 - (0.271 + 0.152) = 0.577$ 
Also from Equation 11, we have that 
$$E(W) = n(\hat{\pi}^+ - \hat{\pi}^-) = 151(0.271 + 0.152) = 151(0.119) = 17.969$$

From Equation 17, we have that Var(W) = 151[(0.271 + 0.152) -

$$(0.271 - 0.152)^2$$
] = 61.735

Hence from Equation 23, we have that

$$x^{2} = \frac{\left((31-16)+(4-3)+(6-4)\right)^{2}}{31+4+6+16+3+4-\left(\frac{(81-16)+(4-8)+(6-4)}{151}\right)^{2}} = \frac{324}{63.986} = 5.064 \left(p-value = 0.008\right)$$

Which with c - 1 = 3 - 1 = 2 degrees of freedom is highly statistically significant at  $\alpha = 0.01$ .

We may therefore conclude at the 1 percent significance level that the treatments have differential effects on the patients.

If we had used the Stuart/Maxwell method to analyse the data have from Equation  $d_1 = 95 - 79 = 16, d_2 = 46 - 59 = -13, d_3 =$ 10 - 13 = -3

Also letting 
$$\bar{n}_{ij} = \frac{n_{ij} + n_{ji}}{2}$$
,  $i = 1,2,3; j = 1,2,3; i \neq j$ , we have  $\bar{n}_{12} = \frac{31 + 16}{2} = \frac{47}{2} = 23.5; \ \bar{n}_{13} = \frac{4 + 3}{2} = \frac{7}{2} = 3.5; \ \bar{n}_{23} = \frac{6 + 4}{2} = \frac{10}{2} = 5$ 

Hence using the Stuart Maxwell test, we have  $x^2 = \frac{5(16)^2 + 3.5(-13)^2 + 23.5(-3)^2}{2((23.5)(3.5) + (23.5)(5) + (3.5)(5))} = \frac{1894}{434.5} =$ 4.359 (p - value = 0.12)

Which, with 2 degrees of freedom, is statistically significant at the 2 percent level of significance but not statistically significant at the 1 percent level of significance, the usually used norm in medical research?

Thus the present (extended) method leads to a rejection of the null hypothesis H<sub>0</sub> while the Stuart/Maxwell test statistic

leads to an acceptance of the null hypothesis at the 1 percent significance level. Hence the Stuart/Maxwell Test is likely to lead to an acceptance of a false null hypothesis (Type II error). This means that the present test statistic is likely to be more efficient and more powerful than the Stuart/Maxwell test statistic.

As noted above, the present method may also be used to analyse quantitative or numeric data obtained in matched controlled studies. Often, responses from controlled experiments are reported as numeric scores assuming all possible values on the real line such that scores in the interval  $(c_1, c_2)$  where  $c_1$  and  $c_2$  are real. For example, these responses may be values on the real line such that scores in the interval  $(c_1, c_2)$  where  $c_1$  and  $c_2$  are real numbers  $(c_1 < c_2)$ , indicate that the responses by the subject concerned are normal, negative, condition absent, etc; values less than c<sub>1</sub> indicate that the subjects have abnormally low scores; and values above c2 indicate that the subjects have abnormally high scores. It is also possible to have situations in which subjects have scores that are either some  $c_3$  units below  $c_1$  or some  $c_4$  units above c<sub>2</sub>. These subjects may be concerned to have non specific or non definitive manifestations. Subjects whose scores are below c<sub>3</sub> and above c<sub>4</sub> may be considered to have critically abnormal manifestations, one below the critical minimum and the other above the critical maximum normal scores. If these results are considered important manifestations, then the first set of subjects may be grouped into three response categories, while the second set of subjects may be grouped into five response categories for policy and management purposes.

To illustrate the use of the present method when the case and control subjects in matched controlled studies have quantitative scores with three possible outcomes for instance, we would proceed as follows:

Suppose as above, a random sample of n pairs of case and control subjects are used in a controlled experiment on two procedures  $T_1$  (control, standard) and  $T_2$  (case, experimental procedure). Suppose as before, one member of each pair is randomly assigned treatment  $T_1$  (control, standard) and the remaining member assigned treatment  $T_2$  (case, experimental procedure).

Let  $y_{il}$  and  $y_{i2}$  be respectively the responses or scores with real values, quantitatively measured, by the subjects assigned treatment  $T_1$  (control) and  $T_2$  (case) for the  $i_{th}$  pair of subject for i = 1, 2, ..., n.

Then  $u_i$  of Equation 3 may now be defined as

$$\begin{aligned} u_i &= \\ & \begin{cases} 1, & \text{if either } y_{i2} < c_1 \text{ and } c_1 \leq y_{i1} \leq c_2 \text{ or } y_{i2} < c_1 \\ & \text{and } y_{i1} > c_2 \text{ or } c_1 \leq y_{i2} \leq c_2 \text{ and } y_{i1} > c_2 \end{cases} \\ 0, & \text{if either } y_{i2} < c_1 \text{and } y_{i1} < c_1 \text{ or } c_1 \leq y_{i2} \leq c_2 \\ & \text{and } c_1 \leq y_{i1} \leq c_2 \text{ or } y_{i2} > c_2 \text{ and } y_{i1} > c_2 \end{cases} \\ -1, & \text{if either } c_1 \leq y_{i2} \leq c_2 \text{ and } y_{i1} < c_1 \text{ or } y_{i2} > c_2 \\ & \text{and } y_{i1} < c_1 \text{ or } y_{i2} > c_2 \text{ and } c_1 \leq y_{i1} \leq c_2 \end{cases} \\ -1, & \text{if either } c_1 \leq y_{i2} \leq c_2 \text{ and } c_1 \leq y_{i1} \leq c_2 \end{cases} \\ -1, & \text{if either } c_1 \leq y_{i2} \leq c_2 \text{ and } c_1 \leq c_2 \end{cases} \\ -1, & \text{if either } c_1 \leq c_1 \leq c_2 \end{cases} \\ -1, & \text{if either } c_1 \leq c_2 \leq c_2 \text{ and } c_1 \leq c_2 \leq c_2 \end{cases}$$

Note that this specification may be depicted in a 3 x 3 table if we let  $n_{ij}$  be the number of paired case and control subjects

in the (i, j)th case – control response classification for i = 1,2,3 and j = 1,2,3.

Specifications similar to Equation 28 can also be easily developed for more than three quantitative response categories if of interest. Now to use Equation 20 to analyse these data, we would again simply define  $\pi^+$ ,  $\pi^0$ ,  $\pi^-$  and W as in Equations 6-8. Then data analysis proceeds as usual.

#### B. Illustrative Example 2

A medical researcher is interested in knowing the relationship between heart disease and low density Lipo-Protein Levels (LPL). Using a random sample of 36 non-heart disease patients and another random sample of 36 heart disease patients, she paired each non heart disease patient with a heart disease patient matched in age, gender, body weight and occupation and then measured the LPL of each subject in the pair. The results are presented in Table 3

**Table 3:** LPL levels of Paired Samples of Patients in a Clinical Trial

S/N	Paired LPL levels	Scores $(u_i)$
1	(1.97,4.14)	0
2	(3.70,1.57)	-1
3	(5.40,5.60)	0
4	(2.60,5.10)	1
5	(3.10,1.50)	-1
6	(1.48,4.56)	1
7	(1.69,1.70)	0
8	(4.97,1.21)	-1
9	(2.34,2.51)	0
10	(3.95,1.55)	-1
11	(4.84,1.25)	-1
12	(4.65,4.59)	0
13	(1.29, 1.37)	0
14	(1.15,6.24)	1
15	(5.41,1.20)	-1
16	(4.62,1.25)	-1
17	(2.02,1.53)	-1
18	(1.45, 1.30)	0
19	(5.31,1.07)	-1
20	(5.18,4.37)	-1
21	(4.52,5.38)	1
22	(5.03,3.34)	-1
23	(5.21,4.55)	0
24	(4.74,5.59)	0
25	(3.76, 3.96)	0
26	(5.21,3.50)	-1
2 27	(5.09,4.66)	0
28	(1.97, 4.14)	0
29	(2.60,5.10)	1
30	(1.69,1.70)	0
31	(3.95,1.55)	-1
32	(1.29,1.37)	0
33	(4.62,1.25)	-1
34	(5.31,1.07)	-1
35	(5.03,3.34)	-1
36	(3.76,3.96)	0

LPL Normal range (1.68, 4.53)

Applying the specification of Equation 28 to the LPL levels of Table 3 with  $c_1 = 1.68$ , the lowest and 4.53 the highest normal values respectively we obtain the corresponding scores  $u_{i of} 1_s$ ,  $0_s$  and  $-1_s$  shown in the 3rd column of this table.

Thus we have  $f^+=5$ ,  $f^0=15$  and  $f^-=16$ . Hence, we have from Equations 12 - 15 that

$$\hat{\pi}^+ = \frac{5}{36} = 0.139$$
;  $\hat{\pi}^0 = \frac{15}{36} = 0.417$  and  $\hat{\pi}^- = \frac{16}{36} = 0.444$ 

From Equation 18, we have that the estimated variance of Wis

$$Var(W) = 36(0.139 + 0.444) - \frac{(-11)^2}{36} = 20.988 - 3.361 = 17.627$$

The null hypothesis to be tested is that heart disease patients and non-heart disease patients do not differ in their LPL which is equivalent to testing

$$H_0: \hat{\pi}^+ - \hat{\pi}^- = 0$$
 versus  $H_1: \hat{\pi}^+ - \hat{\pi}^- \neq 0$   
Using the test statistic of equation 20 or 23, we have that

$$x^2 = \frac{(-11)^2}{17.627} = \frac{121}{17.627} = 6.864 \ (p \ value = 0.0391)$$

which with c - 1 = 3 - 1 = 2 degrees of freedom is statistically significant at the 5 percent level  $(x_{0.95,2}^2 = 5.991)$ 

We may therefore conclude that heart disease patients and non- heart disease patients do infact differ in their LPL.

The data of Table 2 may infact be represented by a 3 x 3 table and following the specifications of Equation 28 with  $c_1 = 1.68$ 

and  $c_2 = 4.53$  to aid in clearer analysis as in Table 3.

**Table 4:** Distribution of Scores  $u_i$  of Matched pairs of case and control subjects of Table 3

	Control (T <sub>1</sub> ) Scores				
Case(T <sub>2</sub> ) Scores	Below Normal $(y_{il} < 1.68)$	Normal $(1.68 \le y_{il} \le .53)$	Above Normal $(y_{il} > 4.53)$	Total	
Below Normal ( $y_{i2} < 1.68$ )	4	0	2	6	
Normal $(1.68 \le y_{i2} \le 4.53)$	5	6	3	14	
Above Normal $(y_{i2} > 4.53)$	7	4	5	16	
Total	16	10	10	36	

To re-analyse these data consistent with the generalized

method, we have from Equation 13 that 
$$\hat{\pi}^+ = \frac{0+2+3}{36} = \frac{5}{36} = 0.139$$

$$\hat{\pi}^0 = \frac{4+6+5}{36} = \frac{15}{36} = 0.417$$

From Equation 14, we have that
$$\hat{\pi}^0 = \frac{4+6+5}{36} = \frac{15}{36} = 0.417$$
And from Equation 15, we have that
$$\hat{\pi}^- = \frac{5+7+4}{36} = \frac{16}{36} = 0.444$$

These are the same results obtained earlier using the scores in Table 3. We would therefore obtain the same values of W(-11) and chi-square (6.864) and arrive at the same conclusions. Hence, the present example illustrates how to analyse matched quantitative test scores without first converting them into frequency data.

The data of Example 2 as presented in Table 4 may also be analysed using the Stuart-Maxwell test. However as already pointed out, the Stuart/Maxwell test statistic is almost as powerful as the test statistic used in the proposed method presented here when the two methods are used with data of equal sample sizes

# III. SUMMARY AND CONCLUSION

We have in this paper presented and discussed a generalisable statistical method for the analysis of three responses or outcomes in case - control studies, including situations in which the data being analysed are either quantitative or qualitative frequency data.

Test statistics are developed for testing the statistical significance of differences between responses.

The proposed methods are illustrated with sample data and shown to be more powerful than the usual Stuart/Maxwell test when the two methods are equally applicable to a set of data.

#### REFERENCES

- [1] Box G. E. P. and Cox D. R: "An analysis of transformations. Journal of the Royal Statistical Society (B), 1964, 26, 211 - 252.
- Everitt H. B. S: The Analysis of Contingency Tables. London: Chapman and Hall, 1977.
- Fleiss, J. L. (1981): Statistical Methods for Rates and Proportions (Second Edition), New York. Wiley 1981.
- Gibbons, J. D. (1971): Nonparametric Statistical Inference. McGraw-Hill Book Company, New York.
- Maxwell, A. E: Comparing the classification of subjects by two independent judges. British Journal of Psychiatry, 1970, 116, 651 -
- McNemar Q: Note on the sampling error of the difference between [6] correlated proportions or percentages. Psychometrika 1947, 12, 153 -
- Miettinen O. S: Individual Matching with multiple controls in the case of all - or - more response. Biometrics, 1969, 22, 339 - 355.
- Robertson C, Boyle P, Hseih C, Macfarlane, G. J, and Maisonnesuve P: Some Statistical considerations in the analysis of case-control studies when the exposure variables are continuous measurements. Epidemiology, 1974, 5, 164-170.
- Schlesselman J. J. (1992): Case-Control studies. Oxford University Press, New York.
- [10] Sheskin D. J. (2000): Handbook of Parametric and Non-parametric Statistical procedures (Second Edition). Boca Raton Chapman and Hall, 491 - 508.
- [11] Somes, G: McNemar Test Encyclopaedia of Statistical Sciences, S. Kotz and N. Johnson eds, 1983, 5, 361 - 363. New York. Wiley.

# International Journal of Engineering and Applied Sciences (IJEAS) ISSN: 2394-3661, Volume-2, Issue-1, January 2015

- [12] Stuart, A. A: A test for homogeneity of the marginal distributions in a two way classification. Biometrika, 1955, 42, 412 416.
- [13] Zhao L. P and Kolonel L. N: "Efficiency Loss from Categorizing quantitative exposure into qualitative exposures in case-control studies". American Journal of Epidemiology, 136, 464 474. 1992.