

Solving Fuzzy Linear Systems

Dequan Shang, Jinglun Li, Xiaobin Guo

Abstract—In paper the fuzzy linear equation $A\tilde{x} = \tilde{b}$ in which A is a crisp matrix and \tilde{b} is an arbitrary fuzzy numbers vector, is investigated in detail. Numerical procedure for calculating the solution is designed and the sufficient condition for the existence of strong fuzzy solution is derived. Some examples are given to illustrate the proposed method.

Index Terms—Fuzzy numbers, Matrix analysis, Fuzzy linear system, Fuzzy approximate solution.

I. INTRODUCTION

System of simultaneous linear equations plays a major role in various areas such as mathematics, physics, statistics, engineering and social sciences. In many linear systems, some of the system parameters are vague or imprecise, and fuzzy mathematics is a better tool than crisp mathematics for modeling these problems, and hence how to design a better method for solving a fuzzy linear system is becoming more and more important thing. The concept of fuzzy numbers and their arithmetic operations were first introduced and investigated by Zadeh [29], Dubois et al.[15] and Nahmias [22]. A different approach to fuzzy numbers and the structure of fuzzy number spaces was given by Puri and Ralescu[26], Goetschell et al.[18] and Wu Congxin et al.[27, 29].

Since Friedman et al.[16] proposed a general model for solving an $n \times n$ fuzzy linear systems $A\tilde{x} = \tilde{b}$ by an embedding approach in 1998, lots of works have been done about some advanced fuzzy linear systems such as dual fuzzy linear systems (DFLS), general fuzzy linear systems (GFLS), fully fuzzy linear systems (FFLS) and general dual fuzzy linear systems (GDFLS) see [1-6, 9,13-14,30]. Some new theory and method for fuzzy linear systems still appeared [7, 10, 17, 19-20, 25-26] recently.

In this paper we propose a general model for solving the fuzzy linear system $A\tilde{x} = \tilde{b}$ by using a matrix approach. We extend the fuzzy linear $A\tilde{x} = \tilde{b}$ system into a crisp system of linear equations. The fuzzy minimal solution of the fuzzy matrix equation is derived from solving the crisp function system. Moreover, the existence condition of the strong minimal fuzzy solution is discussed. Finally, some examples are given to illustrate our method.

Dequan Shang, College of Public Health, Gansu Traditional Chinese Medicine University, Lanzhou 730000, China

Jinglun Li, College of Mathematics and Statistics, Northwest Normal University, Lanzhou 730070, China.

Xiaobin Guo, College of Mathematics and Statistics, Northwest Normal University, Lanzhou 730070, China

The work is supported by the Natural Scientific Funds of PR China (no: 61262022).

II. PRELIMINARIES

There are several definitions for the concept of fuzzy numbers.

2.1. The fuzzy number

Definition 2.1. A fuzzy number is a fuzzy set like $u: R \rightarrow I = [0, 1]$ which satisfies:

- (1) u is upper semi-continuous,
- (2) u is fuzzy convex,

i.e., $u(\lambda x + (1 - \lambda)y) \geq \min\{u(x), u(y)\}$ for all $x, y \in R, \lambda \in [0, 1]$,

- (3) u is normal, i.e., there exists $x_0 \in R$ such that

$$u(x_0) = 1,$$

- (4) $\text{supp } u = \{x \in R \mid u(x) > 0\}$ is the support of the u ,

and its closure $\text{cl}(\text{supp } u)$ is compact.

Let E^1 be the set of all fuzzy numbers on R .

Definition 2.2. A fuzzy number \tilde{u} in parametric form is a pair (\underline{u}, \bar{u}) of functions $\underline{u}(r), \bar{u}(r), 0 \leq r \leq 1$; which satisfies the requirements:

- (1) $\underline{u}(r)$ is a bounded monotonic increasing left continuous function,

- (2) $\bar{u}(r)$ is a bounded monotonic decreasing left continuous function,

- (3) $\underline{u}(r) \leq \bar{u}(r), 0 \leq r \leq 1$.

A crisp number x is simply represented

by $(\underline{u}(r), \bar{u}(r)) = (x, x), 0 \leq r \leq 1$. By appropriate

definitions the fuzzy number space $\{\underline{u}(r), \bar{u}(r)\}$ becomes a convex cone E^1 which could be embedded isomorphically and isometrically into a Banach space.

Let

$$\tilde{x} = (\underline{x}(r), \bar{x}(r)), y = (\underline{y}(r), \bar{y}(r)), 0 \leq r \leq 1, \text{ and } k \in R. \text{ Then}$$

Then

$$(1) \tilde{x} = \tilde{y} \text{ iff } \underline{x}(r) = \underline{y}(r), \bar{x}(r) = \bar{y}(r),$$

$$(2) \tilde{x} + \tilde{y} = (\underline{x}(r) + \underline{y}(r), \bar{x}(r) + \bar{y}(r)),$$

$$(3) \tilde{x} - \tilde{y} = (\underline{x}(r) - \underline{y}(r), \bar{x}(r) - \bar{y}(r)),$$

$$(4) k\tilde{x} = \begin{cases} (k\underline{x}, k\bar{x}), & k \geq 0 \\ (k\bar{x}, k\underline{x}), & k < 0 \end{cases}$$

Definition 2.3. A matrix

$$\tilde{A} = (\tilde{a}_{ij}), 0 \leq i \leq m, 0 \leq j \leq n$$

is called a fuzzy matrix, if each element \tilde{a}_{ij} of \tilde{A} is a fuzzy number. Particularly, a vector $\tilde{x} = (\tilde{x}_j), 0 \leq j \leq n$ is called a fuzzy vector, if each element \tilde{x}_j of \tilde{x} is a fuzzy number

Definition 2.4. Let $A = (a_{ij})$ be a $m \times n$ crisp matrix and be a $\tilde{B} = (\tilde{b}_{ij}), 0 \leq i \leq n, 0 \leq j \leq p$ fuzzy matrix. The size of the product of two matrices is $m \times p$ and is written as follows:

$$A\tilde{B} = \tilde{C} = (\tilde{c}_{ij}),$$

where

$$\tilde{c}_{ij} = \sum_{k=1, \dots, n}^+ a_{ik} \times \tilde{b}_{kj}$$

Definition 2.5. The linear system equation

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \begin{pmatrix} \tilde{x}_1 \\ \tilde{x}_2 \\ \vdots \\ \tilde{x}_n \end{pmatrix} = \begin{pmatrix} \tilde{b}_1 \\ \tilde{b}_2 \\ \vdots \\ \tilde{b}_m \end{pmatrix}, \quad (2.1)$$

Where $a_{ij}, 0 \leq i \leq m, 0 \leq j \leq n$ are crisp numbers and $\tilde{b}_i, 1 \leq i \leq m$ are fuzzy numbers, is called a fuzzy linear system (FLS).

Using matrix notation, we have

$$A\tilde{x} = \tilde{b} \quad (2.2)$$

A fuzzy numbers vector

$$\tilde{x} = (\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n)^T$$

is called a fuzzy solution of the fuzzy linear system if and only if \tilde{x} satisfies (2.6).

III. SOLVING FUZZY LINEAR SYSTEMS

Definition 3.1. A fuzzy numbers vector \tilde{u} in parametric form is a pair (\underline{u}, \bar{u}) of vector functions $\underline{u}(r), \bar{u}(r)$ where

$$\underline{u}(r) = (u_1(r), u_2(r), \dots, u_n(r))^T,$$

$$\bar{u}(r) = (\bar{u}_1(r), \bar{u}_2(r), \dots, \bar{u}_n(r))^T,$$

$0 \leq r \leq 1$, which satisfies the requirements:

- (1) $\underline{u}(r)$ is a bounded monotonic increasing left continuous vector function,
- (2) $\bar{u}(r)$ is a bounded monotonic decreasing left continuous vector function,
- (3) $\underline{u}(r) \leq \bar{u}(r), 0 \leq r \leq 1$.

Definition 3.2. Let

$\tilde{u} = (\underline{u}(r), \bar{u}(r)), v = (\underline{v}(r), \bar{v}(r)) \in E^n, 0 \leq r \leq 1$ be two fuzzy vectors and $k \in R$. Then

- (1) $\tilde{u} = \tilde{v}$ iff $\underline{u}(r) = \underline{v}(r)$ and $\bar{u}(r) = \bar{v}(r)$
- (2) $\tilde{u} + \tilde{v} = (\underline{u}(r) + \underline{v}(r), \bar{u}(r) + \bar{v}(r))$,
- (3) $\tilde{u} - \tilde{v} = (\underline{u}(r) - \underline{v}(r), \bar{u}(r) - \bar{v}(r))$,

$$(4) k\tilde{u} = \begin{cases} (k\underline{u}, k\bar{u}), k \geq 0. \\ (k\bar{u}, k\underline{u}), k < 0. \end{cases}$$

Theorem 3.1. The fuzzy matrix equation (2.6) can be extended to a crisp function linear system as follows

$$Sx(r) = b(r) \quad (3.1)$$

where

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}, x(r) = \begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix}, b(r) = \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix},$$

in which the elements a_{ij}^+ of matrix A^+ and a_{ij}^- of matrix A^- are determined by the following way:

if $a_{ij} \geq 0, a_{ij}^+ = a_{ij}$ else $a_{ij}^+ = 0, 1 \leq i \leq m, 1 \leq j \leq n$;

if $a_{ij} < 0, a_{ij}^- = a_{ij}$ else $a_{ij}^- = 0, 1 \leq i \leq m, 1 \leq j \leq n$.

Proof. We denote the right fuzzy vector of Eqs.(2.2) \tilde{b}

with $\tilde{b} = [\underline{b}(r), \bar{b}(r)], 0 \leq r \leq 1$ and the unknown fuzzy vector denoted by

$$\tilde{x} = [\underline{x}(r), \bar{x}(r)] = [\underline{x}_j(r), \bar{x}_j(r)]_{n \times 1}.$$

We also suppose $A = A^+ + A^-$ in which the elements a_{ij}^+ of matrix A^+ and a_{ij}^- of matrix A^- are determined by the following way:

if $a_{ij} \geq 0, a_{ij}^+ = a_{ij}$ else $a_{ij}^+ = 0, 1 \leq i \leq m, 1 \leq j \leq n$;

if $a_{ij} < 0, a_{ij}^- = a_{ij}$ else $a_{ij}^- = 0, 1 \leq i \leq m, 1 \leq j \leq n$.

For fuzzy linear equation $A\tilde{x} = \tilde{b}$, we can express it as

$$(A^+ + A^-)[\underline{x}(r), \bar{x}(r)] = [\underline{b}(r), \bar{b}(r)]. \quad (3.2)$$

Since

$$k\tilde{x}_j = \begin{cases} (k\underline{x}_j, k\bar{x}_j), k \geq 0 \\ (k\bar{x}_j, k\underline{x}_j), k < 0 \end{cases},$$

we have

$$A\tilde{x}_j = \begin{cases} (A\underline{x}_j, A\bar{x}_j), k \geq 0 \\ (A\bar{x}_j, A\underline{x}_j), k < 0 \end{cases},$$

so the Eqs.(3.2) be rewritten is

$$[A^+ \underline{x}(r) + A^- \bar{x}(r), A^+ \bar{x}(r) + A^- \underline{x}(r)] = [\underline{b}(r), \bar{b}(r)]$$

Thus we get

$$\begin{cases} A^+ \underline{x}(r) + A^- \bar{x}(r) = \underline{b}(r) \\ A^+ \bar{x}(r) + A^- \underline{x}(r) = \bar{b}(r) \end{cases} \quad (3.3)$$

or

$$\begin{cases} A^+ \underline{x}(r) - A^- (-\bar{x}(r)) = \underline{b}(r) \\ -A^+ \bar{x}(r) - A^- \underline{x}(r) = -\bar{b}(r) \end{cases}$$

Expressing Eqs.(3.3) in matrix form, we have

$$\begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix} \begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix} = \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix}. \quad \square$$

In order to solve the fuzzy linear equation (2.2), we need to consider the systems of linear equations (3.1). We consider a specious case, that is, the coefficient matrix A is a $n \times n$ matrix. By the M. Freidman[16], the following results are

apparent.

Theorem 3.2. The matrix S is nonsingular if and only if the matrices $A^+ + A^-$ and $A^+ - A^-$ are both nonsingular. If S is nonsingular, then

$$S^{-1} = \frac{1}{2} \begin{pmatrix} (A^+ - A^-)^{-1} + (A^+ + A^-)^{-1} & (A^+ - A^-)^{-1} - (A^+ + A^-)^{-1} \\ (A^+ - A^-)^{-1} - (A^+ + A^-)^{-1} & (A^+ - A^-)^{-1} + (A^+ + A^-)^{-1} \end{pmatrix} \quad (3.4)$$

where $(A^+ + A^-)^{-1}, (A^+ - A^-)^{-1}$ are inverse matrices of matrices $A^+ + A^-$ and $A^+ - A^-$, respectively. In this case, the solution of model equation (3.1) is

$$\begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix} = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}^{-1} \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix} \quad (3.5)$$

By matrix theory analysis, we obtained the minimal solution [3] of the function linear system (3.1) as

$$x(r) = S^\dagger b(r),$$

i.e.

$$\begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix} = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}^\dagger \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix}. \quad (3.6)$$

Where S^\dagger is the Moore-Penrose generalized inverse of matrix S .

However, the solution matrix may still not be an appropriate fuzzy numbers vector. Restricting the discussion to triangular fuzzy numbers, i.e., $\underline{b}_j(r), \bar{b}_j(r), 1 \leq j \leq n$ and consequently $\underline{x}_i(r), \bar{x}_i(r), 1 \leq i \leq n$ are all linear functions of r , and having calculated $x(r)$ which solves (3.1), we define the fuzzy minimal solution to the fuzzy linear systems (2.2) as follows.

Definition 3.3. Let $x(r) = \underline{x}_i(r), \bar{x}_i(r), 1 \leq i \leq n$ be the minimal solution of (3.1). The fuzzy number matrix $\underline{u}_i(r), \bar{u}_i(r), 1 \leq i \leq n$ defined by

$$\begin{cases} \underline{u}_i(r) = \min\{\underline{x}_i(r), \bar{x}_i(r), \underline{x}_i(1), \bar{x}_i(1)\}, \\ \bar{u}_i(r) = \max\{\underline{x}_i(r), \bar{x}_i(r), \underline{x}_i(1), \bar{x}_i(1)\}, \\ 1 \leq i \leq n, 0 \leq r \leq 1. \end{cases} \quad (3.7)$$

is called the fuzzy minimal solution of the function linear systems (3.1). If $\underline{x}_i(r), \bar{x}_i(r), 1 \leq i \leq n$ are all fuzzy numbers then $\underline{u}_i(r) = \underline{x}_i(r), \bar{u}_i(r) = \bar{x}_i(r), 1 \leq i \leq n$ and U is called a strong minimal fuzzy solution of the fuzzy linear systems (2.6). Otherwise, U is called a weak minimal fuzzy solution.

To illustrate the expression (3.6) to be a fuzzy solution vector, we now discuss the generalized inverses of non negative matrix

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}$$

in a special structure.

Lemma 3.1 [11]. Let

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}$$

Then the matrix

$$S^\dagger = \frac{1}{2} \begin{pmatrix} (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger & (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger \\ (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger & (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger \end{pmatrix}$$

(3.8)

is the Moore-Penrose inverse of the matrix S ,

where $(A^+ - A^-)^\dagger, (A^+ + A^-)^\dagger$ are Moore-Penrose inverses of matrices $A^+ - A^-$ and $A^+ + A^-$, respectively.

The key points to make the solution matrix being a strong fuzzy solution is that $S^\dagger b(r)$ is fuzzy matrix, i.e., each element in which is a triangular fuzzy number. By the analysis, it is equivalent to the condition $S^\dagger \geq O$.

Theorem 3.3. If

$(A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger \geq O, (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger \geq O$, the fuzzy linear equation (2:6) has a strong fuzzy minimal solution as follows:

$$\tilde{x} = [\underline{x}(r), \bar{x}(r)]$$

where

$$\begin{cases} \underline{x}(r) = E \underline{b}(r) - \bar{b}(r), \\ \bar{x}(r) = -F \underline{b}(r) + E \bar{b}(r) \\ E = \frac{1}{2} ((A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger) \\ F = \frac{1}{2} ((A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger) \end{cases} \quad (3.9)$$

Proof. Let

$$S^\dagger = \begin{pmatrix} E & F \\ F & E \end{pmatrix} = \frac{1}{2} \begin{pmatrix} (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger & (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger \\ (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger & (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger \end{pmatrix}$$

, we know the condition that $S^\dagger \geq O$ is equivalent to $E \geq O, F \geq O$.

Since $\tilde{b} = [\underline{b}(r), \bar{b}(r)], \underline{b}(r)$ is a bounded monotonic increasing left continuous function vector and $\bar{b}(r)$ a bounded monotonic decreasing left continuous function vector with $\underline{b}(r) \leq \bar{b}(r), 0 \leq r \leq 1$ by Definition 3.1.

According to Eqs.(3.6), we have

$$\begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix} = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}^\dagger \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix},$$

i.e.,

$$\underline{x}(r) = E \underline{b}(r) - \bar{b}(r),$$

$$\bar{x}(r) = -F \underline{b}(r) + E \bar{b}(r).$$

Now that $E \geq O, F \geq O$ and $\underline{b}(r), -\bar{b}(r)$ are bounded monotonic increasing left continuous function vectors, we know that $\underline{x}(r)$ is a bounded monotonic increasing left continuous function vector and $\bar{x}(r)$ is a bounded monotonic decreasing left continuous function vector. And $\bar{x}(r) - \underline{x}(r) = E(\bar{b}(r) - \underline{b}(r)) + F(\bar{b}(r) - \underline{b}(r)) = (E + F)(\bar{b}(r) - \underline{b}(r)) = (A^+ - A^-)^\dagger(\bar{b}(r) - \underline{b}(r)) \geq O$.

Thus the fuzzy matrix equation (2.6) has a strong minimal fuzzy solution. □

The following Theorems give some results for such S^{-1} and S^\dagger to be nonnegative. As usual, $(\cdot)^T$ denotes the transpose of a matrix (\cdot) .

Theorem 3.6[25]. The inverse S^{-1} of a nonnegative matrix S is nonnegative if and only if S is a generalized permutation matrix.

Theorem 3.7[12]. Let S be an $2n \times 2n$ nonnegative matrix with rank r . Then the following assertions are equivalent:

(a). $S^\dagger \geq 0$:

(b). There exists a permutation matrix P , such that PS has the form

$$PS = \begin{pmatrix} T_1 \\ T_2 \\ \vdots \\ T_r \\ O \end{pmatrix},$$

where each T_i has rank 1 and the rows of T_i are orthogonal to the rows of T_j , whenever $i \neq j$, the zero matrix may be absent.

(c). $S^\dagger = \begin{pmatrix} GC^T & GD^T \\ GD^T & GC^T \end{pmatrix}$

for some positive diagonal matrix G . In this case, $(C + D)^\dagger = G(C + D)^T, (C - D)^\dagger = G(C - D)^T$.

IV. NUMERICAL EXAMPLES

Example 4.1. Consider the following fuzzy linear system

$$\begin{pmatrix} 1 & -1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} \tilde{x}_1 \\ \tilde{x}_2 \end{pmatrix} = \begin{pmatrix} (r, 2-r) \\ (4+r, 7-2r) \end{pmatrix}.$$

Let

$$\tilde{x} = [\underline{x}(r), \bar{x}(r)] = \left[\begin{pmatrix} \underline{x}_1(r) \\ \underline{x}_2(r) \end{pmatrix}, \begin{pmatrix} \bar{x}_1(r) \\ \bar{x}_2(r) \end{pmatrix} \right], A = A^+ + A^- = \begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} + \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}$$

and

$$\tilde{b} = [\underline{b}(r), \bar{b}(r)] = \left[\begin{pmatrix} r \\ 4+r \end{pmatrix}, \begin{pmatrix} 2-r \\ 7-2r \end{pmatrix} \right].$$

By the Theorem 3.1., the original fuzzy matrix equation is equivalent to the following function linear system

$$Sx(r) = b(r),$$

where

$$x(r) = \begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix}, b(r) = \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix}$$

and

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 2 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 2 \end{pmatrix}.$$

The matrix S is obviously invertible, its inverse is

$$S^\dagger = \begin{pmatrix} 1.3333 & -0.3333 & 0.6667 & -0.6667 \\ -0.6667 & 0.6667 & -0.3333 & 0.3333 \\ 0.6667 & -0.6667 & 1.3333 & -0.3333 \\ -0.3333 & 0.3333 & -0.6667 & 0.6667 \end{pmatrix}.$$

From Eqs.(3.6), the solution of model is as follows:

$$\begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix} = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}^\dagger \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix} \\ = \frac{1}{10} \begin{pmatrix} 1.3333 & -0.3333 & 0.6667 & -0.6667 \\ -0.6667 & 0.6667 & -0.3333 & 0.3333 \\ 0.6667 & -0.6667 & 1.3333 & -0.3333 \\ -0.3333 & 0.3333 & -0.6667 & 0.6667 \end{pmatrix} \begin{pmatrix} r \\ 4+r \\ -2+r \\ -7+2r \end{pmatrix} \\ = \begin{pmatrix} (2.0000 + 0.3333r) \\ (1.0000 + 0.3333r) \end{pmatrix}, \begin{pmatrix} -3.0000 + 0.6667r \\ -2.0000 + 0.6667r \end{pmatrix}$$

i.e.,

$$\begin{pmatrix} \tilde{x}_1 \\ \tilde{x}_2 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} (2.0000 + 0.3333r, 3.0000 - 0.6667r) \\ (1.0000 + 0.3333r, 2.0000 - 0.6667r) \end{pmatrix}.$$

Since \tilde{x} is an appropriate triangular fuzzy numbers vector, we obtained the solution of the fuzzy linear system is

$$\tilde{U} = \begin{pmatrix} \tilde{u}_1 \\ \tilde{u}_2 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} (2r, 4 - 2r) \\ (-2 + r, -r) \end{pmatrix}.$$

it admits a strong fuzzy solution by Definition 3.3.

Example 4.2. Consider the following fuzzy linear system

$$\begin{pmatrix} 2 & -1 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} \tilde{x}_1 \\ \tilde{x}_2 \end{pmatrix} = \begin{pmatrix} (r, 2-r) \\ (-2+r, -r) \end{pmatrix}.$$

Let

$$\tilde{x} = [\underline{x}(r), \bar{x}(r)] = \left[\begin{pmatrix} \underline{x}_1(r) \\ \underline{x}_2(r) \end{pmatrix}, \begin{pmatrix} \bar{x}_1(r) \\ \bar{x}_2(r) \end{pmatrix} \right], A = A^+ + A^- = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & -1 \\ -2 & 0 \end{pmatrix}$$

and

$$\tilde{b} = [\underline{b}(r), \bar{b}(r)] = \left[\begin{pmatrix} r \\ -2+r \end{pmatrix}, \begin{pmatrix} 2-r \\ -r \end{pmatrix} \right].$$

By the Theorem 3.1., the original fuzzy matrix equation is equivalent to the following function linear system

$$Sx(r) = b(r),$$

where

$$x(r) = \begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix}, b(r) = \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix}$$

and

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 2 & 2 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

The matrix S is obviously not invertible, its Moore-Penrose inverse is

$$S^\dagger = \frac{1}{10} \begin{pmatrix} 2 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 2 & 2 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \geq O.$$

From Eqs.(3.6), the minimal solution of model is as follows:

$$\begin{aligned} \begin{pmatrix} \underline{x}(r) \\ -\bar{x}(r) \end{pmatrix} &= \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}^\dagger \begin{pmatrix} \underline{b}(r) \\ -\bar{b}(r) \end{pmatrix} \\ &= \frac{1}{10} \begin{pmatrix} 2 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 2 & 2 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \left[\begin{pmatrix} r \\ -2+r \end{pmatrix}, \begin{pmatrix} -2+r \\ r \end{pmatrix} \right] \\ &= \frac{1}{5} \left[\begin{pmatrix} 2r \\ -1+2r \end{pmatrix}, \begin{pmatrix} -4+2r \\ r \end{pmatrix} \right] \end{aligned}$$

i.e.,

$$\begin{pmatrix} \tilde{x}_1 \\ \tilde{x}_2 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} (2r, 4-2r) \\ (-2+r, -r) \end{pmatrix}.$$

Since \tilde{x} is an appropriate triangular fuzzy numbers vector, we obtained the solution of the fuzzy linear system is

$$\tilde{U} = \begin{pmatrix} \tilde{u}_1 \\ \tilde{u}_2 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} (2r, 4-2r) \\ (-2+r, -r) \end{pmatrix}.$$

it admits a strong fuzzy solution by Definition 3.3.

V. CONCLUSION

In this paper we proposed a general model for solving the fuzzy linear equation $A\tilde{x} = \tilde{b}$ where A is a crisp matrix and \tilde{b} is an arbitrary fuzzy numbers matrix respectively by a matrix method. We extended the fuzzy linear system into a crisp system of linear equations and obtained the fuzzy minimal solution by solving the crisp function system. Moreover, the existence condition of the strong minimal fuzzy solution is discussed respectively. Numerical examples showed that our method is feasible to solve the fuzzy linear system.

ACKNOWLEDGMENT

This research was supported by the National Natural Science Foundation of China (no. 61262022) and the Youth Scientific Research Ability Promotion Project of Northwest Normal University (NWNLU-LKQN-15-20).

REFERENCES

[1] S. Abbasbandy, A. Jafarian, R. Fzzati, Conjugate gradient method for fuzzy symmetric positive definite system of linear equations, Applied Mathematics and Computation, 171(2005) 1184-1191.
 [2] S. Abbasbandy, R. Ezzati, A. Jafarian, LU decomposition method for solving fuzzy system of linear equations, Applied Mathematics and Computation, 172 (2006) 633-643.
 [3] S. Abbasbandy, M. Otadi, M. Mosleh, Minimal solution of general dual fuzzy linear systems, Chaos, Solitons and Fractals, 29 (2008) 638-652.

[4] T. Allahviranloo, Numerical methods for fuzzy system of linear equations, Applied Mathematics and Computation, 153 (2004) 493-502.
 [5] T. Allahviranloo, Successive over relaxation iterative method for fuzzy system of linear equations, Applied Mathematics and Computation, 162 (2005) 189-196.
 [6] T. Allahviranloo, The adomian decomposition method for fuzzy system of linear equations, Applied Mathematics and Computation, 163 (2005) 553-563
 [7] T. Allahviranloo, F. Hosseinzadeh Lot_, M. KhorasaniKiasari, M. Khezerloo, On the fuzzy solution of LR fuzzy linear systems, Applied Mathematical Modelling, 37(2013) 1170-1176.
 [8] T. Allahviranloo, M. Ghanbari, R. Nuraei, A note on /Fuzzy linear systems, Fuzzy Sets and Systems, 177 (2011) 876-92.
 [9] B. Asady, S. Abbasbandy, M. Alavi, Fuzzy general linear systems, Applied Mathematics and Computation, 169 (2005) 34-40.
 [10] N. Babbar, A. Kumar, Abhinav Bansal, Solving fully fuzzy linear system with arbitrary triangular fuzzy numbers, Soft Computing, 17 (2013) 6916702.
 [11] A. Ben-Israel, T.N.E. Greville, Generalized inverses: theory and applications, second edition, Springer-Verlag, New York, 2003.
 [12] A. Berman, R.J. Plemmons, Nonnegative matrices in the Mathematical Sciences, Academic press, New York, 1979.
 [13] M. Dehghan, B. Hashemi, M. Ghatsee, Solution of the full fuzzy linear systems using iterative techniques, Chaos, Solitons and Fractals, 34 (2007) 316-336.
 [14] M. Dehghan, B. Hashemi, Iterative solution of fuzzy linear systems, Applied Mathematics and Computation, 175 (2006) 645-674.
 [15] D. Dubois, H. Prade, Operations on fuzzy numbers, Journal of Systems Science, 9 (1978)613-626.
 [16] M. Friedman, M. Ma, A. Kandel, Fuzzy linear systems, Fuzzy Sets and Systems, 96 (1998)201-209.
 [17] R. Ghanbari, Solutions of fuzzy LR algebraic linear systems using linear programs, Applied Mathematical Modelling, 39 (2015) 516465173.
 [18] R. Goetschel, W. Voxman, Elementary calculus, Fuzzy Sets and Systems, 18 (1986) 31-43.
 [19] Z.T. Gong, X.B. Guo, Inconsistent fuzzy matrix equations and its fuzzy least squares solutions, Applied Mathematical Modelling, 35 (2011) 1456-1469.
 [20] X.B. Guo, K. Zhang, Solving fuzzy matrix equation of the form $\tilde{X}A = \tilde{B}$, Journal of Intelligent and Fuzzy Systems, 32 (2017) 2771-2778.
 [21] M. Ma, M. Friedman, A. Kandel, Duality in Fuzzy linear systems, Fuzzy Sets and Systems, 109 (2000) 55-58.
 [22] S. Nahmias, Fuzzy variables, Fuzzy Sets and Systems, 1(2)(1978) 97-111.
 [23] R. Nuraei, T. Allahviranloo, M. Ghanbari, Finding an inner estimation of the solution set of a fuzzy linear system, Applied Mathematical Modelling, 37(2013) 5148-5161.
 [24] M. Otadi, Maryam Mosleh, Solving fully fuzzy matrix equations, Applied Mathematical Modelling, 36 (2012) 6114-6121.
 [25] R.J. Plemmons, Regular nonnegative matrices, Proceedings of the American Mathematical Society, 39 (1973) 26-32.
 [26] M.L. Puri, D.A. Ralescu, Differentials for fuzzy functions, Journal of Mathematics Analysis and Application, 91 (1983) 552-558.
 [27] C.X. Wu, M. Ma, Embedding problem of fuzzy number space: Part I, Fuzzy Sets and Systems, 44 (1991) 33-38.
 [28] C.X. Wu, M. Ma, Embedding problem of fuzzy number space: Part III, Fuzzy Sets and Systems, 46 (1992) 281-286.
 [29] L.A. Zadeh, Fuzzy sets, Information and Control, 8 (1965) 338-353.
 [30] B. Zheng, K. Wang, General fuzzy linear systems, Applied Mathematics and Computation, 181 (2006) 1276-1286.