Review on Phytochemistry and Pharmacological Aspects of Guiera Senegalensis J. F. Gmel (Combretaceae)

Adama Dénout, Adiaratou Togola, Mahamane Haïdara, Rokia Sanogo, Drissa Diallo ,Mamadou Koumaré

Abstract— Medicinal plants are the local heritage with global importance. They have curative properties due to presence of various complex chemical substances of different composition, which are found as secondary plant metabolites in one or more parts of these plants. These plant metabolites according to their composition are grouped as flavonoids, tannins, alkaloids, saponins etc...Guiera senegalensis J. F. Gmel. (Combretaceae) is one of the most important West African medicinal plants, often used to treat a variety of microbial infections.

Index Terms— Phytochemistry, Pharmacological Aspects, Guiera Senegalensis.

I. INTRODUCTION

Guiera senegalensis is called Sabara" (Hausa), [1]. It is a shrubby and can grow to a height of 3 to 5 m depending on the habitat [2]. The leaves which are 3 to 5 cm long and 1.5 to 3.0 cm broad are arranged opposite or sub opposite on the stem [3]. It is widely distributed in the savannah region of west and central Africa, Nigeria, Senegal, Gambia, Mali, Niger, Burkina Faso and Ghana, [4-6]. It is active against cough, respiratory congestion and fever [7,8], and is prescribed as an antitussive, hypertension and hypotension as well as venereal diseases [7], to ease breathing and to treat lung and bronchial disorders. It is also used against malaria fever [9].

II. BOTANICAL DESCRIPTION

A. Scientific Classification

Kingdom: Plantae Division: Magnoliophyta Class: Magnoliopsida Order: Myrtales Family: Combretaceae Genus: Guiera

Species: senegalensis

Adama Dénou, Faculty of Pharmacy, University of Sciences Techniques and Technologies of Bamako (Mali);

Adiaratou Togola, Department of Traditional Medicine, National Institute of Research in Public Health (Mali);

Mahamane Haïdara, Faculty of Pharmacy, University of Sciences, Techniques and Technologies of Bamako (Mali);

Rokia Sanogo, Faculty of Pharmacy and Department of Traditional Medicine (Mali);

Drissa Diallo, Faculty of Pharmacy and Department of Traditional Medicine (Mali):

 ${\bf Mamadou\ Koumar\'e, Malian\ Society\ of\ Phytotherapy\ (Bamako-Mali).}$

B. Local names [10]

Bamanan : N'Kunje

Bomou : Suncawe, Sunlayi

Dogon : Guru, Tulu Hausa : Sabara Kasonké : Kangano Peulh : N'Geloki

Senoufo : Konifire, Kogbwe

Sonraï : Sabara Wolof : N'Ger Tamacheck : Subara

C. Vernacular name: Nuger

D. Morphology

Guiera senegalensis, very well known in its native area, generally occurs as a shrub that can grow to a height of 3 to 5 m according to habitat. Its stem presents numerous knots that send out branches. The ash-grey stem and branches have fibrous or pubescent bark and bear opposing, short petiolated oval leaves, sometimes mucronate, sometimes even cordate at their base, about 2 to 4 cm long by 1 to 2 cm wide. These grey-green leaves, darker on their upper surface, display black spots on their lower surface and are slightly downy on both sides. These features lend the plant an overall silver green colour that is conspicuous in brushland [2]. Flowering occurs almost throughout the year, when it is leafy. Often blooms twice a year, during the dry season and the rainy season. Each flower has a calcinal tube ovoid, welded to the ovary. This tube is topped by a bellflower blade with 5 teeth screened black and persistent points to fruiting (Figure 1). The ligulform corolla is composed of 5 petals also riddled with black spots. The stamens are 10 on two rows of 5, all inserted on the calyx. The ovary has a single box containing 4 to 6 eggs [11].

The fruit is an achene (Figure 1) around 3 cm long brown or green ash, spindle-shaped, hairy with sides and the remains of the calyx [11].



Fig 1: Flowery and fruity branches of Guiera senegalensis J.F. Gmel (Combretaceae) by Mamadou Koumaré



III. HABITAT AND REPARTITION

It is a plant that grows primarily in Sudanian Sahel area, on soils sandy, leached or exhausted, fallow and dry stations. Plant pioneer disseminated by cattle in the fallow land, it is also indicative of overgrazing. It is found from Senegal to Cameroon to Sudan. Widespread and common, locally gregarious and very abundant [12].

IV. PHYTOCHEMISTRY

Earlier works showed that leaves and roots of G. senegalensis contained harman, tetrahydroharman, harmalan and Guieranone A [1]. The plant has been found to contain carbohydrates, steroids, flavonoids, saponins, alkaloids, tannins and mucilage [24, 26]. Ficarra et al. in 1997 found four flavonoids in the leaves of G. senegalensis, namely catechin, myricitrin, rutin and querterin [13]. Some findings on the plant elsewhere showed the presence of alkaloids (Hyoscyamin [I] and solanine, [II]), tannins, terpenoids menthol, coumarins, saponins, flavonoids (quercetin,[III]), cardiotonics and cyanogenic heterosides which were assayed in various organs of the plant leaves, stem bark, fruits and roots [12]. Hyoscyamin [I], solanine [II] and quercetin [III] are represented in figure 2. While in the gall of the plant from Ouagadougou, Burkina Faso, alkaloids, polyphenols and saponins were detected [23]. The ashes are poor in alkali but rich in alkaline earth metals. We found especially Mg, Ca, Sr, Ti, Fe, Al and in lesser amounts sometimes traces: Cu, Ni, Co, Zn [11]. Previous studies indicated the presence in leaves of two alkaloids (harman, tetrahydroharman or eleagnine), flavonoids, naph-thopyrans, tannins, and a naphthyl butenone (guieranone A); in roots, were only obtained tannins and the same beta-carboline alkaloids than in leaves [9, 11, 29, 30, 31, 32, 33, 34].




Fig 2:Chemical structure from Guiera senegalensis [12].

V. TRADITIONAL USES

G. senegalensis is considered by traditional practitioners of Burkina Faso, Senegal, and Mali as a panacea, both its medicinal properties are significant and varied.

The usual form of preparation for internal use is in decoction or mixed with food preparations. G. senegalensis leaves are widely administered for pulmonary and respiratory complaints, for coughs, as a febrifuge, colic and diarrhea, syphilis, beriberi, leprosy, impotence, rheumatism, diuresis and expurgation [3,4].

VI. PARMACOLOGICAL AND TOXICOLOGICAL ACTIVITIES

Guiera senegalensis is a popular medicinal plant for both human and veterinary use in West Africa. Its leaves have been reportedly used to treat rhinitis, bronchitis and fever, and the roots to treat diarrhea and dysentery. It is recognized as being active against cough, respiratory congestion and fever [14], and is prescribed as an antitussive [11,15,16,17,18], to ease breathing and to treat lung and bronchial disorders. It is also used against malaria [9,19,20]. The syrup D2 from G. senegalensis has been screened for its antitussive clinical essay [10]. Aqueous extracts from its roots and leaves have also been screened for toxicity [21]. The galls of G. senegalensis are used in Burkina Faso as an ethnoveterinary product to increase milk production in cows and to treat fowlpox infection in chickens [22].

Guiera senegalensis leaves extract or fractions may be used for the treatment of various disease caused by Escherichia coli, Pseudomonas aeruginosa and Klebsiella pneumonia [24].

View toxicity aqueous extracts are overall low toxicity; "extract of leaves being slightly more active than the extract from roots in the Guinea pig, rabbit, dog and less active in mice and rats [11]. G. Senegalensis at lower doses is not harmful to the liver and therefore can be exploited as it is served in the treatments of some illnesses [25]. Treatment with water and methanol extracts from G. senegalensis leaves resulted in endotheliotoxicity, hepatonephropathy and pancreatic hyperplasia [27]. Guieranone A from Guiera senegalensis showed a strong antiplasmodial activity associated with a high cytotoxicity toward human monocytes [1]. Methanol fraction from leaves extract only had antiplasmodial activity [28].

VII. CONCLUSION

Herbal drug which are used in various traditional medicine, needs detailed investigation with ethno-pharmacological approach. In the present review we have made to explorer the all details of Guiera senegalensis information its botany, habitat, ethnoveterinary, traditional and modern uses.

REFERENCES

- [1] Fiot J, Sanon S, Azas N, et al. Phytochemical and pharmacological study of roots and leaves of *Guiera senegalensis* J.F. Gmel (Combretaceae). Journal of ethnopharmacology 2006; 106(2):173-178.
- [2] Silva O, Serrano R, Gomes ET. Botanical characterization of *Guiera senegalensis* leaves. Microscopy and Microanalysis. Official journal of microscopy society of America 2008; 14(5):398-404.
- [3] Hutchinson J, Dalziel JM. Flora of West Tropical Africa. London: Crown Agents, 1972.
- [4] Zeljan M, Marica M, Franz B. Flavonoida of G. senegalensis-Thin layer Chromatography and Numerical Methods. Croatica Chemica Acta 1998; 71(1):69-79.
- [5] Shettima YA, Tijjani MA, Kanumi Y, et al. Phytochemical and antidiarrhoeal properties of methanol extract of Guiera senegalensis J.F. Gmel International Journal of Pharmacy, 2012, 3(11).
- [6] Tijjani A, Sallau MS, Sunusi I. Synergistic activity of methanolic extract of Adenium obesum (Apocynaceae) stem-bark and xytetracycline against some clinical bacterial isolates. Bayero Journal of Pure and Applied Sciences 2011; 4(1):79-82.
- [7] Adedapo AA, Sofidiya MO, Afolayan AJ. Antiinflammatory and analgesic activities of the aqueous extracts of Margaritaria discoidea (Euphorbiaceae) stem bark in experimental animal models. (Int J Trop Biol. 2009; 57(4):1193-1200.
- [8] Ali AJ, Akanya HO, Dauda BEN, et al. Antiplasmodial, analgesic and anti-inflammatory effects of crude Guiera senegalensis J.F. Gmel (Combretaceae) leaf extracts in mice infected with Plasmodium berghei. Journal of Pharmacognosy and Phytotherapy, 2011, 3(10).



31 www.ijntr.org

- [9] Ancolio C, Azas N, Mahiou V, et al. Antimalarial activity of extracts and alkaloids isolated from six plants used in traditional medicine in Mali and Sao Tome. Phytotheraphy Research. 2002; 16(7):646-649.
- [10] Dénou A. Contribution à l'étude des problèmes de conservation de la forme sirop des médicaments traditionnels. Thèse de Pharmacie, FMPOS, Bamako. 2008, 82p.
- [11] Koumaré M. Contribution à l'étude pharmacologique du *Guiera senegalensis* Lam, Combretaceae. Thèse de Doctorat en pharmacie, Toulouse –France, 1968.
- [12] Somboro A A, Patel K, Diallo D, Sidibe L, Chalchat JC, Figueredo G, Ducki S, Troin Y, Chalard P. An ethnobotanical and phytochemical study of the African medicinal plant *Guiera senegalensis* J. F. Gmel. Journal of Medicinal Plants Research Vol. 2011, 5(9):1639-1651.
- [13] Ficarra R, Ficarra P, Tommasini S, Carulli M, Melardi S, Di Bella MR, Calabro ML, De Pasquale R, Germano MP, Sanogo R, Casuscelli. Isolation characterization of *Guiera senegalensis* J. F.Gmel., active principales. Boll. Chim. Farm. 1997, 136(5): 454-459.
- [14] Kerharo J, Adam JG. La Pharmacopée Sénégalaise Traditionnelle. Plantes Médicinales et Toxiques. Editions Vigot Frères, Paris. 1974, p. 1011.
- [15] Negrevergne G. New medicament based on a dry aqueous extract of Guiera senegalensis, especially useful as a hypotensive and antitussive agent. Fr. M. 1968, p. 2.
- [16] Faye O, Olschwamg D, Giono-Barber H, Pousset JL. Action antitussive d'un extrait lyophilisé de *Guiera senegalensis*. Plantes médicinales Africaines II, 1980, 25(4): 285-292.
- [17] Sanogo R, De Pasquale R, Germano MP. The Antitussive Activity of Guiera senegalensis J. F. Gmel. (Combretaceae). Phytother. Res., 1998, 12(2): 132-134.
- [18] Diatta W, Fall AD, Dieye AM, Faty S, Bassene E, Faye B. Experimental evidence of cough activity of total alkaloids from *Guiera senegalensis* Lam., In guinea pig. Dakar Med., 2007,52(2): 130-134.
- [19] Benoit F, Valentin A, Pelissier Y, Diafouka F, Marion C, Kone-Bamba D, Kone M, Mallie M, Yapo A, Bastide JM. *In vitro* antimalarial activity of vegetal extracts used in West African traditional medicine. Am. J. Trop. Med. Hyg., 1996,54(1): 67-71.
- [20] Azas N, Laurencin N, Delmas F, Di Giorgio C, Gasquet M, Laget M, Timon-David P, Synergistic in vitro antimalarial activity of plant extracts used as traditional herbal remedies in Mali. Parasitol. Res., 2002,88(2): 165-171.
- [21] Pousset JL. Plantes médicinales africaines: Utilisation pratiques. Paris: Ellipses- ACCT. 1989.
- [22] Nacoulma OG. Plantes médicinales et pratiques médicales traditionnelles au Burkina Faso. Cas du plateau central. Tome2. Ph.D. thesis, University of Ouagadougou. 1996.
- [23] Lamien CE, Meda A, Couacy-Hymann E, et al. The phytochemical composition and in vitro antiviral activity of decoctions from galls of Guiera senegalensis J.F. Gmel. (Combretaceae) and their relative non-toxicity for chickens. Onderstepoort Journal of Veterinary Research 2005; 72:111-118.
- [24] Salihu SO, Usman AA. Antimicrobial and phytochemical study of the bioactive fractions of *Guiera senegalensis* from Alasan Tambuwal, Nigeria. Journal of Pharmacognosy and Phytochemistry 2015; 3(6): 106-111.
- [25] Oshobu MA, Geidam MA. Effects of Aqueous Ethanolic Extract of the leaves of *Guiera senegalensis* J.F. Gmel (Combretaceae) on Liver Function in Wister Strain Albino Rats. International Journal of scientific research and management (IJSRM); 2014,2(3) 611-615.
- [26] Mohammed SY. Quantitative phytochemical and elemental analysis of Guiera senegalensis leaf extract. Journal of Pharmacognosy and Phytotherapy. 2013,5(12), pp. 204-207.
- [27] Fatih Elrahman AO, Abuelgasim AI, Galal M. Toxicopathological effects of *Guiera senegalensis* extracts in wistar albino rats. Journal of Medicinal Plants Research. 2008, 2(1), pp. 001 – 004.
- [28] Jigam AA, Akanya HO, Dauda BEN, Ogbadoyi EO. Antiplasmodial, analgesic and anti-inflammatory effects of crude *Guiera senegalensis* Gmel (Combretaceae) leaf extracts in mice infected with *Plasmodium* berghei. Journal of Pharmacognosy and Phytotherapy. 2011, 3(10), pp. 150-154.
- [29] Bouchet N, Levesque J, Pousset J.L. HPLC isolation, identification and quantification of tannins from *Guiera senegalensis*. Phytochemical A nalysis. 2000,11, 52–56.
- [30] Bucar F, Schubert-Zsilavecz M, Knauder E. Flavonoids of *Guiera senegalensis*. Pharmazie. 1996,51, 517–518.
- [31] Combier H, Becchi M, Cave A. Alcaloïdes du Guiera senegalensis Lam. Plantes Médicinales et Phytothérapie .1977,11, 251–253.
- [32] Mahmoud EN, Sami AK, 5-Methyldihydroflavasperone, a dihydronaphthopyran from *Guiera senegalensis*. Phytochemistry. 1997,46, 793–794.

[33] Silva O, Gomes ET. Guieranone A, a naphthyl butenone from the leaves of Guiera Senegalensis with antifungal activity. Journal of Natural Products. 2003, 66, 447–449.

Adama Denou, Pharmacist since 2008 (Faculty of Medicine, Pharmacy and Odonto-Stomatology, University of Bamako, Mali), this author became Assistant of Pharmacognosy in 2009, 2010-2012: MSc science field: biology, science branch: pharmacognosy (Faculty of Sciences, University of Lome, Togo).List of publications: google.scholar



Adiaratou Togola, Pharmacist since 2002 (Faculty of Medicine, Pharmacy and Odonto-Stomatology, University of Bamako, Mali). Researcher at Department of Traditional Medicine. Education: 2004-2008 PhD, science field: pharmacy, science branch: pharmacognosy (Faculty of Mathematics and Natural Sciences School of Pharmacy, School of Pharmacy, University of Oslo).List of publications: google.scholar

Mahamane Haïdara, Pharmacist since 2008 (Faculty of Medicine, Pharmacy and Odonto-Stomatology, University of Bamako, Mali), this author became Assistant of Pharmacognosy in 2009, 2011-2013: MSc science field: chemistry, science branch: pharmacognosy (University of Paul Sabatier, Toulouse, France). List of publications: google.scholar

Rokia Sanogo, Pharmacist since1988 (National School of Medicine and Pharmacy, Mali). Education: Professor of pharmacognosy since 2008 (Faculty of Pharmacy, University of Sciences, Techniques and Technologies of Bamako, Mali), this author became the chef of the Department of Traditional Medicine from 29th March 2016.List of publications: google.scholar

Drissa Diallo, Pharmacist since 1982 (National School of Medicine and Pharmacy, Mali). Education: Professor of pharmacognosy since 2004 (Faculty of Pharmacy, University of Sciences, Techniques and Technologies of Bamako, Mali), 2001-2016: Chief of the Department of Traditional Medicine. April 2006 - May 2010: Vice Dean of Faculty of Medicine, Pharmacy and Odonto-Stomatology, University of Bamako, Mali. List of publications: google.scholar.

Mamadou Koumaré, Pharmacist, Doctor in pharmaceutical es-science since 1968 (University of Toulouse, France). Education: Professor of pharmacognosy since 1975 (National School of Medicine and Pharmacy, Mali). Honoris Causa Doctor of International Alternative Medicine in 1983 (University of Srilanka). President of Malian and African Societies of Phythotherapy. List of publications: google.scholar.



32 www.ijntr.org