

# Effect of Endophytic and Plant Growth Promoting Rhizobacteria against Foot Rot Disease of *Piper nigrum* L.

Shobha M.S.<sup>1</sup>, Mahadeva Murthy S.<sup>2\*</sup>

<sup>1</sup>Department of Microbiology, Maharani's Science College for Women, Mysuru-570005, India

Email:shobamanoj2014@gmail.com

<sup>2</sup>Department of Microbiology, Yuvaraja College (Autonomous), University of Mysore, Mysuru- 570005, India

Email:smmurthy2025@gmail.com

**Abstract**— Crop loss in black pepper (*Piper nigrum* L.) due to pathogenic diseases is mainly induced by soil borne fungi, bacteria, nematodes and viruses. Foot rot disease caused by *Phytophthora capsici* Leonian is a major production constraint in South India and other south East Asian countries. Combination of biocontrol agents that are compatible with each other is one of the emerging strategies to control plant disease and pest. The present study was designed to evaluate the protective effects of compatible endophytic fungal (*Trichoderma harzianum* Th16 and Th5) and rhizobacterial (*Pseudomonas fluorescens* Pfl) strains against pepper foot rot disease. Our results showed that *T. harzianum* (Th16 and Th5) and *P. fluorescens* (Pfl) were compatible and effectively inhibited the growth of *P. capsici*. The application of endophytic and rhizobacterial strains, alone and in combination in green house and field conditions were found to be effective in controlling the foot rot of pepper caused by *P. capsici* by inducing systemic resistance (ISR) as evidenced by enhanced activities of PO, PPO, PAL,  $\beta$ -1,3-glucanase, chitinase and total phenolics involved in the synthesis of phytoalexins thereby promoting the growth of plants. However, combinations of Th16 + Th5 + Pfl were more effective than individual treatments. The findings suggest that synergistic interactions of biocontrol agents may be responsible for the management of foot rot of pepper caused by *P. capsici*.

**Keywords**— *Azadiracta indica*, Endophytes, Foot rot of pepper, *Phytophthora capsici* Leonian, PGPR and PGPE bio-formulations.

## I. INTRODUCTION

Black pepper (*Piper nigrum* L.) is a commercial spice crop cultivated in India. India ranks first in the world in terms of production, consumption and exports and the crop is grown in an area of 131230 ha with an annual production of 55500 tonnes (Indian Spice Board,

India, Feb 2017). Foot rot disease of black pepper caused by *Phytophthora capsici* is known to affect 90% of yield in India (Nair and Gupta, 2003; Krishnamoorthy and Parthasarathy 2011). The control of foot rot disease has been almost exclusively based on the application of chemical pesticides that effectively kill the *P. capsici*. Although several effective pesticides have been recommended for use against this pathogen, they are not considered to be long-term solutions due to concerns of expense, exposure risks, fungicide residues, toxicity to non-target organisms and other health and environmental hazards. Therefore, recent efforts have been focused on developing eco-friendly safe, long lasting and effective management strategies against plant pathogens.

Use of biocontrol agents has been shown to be eco-friendly and effective against many plant pathogens and pest. Several biocontrol agents have been documented to prevent foot rot disease by inducing systemic resistance of pepper plants against *P. capsici*. Induced systemic resistance (ISR) activates multiple defense mechanisms that include increased activity of pathogenesis related (PR) proteins like Chitinase,  $\beta$ -1,3-glucanase and peroxidase (PO) (Maurhofer *et al.*, 1994; Xue *et al.*, 1998), and also the accumulation of low molecular weight substances called phytoalexins (Van Peer and Schippers, 1992). Chitinases and  $\beta$ -1,3-glucanases are a structurally and functionally diverse group of hydrolytic enzymes involved in defense reactions of plants against pathogens (Jackson and Taylor, 1996), while PO and PAL are the key enzymes involved in phenylpropanoid metabolism (Vidhyasekaran *et al.*, 1997).

Accumulating evidence suggest that the organisms under most scrutiny for potential use in biological control of pest and diseases are bacteria belonging to the genera *Pseudomonas* and *Bacillus* (Ramamoorthy *et al.*, 2001). Further, plant growth promoting endophytic bacteria (PGPE), especially *Bacillus subtilis* (EPCO16 and EPC5) and plant growth

promoting rhizobacteria (PGPR), especially *Pseudomonas fluorescens* (Pf1) strains have been developed commercially as a talc based formulation and tested against several crop diseases (Vivekananthan *et al.*, 2004; Rajendran *et al.*, 2007; Kavino *et al.*, 2007 and Harish *et al.*, 2008). Sundaramoorthy *et al.*, 2011 reported that combination of *P. fluorescens* strains and *B. subtilis* strain together resulted in significant growth promotion that was correlated with induced resistance in *Capsicum annum* L. Several approaches have been made to manage the foot rot of pepper. However, no attempts have been made for the management of *P. capsici* disease using the mixtures of both PGPR and endophytes. Therefore, the present study was designed to evaluate protective effect of endophytic fungal strains, *T. harzianum* (Th16 and Th5) and rhizobacterial strain *P. fluorescens* (Pf1) against foot rot disease.

## II. MATERIAL AND METHODS

### 2.1. Plant materials and pathogen

The black pepper samples were obtained from Pepper Research Center, Appangala, Mercara, Karnataka. The pathogen was isolated from the black pepper foot showing typical symptoms of *P. capsici* by using Oat meal agar (OMA) medium and the fungal culture was identified based on morphological, conidial and culture characterization using standard fungal manual was send to Department of Studies in Biotechnology, University of Mysore, Mysuru– India for identification purpose and it was identified as *P. capsici*.

### 2.2. Biocontrol agents

The two endophytic fungal strains of *T. harzianum* namely Th16 and Th15 were isolated from bark of *Azadiracta indica*. In addition, antagonistic strains of *P. fluorescens* Pf1 were collected from Department of Studies in Biotechnology, University of Mysore, Mysuru. Pure strains of endophytic *T. harzianum* strains (Th16 and Th15) were maintained on potato dextrose agar (PDA) slants and *P. fluorescens* strain (Pf1) was maintained on King's B (KB) agar slants at 4 °C.

### 2.3. Efficacy of individual and mixtures of biocontrol agents on radial growth of *P. capsici*

*Pseudomonas fluorescens* strain (Pf1) and *T. harzianum* (Th16 and ThC5) strains were tested individually and in combination against *P. capsici* by dual culture technique (Dennis and Webster, 1971). The mycelial disc (9 mm) from 7 days old culture of *P. capsici* was placed in one side of the Petri plate containing 15 ml of PDA medium. After three days of pathogen inoculation, 72 h of old bacterial strains *P. fluorescens* and *T. harzianum* strains were streaked on the opposite of the petri plate by the help of sterilized inoculation needle individually on each plate. Three replications were

maintained for each treatment. The plates were incubated at room temperature (28 ± 2 °C) for three days and inhibition zone was measured. The radial growth of the pathogen and per cent reduction over control was calculated by using the following formula

$$\text{Per cent reduction over control} = \frac{C - T}{C} \times 100$$

Where, C – Mycelial growth of the pathogen in control (mm) and T – Mycelial growth of the pathogen in dual plate (mm).

### 2.5. Preparation of individual and mixtures of PGPR and PGPE bio-formulations

For individual strains of *P. fluorescens* (Pf1) and *T. harzianum* (Th16 and Th5) were inoculated into the sterilized KB and Potato dextrose broth, respectively and incubated in a rotary shaker at 150 rpm for 48 h at room temperature (28 ± 2 °C). After 48 h and 7 days of incubation, the broth containing  $9 \times 10^8$  cfu/ml was used for the preparation of talc-based formulation for bacterial culture. To the 400 ml of bacterial suspension, 1 kg of the talc powder (sterilized at 105 °C for 12 h), calcium carbonate 15 g (to adjust the pH to neutral) and Carboxymethyl cellulose (CMC) 10 g (adhesive) were mixed under sterile conditions, following the method described by (Nandakumar *et al.*, 2001). After shade drying overnight, it was packed in polypropylene bag and sealed.

### 2.6. Greenhouse studies

#### 2.6.1. Effects of bio-formulation mixtures on the incidence of foot rot disease

To study the induced systemic resistance (ISR) against *P. capsici* of pepper veins, a pot culture experiment was conducted with single and combination of rhizosphere bacteria and endophytic fungus. Veins are grown in earthen pots (Size-0.35 m diameter, 0.50 m height, volume of soil: 0.04 m<sup>3</sup>) filled with sterilized potting soil containing the spore suspension ( $2 \times 10^5$  spores/g of soil) of *P. capsici*.

In all treatments, the talc-based bio-formulation mixture was applied as seedling root dip and soil application. The fungicide **RIDOMIL GOLD® MZ** was used as a positive control. For treatment, the pepper vine cv. Subhakara, hence the cultivar is susceptible to foot rot disease the same cultivar was used for further studies. After 25 days, the seedlings were pulled out from the pots and transplanted at the rate of five vine per pot (Size-0.35 m diameter, 0.50 m height, volume of soil: 0.04 m<sup>3</sup>) containing sterilized soil mixture (cow dung: sand: soil in 1:1:1 ratio) inoculated with the spore suspension ( $2 \times 10^5$  spores/g of soil mixture) of *P. capsici* (Sundaramoorthy *et al.*, 2012). For root dipping, pepper veins were dipped in 250 ml of *Trichoderma* suspension ( $9 \times 10^8$  cfu/ml) for 2 h, ensuring that roots alone were immersed in the

inoculum and planted in pots. In soil application, 25 ml of *Trichoderma* suspension ( $9 \times 10^8$  cfu/ml) per pot was poured 30 and 60 days after planting (40–45 cm height). Vine treated with **RIDOMIL GOLD® MZ** (2 g/litre) as well as soil drench (0.1%) at 30 and 60 days after planting served as a positive control.

The pepper vine inoculated with the pathogen alone served as inoculated control. The observation on development of *P. capsici* symptoms was recorded at the time of harvest. Each treatment was replicated thrice in Completely Randomized Block Design (CRD). The percent disease index (PDI) was estimated using the formula suggested by McKinney (1923).

#### **Induction of defense-related protein and experimental design**

*Pseudomonas fluorescens* and endophytic (*T. harzianum*) fungal strains in single and in combinations (Treatments T1-Th16, T2-Th15, T3-Pf1, T4-Th16 + Th15, T5-Th15 + Pf1, T6-Pf1 + Th16, T7-Th16 + Th5 + Pf1, T8-Ridomil gold, T9-inoculated control, T10-healthy control) were used in the induction of defense reactions in pepper. The bio-formulations treated vine were sown at the rate of one veins per pot (Size-0.35 m diameter, 0.50 m height, volume of soil: 0.04 m<sup>3</sup>) filled with sterilized potting soil containing the spore suspension ( $2 \times 10^5$  spores/g of soil) of *P. capsici*. Bio-formulation treated plants were challenge inoculated with *P. capsici*, while the other set were not challenge inoculated. The plants neither treated with bio-formulation nor challenged by the pathogen were kept as a control. Three replications were maintained in each treatment. Each replicate consisted of five pots and in each pot one vine were maintained. The experiments were conducted using CRD in greenhouse. The humidity in the greenhouse was maintained at around RH 70%. The temperature was adjusted to 26 °C (day)/20 °C (night).

#### **2.6.3. Sample collection and assay of defense-related proteins**

Vine leaf tissues were collected at different time intervals (0, 4, 8, 16, 24, 48 and 72 hours after pathogen inoculation). Four pepper vines were sampled from each replication of the treatment separately and were maintained for biochemical analysis. Leaf samples were homogenized with liquid nitrogen in a pre-chilled mortar and pestle. One gram of leaf sample was homogenized with 2 ml of 0.1 M sodium phosphate buffer (pH 7.0) at 4 °C. The homogenate was centrifuged for 20 min at 10,000 rpm. The supernatant was used as a crude enzyme extract for assaying PO (Hammerschmidt *et al.*, 1982), polyphenol oxidase (PPO) (Mayer *et al.* 1965) and PAL (Dickerson *et al.*, 1984). Enzyme extracted in 0.1 M sodium citrate buffer (pH 5.0) was used for the estimation of chitinase (Boller and Mauch 1988) and  $\beta$ -1,3-glucanase

(Pan *et al.*, 1991). The total phenol content was estimated as per the procedure given by (Zieslin and Ben-Zaken 1993).

#### **2.6.4. Native polyacrylamide gel electrophoresis analysis**

The isoform profiles of PO and PPO were studied by discontinuous Native polyacrylamide gel electrophoresis analysis (PAGE) (Laemmli 1970). The protein extract was prepared by homogenizing 1 g of leaf sample in 2 ml of 0.1 M sodium phosphate buffer (pH 7.0) and centrifuged at 18,000 rpm for 20 min at 4 °C. The protein content of the sample was determined (Bradford 1976) and samples (50  $\mu$ g protein) were loaded into 8% polyacrylamide gels (Sigma, USA). After electrophoresis, PO isoforms were visualized by soaking the gels in staining solution containing 0.05% benzidine (Sigma–Aldrich, Mumbai, India) and 0.03% H<sub>2</sub>O<sub>2</sub> in acetate buffer (20 mM, pH 4.2) (Nadolny and Sequeira 1980). For assessing the PPO isoform profiles, the gels were equilibrated for 30 min in 0.1% *p*-phenylene diamine, followed by the addition of 10 mM catechol in the same buffer (Jayaraman *et al.*, 1987).

#### **2.7. Statistical analysis**

The data on effect of the treatments on the growth of pathogens, severity of diseases and activity of enzymes in pepper vine were analyzed by analysis of variance (ANOVA), and treatment means were compared by Duncan's Multiple Range Test (DMRT). The data on disease severity was arcsine transformed before undergoing statistical analysis (Gomez and Gomez 1984).

### **III. RESULTS**

#### **3.1. Compatibility among bacterial strains**

PGPR strain of *P. fluorescens* (Pf1) and PGPE strains of *T. harzianum* (Th16 and Th15) were tested for their compatibility *in vitro*. None of the antagonistic bacteria were inhibited by each other, So the absence of inhibition zone suggesting that these three biocontrol agents were compatible with each other.

#### **3.2. Effect of biocontrol agents on radial growth of *P. capsici***

One strain of *P. fluorescens* (Pf1) and two strains of *T. harzianum* (Th16 and Th15) were tested individually and in combination to assess the effect of biocontrol agents on radial growth of *P. capsici*. All the treatments were effective in reducing the mycelial growth of the pathogen. However, the combined application of Th16 + Th15 + Pf1, Th16 + Th15 and Th16 + Pf1 had resulted in the least mycelial growth with 49.0, 54.0 and 54.1 mm, respectively. Combined application of Th16 + Th15 + Pf1 and Th16 + Th15 and Th16 + Pf1 recorded the maximum inhibition zone of 26.0, 23.5 and 19.5 mm, respectively. The control plates recorded the highest mycelial growth of 90.00 mm (Table 1).

Table.1: Effect of biocontrol agents on the mycelial growth of *P. capsici*

Sl. No	Treatments	Mycelial growth (mm)	Inhibition zone (mm)	Percent inhibition over control (mm)
1	Th16	61.00 <sup>de</sup>	10.00 <sup>f</sup>	32.22 <sup>de</sup>
2	Th5	63.50 <sup>ef</sup>	8.70 <sup>g</sup>	29.44 <sup>ef</sup>
3	Pf1	70.40 <sup>e</sup>	7.00 <sup>h</sup>	21.77 <sup>g</sup>
4	Th16 + Th5	54.00 <sup>b</sup>	23.50 <sup>b</sup>	40.00 <sup>b</sup>
5	Th5 + Th5	56.20 <sup>bc</sup>	15.20 <sup>d</sup>	37.55 <sup>bc</sup>
6	Pf1 + Th16	54.10 <sup>b</sup>	19.50 <sup>c</sup>	39.89 <sup>b</sup>
7	Th16 + Th5 + Pf1	49.00 <sup>a</sup>	26.00 <sup>a</sup>	45.55 <sup>a</sup>
8	<b>RIDOMIL GOLD</b>	58.00 <sup>bcd</sup>	13.10 <sup>e</sup>	35.55 <sup>cd</sup>
9	Control	90.00 <sup>h</sup>	0.00	0.00

Values are mean of three replications. In a column, mean followed by a common letter (s) are not significantly different at the 5% level by DMRT.

### 3.3. Efficacy of PGPR and PGPE strains on foot rot incidence under greenhouse conditions

Talc-based bio-formulation of *P. fluorescens* (Pf1) and *T. harzianum* (Th16 and Th15) strains individually or in combination were tested for their efficacy against *P. capsici* under pot culture conditions, along with **RIDOMIL GOLD** as a chemical check. Both individual and strain mixtures significantly reduced the foot rot incidence (by 21–35%) compared to untreated

plants upon challenge inoculation of *P. capsici* (Fig. 1). Conspicuously, a combination of Th16 + Th15 + Pf1 together resulted in a significantly lower foot rot disease index (PDI) than any of the strains individually, as well as better germination (96%) and plant height (73.62 cm) (Fig. 2). The results indicated that disease reduction by a combination of antagonistic bacterial and endophytic strains was comparable with the fungicide control, which recorded the PDI of 22% and 18%, respectively.




Fig.1: Efficacy of individual and mixture of biocontrol agents on foot rot incidence in black pepper under greenhouse conditions. Values are mean of three replications. The line on each bar represents  $\pm$ SE and the values in the bars are not significantly different at the 5% level according to DMRT.




Fig.2: Effect of individual and mixture of biocontrol agents on growth promoting activity in black pepper under greenhouse condition. Values are mean of three replications. The line on each bar represents  $\pm$ SE and the values in the bars followed by the common letter(s) are not significantly different at the 5% level according to DMRT.

### 3.4. Activity of defense enzymes and pathogenesis-related (PR) proteins

Induction of defense enzymes and pathogenesis-related (PR) proteins was studied in the rhizobacteria and endophytes treated pepper veins. The results revealed higher expression of defense-related proteins upon challenge inoculation with *P. capsici*. *P. fluorescens* and *T. harzianum* strains individually and in a mixture differ in the ability to stimulate PO and PPO in pepper veins challenge inoculated with *P. capsici*. The results of the study revealed that there was an increase PO and PPO

activities in Th16 + Th15 + Pfl mixtures treated plants compared to untreated control plants upon challenge inoculation. The activities (Th16 + Th15 + Pfl) were found to increase 110 units of protein at 48 hours after inoculation, compared to all other treatments with drastic reduction in enzyme activities in later time points. The pathogen inoculation in untreated control plants also stimulated the enzymes but the level was less than that of mixture of *P. fluorescens* and *T. harzianum* pretreated plants (Fig. 3 and Fig. 4).




Fig.3: Induction of peroxidase activity in pepper veins treated with biocontrol agents against *P. capsici*. The vertical bar indicate the SE at 5% level according to DMRT. Treatments T1-Th16, T2-Th15, T3-Pfl, T4-Th16 + Th15, T5-Th15 + Pfl, T6-Pfl + Th16, T7-Th16 + Th5 + Pfl, T8-Ridomil gold, T9-inoculated control, T10-healthy control




Fig.4:

Induction of polyphenol oxidase activity in pepper veins treated with biocontrol agents against *P. capsici*. The vertical bar indicate the SE at 5% level according to DMRT. Treatments T1-Th16, T2-Th15, T3-Pfl, T4-Th16 + Th15, T5-Th15 + Pfl, T6-Pfl + Th16, T7-Th16 + Th15 + Pfl, T8- Ridomil gold, T9-inoculated control, T10-healthy control

PAL activity was significantly higher in Th16 + Th15 + Pfl mixture treated pepper veins inoculated with *P. capsici* than in untreated controls. PAL accumulation reached a maximum of 3.5 units of protein at 24 hours after inoculation in Th16 + Th15 + Pfl treatment group

when compared to all other treatment groups. Untreated pepper veins inoculated with *P. capsici* recorded the least induction of 1.4 units of protein and showed drastic decline at 48 hours after inoculation (Fig. 5).




Fig.5: Induction of phenylalanine ammonia-lyase activity in pepper veins treated with biocontrol agents against *P. capsici*, the vertical bar indicate the SE at 5% level according to DMRT. Treatments T1-Th16, T2-Th15, T3-Pfl, T4-Th16 + Th15, T5-Th15 + Pfl, T6-Pfl + Th16, T7-Th16 + Th15 + Pfl, T8- Ridomil gold, T9-inoculated control, T10-healthy control






Fig.6: Induction of  $\beta$ -1,3-glucanase activity in pepper veins treated with biocontrol agents and endophytes against *P. capsici*, the vertical bar indicate the SE at 5% level according to DMRT. Treatments T1-Th16, T2-Th15, T3-Pfl, T4-Th16 + Th15, T5-Th15 + Pfl, T6-Pfl + Th16, T7-Th16 + Th5 + Pfl, T8- Ridomil gold, T9-inoculated control, T10-healthy control

1,3-glucanase (Th16 + Th15 + Pfl) mixture treated plants inoculated with *P. capsici*. Showed an enhanced activity of 1.7 units of protein at 16 hours after inoculation when compared to untreated control plants which showed 0.8 units of protein. The activities of  $\beta$ -1,3-glucanase increased in plants treated with bio-formulation mixture (Th16 + Th15 + Pfl) up to 3 days after *P. capsici* inoculation, and declined thereafter as shown in (Fig. 6). Maximum accumulation of total phenol content was also

observed in the combination treatments of Th16 + Th5 + Pfl followed by Th16 + Th15 and Th16 + Pfl (Fig. 7). Untreated control, or inoculated with the pathogen alone did not show any remarkable change in the activity of phenolic substances. These results clearly indicate that the defense enzymes and PR proteins were more activated in the mixtures of bacterial bio-formulations treated plants thus strengthening the pepper veins against the abiotic stress.




Fig.7: Induction of chitinase activity in pepper veins treated with biocontrol agents and endophytes against *P. capsici*, the vertical bar indicate the SE at 5% level according to DMRT. Treatments T1-Th16, T2-Th15, T3-Pfl, T4-Th16 + Th15, T5-Th15 + Pfl, T6-Pfl + Th16, T7-Th16 + Th5 + Pfl, T8- Ridomil gold, T9-inoculated control, T10-healthy control

The chitinase activity was found high in Th16 + Th5 + Pfl and Ridomil treated (T8) veins showed 2.5 and 2.0 at 16 hpi and 24 hpi. The activities of chitinase

activity increased in plants treated with bio-formulation mixture (Th16 + Th15 + Pfl) up to 3 days after *P. capsici* inoculation, and declined thereafter as shown in (Fig. 7).

Untreated control, or inoculated with the pathogen alone did not show any remarkable change in the activity as shown in the figure.

#### IV. DISCUSSION

Accumulating evidence from literature has shown that compatible multiple strains appear to be an important pre-requisite for the desired effectiveness of strains and more consistent disease suppression (Young Cheol *et al.*, 2008; Latha *et al.*, 2009). The results of the present study provide evidence that the compatibility of *P. fluorescens* (Pf1) and *T. harzianum* (Th16 and Th15) effectively inhibited the growth of *P. capsici*. Several studies have documented that *Pseudomonas*, *B. subtilis* (CA32) and *Trichoderma harzianum* (RU01) significantly reduced the mycelial growth and conidial production of *P. capsici* (Abeysingne 2007) by producing the wide array of antibiotics such as 2,4-diacetylphloroglucinol, oligomycin, phenazine, pyoleteorin, pyrrolnitrin, pyocyanin, Iturin, bacillomycin, zwittermycin-A and surfactin responsible for antifungal activities (Yu *et al.*, 2002).

Inhibitory effect of biocontrol agents observed in dual culture technique is correlated with the management of wilt disease of foot rot caused by *P. capsici* under green house as well as field conditions. The results of the present study revealed that the talc based bio-formulations prepared by using individual and in combination of Th16 + Th15 + Pf1 strains significantly reduced the wilt disease incidence in pepper. This effect may be due to its stimulatory effects on ISR as evidenced by increased activities of defense enzymes such as PO, PPO and PAL involved in the synthesis of phytoalexins thereby induce systemic resistance against foot rot disease and promote the growth of plants. Our results substantiate the inhibition of various plant pathogen and disease management by using several biocontrol agents through the induction of ISR in plants reported by (Van Peer *et al.*, 1991; Kloepper, 1993; Van Loon, 1997; Chen *et al.*, 2000). Thus, our findings provide evidence that the induction of defense enzymes and PR proteins by application of endophytic fungi and rhizosphere bacterial strains may strengthen the plants against various biotic stresses.

#### V. CONCLUSION

The present study demonstrated that the combinations of endophytic fungi Th16 + Th15 + Pf1 bacterial strains consistently reduced the radial mycelial growth of *P. capsici* by producing various antibiotics and reduced the foot rot of pepper under green house and field conditions by inducing ISR compared to individual agents. Combination of biocontrol agents is a strategic approach to control the plant disease and pest (Nandakumar *et al.*, 2001a and Saravanakumar *et al.*,

2007). Furthermore, interactions among the bacterial strains may have synergistic effects that could induce ISR and promote the growth of the plants (Sundaramoorthy *et al.*, 2007). Several literature have documented that the use of biocontrol agents in combination was more effective for management of plant diseases and pathogens compared to individual agents (Sivakumar, 2012).

#### REFERENCES

- [1] Abeysingne, (2007). Biological control of *Fusarium solani* f. sp. *phaseoli* the causal agent of root rot of bean using *Bacillus subtilis* CA32 and *Trichoderma harzianum* RU01. *Ruhuna Journal of Science*, 2, 82–88.
- [2] Boller T. and Mauch, F. (1988). Colorimetric assay for chitinase. *Methods in Enzymology*, 161, 430–435.
- [3] Chen, C., Belanger, R.R. Benhamou, N. and Paullitz, T.C. (2000). Defense enzymes induced in cucumber roots by treatment with plant-growth promoting rhizobacteria (PGPR). *Physiological and Molecular Plant Pathology*, 56: 13–23.
- [4] Dickerson, D.P. Pascholati, A.E., Hagerman, L.G. Butler and R.L. Nicholson, S.F. (1984). Phenylalanine ammonia-lyase and hydroxy cinnamate CoA ligase in maize mesocotyls inoculated with *Helminthosporium maydis* or *Helminthosporium carbonum*. *Physiology and Plant Pathology*, 25, 111–123.
- [5] Gomez K.A. and Gomez A.A. (1984). *Statistical Procedure for Agricultural Research*, John Wiley and Sons, New York
- [6] Hammerschmidt, R. Nuckles and J. Kuc, E.M. (1982). Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiological Plant Pathology*, 20, 73–82.
- [7] Harish, S. M. Kavino, N. Kumar, D. Saravanakumar, K. Soorianasundaram and Samiyappan, R. (2008). Biohardening with plant growth promoting rhizosphere and endophytic bacteria induces systemic resistance against banana bunchy to virus. *Applied Soil Ecology*, 39, 187–200.
- [8] Indian Spice Board, India, Feb 2017
- [9] Jackson A.O. and Taylor, C.B. (1996). Plant–microbe interactions: life and death at the interface. *Plant Cell*, 8, 1651–1680.
- [10] Jayaraman, K.S. Ramanuja, M.N. Vijayaraghavan P.K. and Vaidyanathan, C.S. (1987). *Oxidative enzyme in pearl millet*. Food Chemistry, Wiley and Sons, New York, 24, 203.
- [11] Kavino, M. S. Harish, N. Kumar, D. Saravanakumar, T. Domodaran, K. Soorianasundaram and Samiyappan, R. (2007). Rhizosphere and endophytic



- bacteria for induction of systemic resistance of banana plantlets against bunchy top virus. *Soil Biology & Biochemistry*, 39, 1087–1098
- [12] Kloepper, J.W. (1993). Plant growth-promoting rhizobacteria as biological control agents, B. Metting, Editor, *Soil Microbial Technologies, Meeting B*, M. Dekker Inc, New York, pp. 255–274.
- [13] Krishnamoorthy, B., and Parthasarathy, V. A. (2011). Improvement of black pepper. *Plant Sciences Reviews 2010*, 37.
- [14] Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680–685.
- [15] Latha, P., Anand, T. N. Ragupathi, V. Prakasam and R. Samiyappan, (2009). Antimicrobial activity of plant extracts and induction of systemic resistance in tomato plants by mixtures of PGPR strains and zimmu leaf extract against *Alternaria solani*. *Biological Control*, 50, pp. 85–93.
- [16] Maurhofer, M. C. Hase, D. Maurwly, J.P. Metraux and Defago, G. (1994). Induction of systemic resistance of tobacco to tobacco necrosis virus by the root colonizing *Pseudomonas fluorescens* strain CHAO: influence of the *gac A* gene and of pyoverdine production. *Phytopathology*, 84, 136–146.
- [17] Mayer, E. Harel and R.B. Shaul, A.M. (1965). Assay of catechol oxidase a critical comparison of methods. *Phytochemistry*, 5, 783–789.
- [18] Mckinney, H.H. (1923). A new system of grading plant diseases. *Journal of Agricultural Research*, 26, 195–218
- [19] Nadolny, L. and Sequeira, I. (1980). Increases in peroxidase activities are not directly involved in induced resistance in tobacco. *Physiology and Plant Pathology*, 16, 1–8.
- [20] Nair, R.R., and Gupta, S.D. (2003). Somatic embryogenesis and plant regeneration in black pepper (*Piper nigrum* L.). Direct somatic embryogenesis from tissue of germinating seeds and ontogeny of somatic embryos. *J. Hortic. Sci. Biotchnol* 78: 416-421.
- [21] Nandakumar, R. Babu, S.R., Viswanathan, J., Sheela, T., Raguchander and Samiyappan, A. (2001). New bio-formulation containing plant growth promoting rhizobacterial mixture for the management of sheath blight and enhanced grain yield in rice. *BioControl*, 46 (4): 493–510.
- [22] Nandakumar, R. S. Babu, R. Viswanathan, J. Sheela, T. Raguchander and R. Samiyappan, A. (2001). A new bio-formulation containing plant growth promoting rhizobacterial mixture for the management of sheath blight and enhanced grain yield in rice. *BioControl*, 46 4, pp. 493–510
- [23] Pan, S.Q., Ye, X.S. and Kuc, J. (1991). Association of  $\beta$ -1,3-glucanase activity and isoform pattern with systemic resistance to blue mold in tobacco induced by stem injection with *Pernospora tabacina* or leaf inoculation with tobacco mosaic virus. *Physiological and Molecular Plant Pathology*, 39, 25–39.
- [24] Rajendran, L. G. Karthikeyan, T. Raguchander and Samiyappan, R. (2007). *In vitro* evaluation of bacterial endophytes influence on *Ganoderma lucidum* (leys) Karst. mycelial growth.. *Journal of Plant Protection Research*, 47 (4), 425–436.
- [25] Ramamoorthy, V. R., Viswanathan, T. Raguchander, Prakasam V. (2001). Induction of systemic resistance by plant growth promoting rhizobacteria in crop plants against pest and diseases. *Crop Protection*, 20, 1–11.
- [26] Saravanakumar, C. Vijayakumar, N. Kumar and R. Samiyappan, (2007). PGPR-induced defense responses in the tea plant against blister blight disease. *Crop Protection*, 26, 556–565.
- [27] Sivakumar, G. (2012). Management of Phytophthora foot rot of black pepper with potassium phosphonate (Akomin) and *Trichoderma harzianum*. *Journal of Mycology and Plant Pathology*, 42, 372-375.
- [28] Sundaramoorthy, S., Raguchander, T., Ragupathi, N., and Samiyappan, R. (2012). Combinatorial effect of endophytic and plant growth promoting rhizobacteria against wilt disease of *Capsicum annum* L. caused by *Fusarium solani*. *Biological Control*, 60(1), 59-67.
- [29] Van Loon L.C. (1997). Induced resistance in plants and the role of pathogenesis-related proteins. *European Journal of Plant Pathology*, 103, 753–765.
- [30] Van Peer, R. and Schippers, B. (1992). Lipopolysaccharides of plant growth promoting *Pseudomonas* spp. strain WCS417r induce resistance in carnation to *Fusarium* wilt. Netherland. *Journal of Plant Pathology*, 98, 129–139.
- [31] Van Peer, R., Neimann, G.J. and Schippers, B. (1991) Induced resistance and phytoalexin accumulation in biological control of *Fusarium* wilt of carnation by *Pseudomonas* sp. strain WCS417r. *Phytopathology*, 81, 728–734.
- [32] Vidhyasekaran, P. Ponmalar, T.R. Samiyappan, R. R. Velazhahan, R. Vimala and A. Ramanathan, A. (1997). Host specific toxin production by *Rhizoctonia solani*, the rice sheath blight pathogen. *Phytopathology*, 87, 1258–1263. |
- [33] Vivekananthan, R. M., Ravi, A. Ramanathan and Samiyappan, R. (2004). Lytic enzymes induced by *Pseudomonas fluorescens* and other biocontrol organisms mediate defence against the anthracnose

- 
- pathogen in mango. *World Journal of Microbiology and Biotechnology*, 20, 235–244
- [34] Xue, L., Charest, P.M. and Jabaji, S.H. (1998). Systemic induction of peroxidases,  $\beta$ -1,3-glucanases, chitinases and resistance in bean plants by binucleate *Rhizoctonia* species. *Phytopathology*, 88, 359–365.
- [35] Young Cheol, K.J. Hyunchae, K. Kil Yong and Seur Kee, P. (2008). An effective biocontrol bioformulation against *Phytophthora* blight of pepper using growth mixtures of combined chitinolytic bacteria under different field conditions. *European Journal of Plant Pathology*, **120**, 373–382
- [36] Yu, G.Y., Sinclair, J.B., Hartman, G.L. and Bertagnolli, B.L. (2002). Production of iturin A by *Bacillus amyloliquefaciens* suppressing *Rhizoctoniasolani*. *Soil Biology & Biochemistry*, 34, 955–963.
- [37] Zieslin, N., and Ben-Zaken, R. (1993). Peroxidase activity and presence of phenolic substances in peduncles of rose flowers. *Plant Physiology and Biochemistry*, 31(3): 333-339.