

Evaluation of safety of *Azadirachta indica* seed oil on Albino rat through haematological and some antioxidants by the rotatable central composite design (RCCD) of the response surface methodology (RSM)

Ojimekwe Phillippa², Udofia Patrick G^{1*}, Anthony Ukom²

²Department of Food Science and Technology, Michael Okpara University of Agriculture, Umudike, P. M. B. 7267, Abia State, Nigeria (philippaco60@yahoo.com).

²Department of Food Science and Technology, Michael Okpara University of Agriculture, Umudike, P. M. B. 7267, Umuahia. Abia State, Nigeria (tony2008gospel@gmail.com)

¹Department of Food Technology, Akwa Ibom State Polytechnic, Ikot Osurua, P. M. B. 1200, Ikot Ekpene, Nigeria.
kesitpatrick1@gmail.com

*Corresponding author: email: kesitpatrick1@gmail.com

Abstract— Raw *Azadirachta indica* seed oil is gradually adopted by local farmers as a protector of stored grains and legumes against the common insects, but with some reservations on its safety. The study investigated the safety of *A. indica* on mammalian subjects using haematological and antioxidant indices. *A. indica* seed was obtained from the trees in the same location in Utu Ikot Ukpang, Essien Udim Local Government Area, Nigeria. The Albino rats were obtained from the Animal House of Biochemistry Department of the University of Calabar, Nigeria. The animals were treated with varied *A. indica* seed oil at varied concentrations, periods of exposure and age. Haematological analyses revealed that models of packed cell volume (PCV), red blood cell (RBC), white cell count (WCC) and hemoglobin content (Hb) were not significant. Catalase (CAT), superoxide dismutase (SOD), and glutathione (GSH) were also not significant ($p > 0.05$). Linearity coefficients of the models however appeared to be significant ($R^2 \leq 0.8000$). The mathematical and pictorial models showed slight influence of the test substance on the parameters. The investigation revealed that *A. indica* seed oil showed no adverse effect on the haematological and antioxidant profile of the animal models at the level of administration, the slight statistical significance could be attributed to other experimental values. *A. indica* oil seems to be safe on mammalian subjects. More work is recommended on the topic using higher levels of the plant materials at a longer period of exposure.

Keywords— *Azadirachta indica*, response surface methodology, rotatable central composite design, haematological indices, enzymes.

I. INTRODUCTION

Insecticides are organic and inorganic substances which are used to kill or inactivate insects to check damaging effects of their activities, application is useful in agriculture, medicine, industry and in the homes (Coats, 1999). Deployment of synthetic chemical insecticides in agriculture is claimed to be a major factor behind the increase in the 21st-Century's agricultural productivity (Gerald, 2006). Long-term applications of synthetic insecticides have resulted in residues accumulating in different environmental components of water, food, air and soil (Lorendo-Pino *et al.*, 2014). In order to circumvent their ill effects, ideal insecticides which are not toxic and more environmentally friendly are sought from the plant kingdom (Ojimekwe, 2008; Lorendo-Pino *et al.*, 2014).

Botanical insecticides in integrated insect management program have greatly reduced environmental pollution, toxicity to users, reduce cost of medical treatment of insecticide related diseases. Safety to farmers and consumers of treated foods has also been demonstrated. Efforts to develop and apply botanicals as alternatives to the synthetic chemical insecticides have been reported by many researchers, for instance, (Ojimekwe and Adler, (1999) reported on the potential of zimaldehyde, 4-Allyl-anisol, linalool, terpinol in the control of confused flour beetle, *Tribolium confusum*. Okpereke and Bunmi, (2006), Okpereke, (2007) reported effectiveness of *Xylopiya aethopia* products against the common maize weevil, Huang and Ho, (1998) demonstrated effectiveness of *Piper guneense* powder against *Tribolium castaneum* (Herbst), *Sitophilus*

zeamais Motsch. Mikhaiel, (2011) showed that some volatile oils of plants could protect packages of irradiated wheat flour against *Ephestia kuheniella* and *Tribolium castaneum*. Insecticidal activities of aromatic plant extracts and essential oils against *Sitophilus oryzae* and *Callosobruchus chinensis* had been demonstrated by (El-Wakil, 2013). Commercial *neem* seed oil is currently in use in the integrated insect management programme (Chaudhary, 2017). *A. indica* is native to Asia, but has now naturalized in Nigeria. *A. indica* is an important item in the traditional medicinal practice in the area of study, according to Pallav *et al.*, (2014) every part of the plant is claimed to possess some medicinal and insecticidal properties, but the seed is grossly underutilized (Schmutterer, 1990). Igbokwe (2017) opined that it is the presence of some biologically active principles like azdirachtin, meliacin, gedunin, salanin, nimbin, valasin and their derivatives that confers the anti-inflammatory, anti-hyper glycaemic, anti-ulcers, anti-malarial, anti-fungal, anti-bacterial, anti-oxidant, anti-mutagenic, and immunomodulatory properties of *neem* seed oil in mammalian models including anti-feedant, repellent, effect in insects (Ndodo, 2013).

Oil of *A. indica* seed is sprayed on grains/legumes for protection against the damaging effect of the common insects. Protected products are either used for food or feed or replanting. Oily nature of *A. indica* seed oil prevents its total removal from the stored product and therefore may contain residue at levels high enough to elicit health problems in susceptible individuals of the population, the oil may become rancid and develop mycotoxin (Yin *et al.*, 2003), which is a known carcinogen especially at high level of accumulation. As a new product, *A. indica* seed oil deserves safety evaluation on mammalian models before injected into the market for human consumption.

In this study, the Rotatable Central Composite Design (RCCD) of the Response Surface Methodology (RSM) was employed to test the linear, combined and quadratic effects of graded levels of age, dose and exposure time on haematological indices and antioxidants of blood of Albino rats. Response surface methodology tests effect of many variables on parameters of interest at the same time, therefore the method is robust, reliable and predictable, the design saves time and cost (Kwak *et al.*, 2015). According to Kanu *et al.*, (2016), chronic diseases affect blood cells adversely, therefore haematological data provides the most important information in the determination of biochemical and physiological state of animal models (Jorum *et al.*, 2016). Damage mediated by free radicals results in high rate of disruption of membrane fluidity, protein denaturation, lipid peroxidation, oxidative deoxyribonucleic acid (DNA) and alteration of platelet functions, the molecular damages

have been linked with diabetes, cancers, inflammation, aging and atherosclerosis (Kanu *et al.*, 2016).

From literature, many works have been done on the safety or toxicity of *A. indica* seed oil products but to the best of our knowledge, few data is available on the effect of *A. indica* seed oil on haematological and antioxidant levels using the rotatable central composite design of the response surface methodology.

Therefore the work evaluated the safety of *A. indica* seed oil through levels of haematological and enzyme indices of Albino rats using the RCCD of the RSM. Data from the work is expected to enhance food security and low cost of insect pest management, cultivation and utilization of *neem* seed can provide employment.

II. MATERIALS AND METHODS

Forty-five (45) Albino rats of 14 - 40 days were obtained from the animal house of the Department of Biochemistry, University of Calabar, Cross River State, Nigeria. The commercial rodent meal was bought from Pfizer, Lagos. *A. indica* seed was obtained from Essien Udim Local Government Area in Akwa Ibom State. All reagents used for the study were of analar grade, double distilled water was used for their preparation.

Preparation of *A. indica* seed powder

Preparation of *A. indica* seed was sorted, decorticated, washed and dried in the sun for 24 hours. It was then dried in hot air oven at 50°C for 24 hours to a moisture content of about 12%. The dried seed was milled to pass through 0.2-0.5 mm sieve (Fritsch GMBH, Germany BRD-6580), packaged and stored in moisture proof container for subsequent use.

Preparation of *A. indica* seed oil

Preparation of *A. indica* seed oil was carried out according to the method adopted by Okigbo, (2008). Five hundred (500) grams of *A. indica* seed powder was soaked in 1000ml of 95% ethanol in a round-bottomed flask and allowed to stand at room temperature for 24 hours. The filtrate was evaporated with a rotary evaporator (Buch Laboratoriums Technik, AG CH - 9230), to free sample from the solvent leaving a dark slurry as the oil.

Animal models

The animal models were handled according to CLSI (2000). They were stabilized for 5 days with water and fed *ad lib*. A dose of ampicillin was administered through the drinking water to check possible infection, they were housed in separate cages, the housing guaranteed adequate lighting, ventilation and standard humidity.

Administration of *A. indica* seed oil

Treatment of Albino rats was done according to the method of (Adewale *et al.*, (2014) with slight

modification to fit the experimental design. Animals with similar blood and antioxidants parameters were randomly assigned to the experimental units; (15 laboratory animal cages) (Table 1, 2), watered with commercial table water (ARSAN Water) and fed standard rodent diet (Pfizer, Lagos) *ad lib* between 0 to 40 days. Graded levels of *A. indica* oil was administered orally at concentrations according to the experimental design in Table 1. At the end of each experimental run, animals in the experimental unit were anesthetized with diethyl ether, blood samples were taken from the heart and preserved with EDTA to prevent clotting. Samples were centrifuged at 3,000 rpm for 10 min to obtain the plasma.

Estimation of haemoglobin and antioxidant assay of blood of treated Albino rats

Pack cell volume (PVC), red blood cell (RBC), white cell count (WCC), haemoglobin content (HB) according to the method of Adebayo *et al.* (2005) using automated haematology analyzer (ERMA) (Model PCE 210 Japan). Catalase was assayed spectrophotometrically according to the method of Aebi *et al.*, (1983), estimation of superoxide dismutase was done according to the method

of Aebi *et al.*, (1983), estimation of reduced glutathione was determined by the method of Kwak *et al.*, (2016).

Background of experimental design and statistical analysis

RSM enables evaluation of effects of multiple independent variables singly or in combination simultaneously on dependent variables (Meyers *et al.*, 2002). Design Expert Version 10 (Statease Inc., Minneapolis, MN, USA) was deployed for the design and analysis of the experimental data. Age of animal model, dose of the test substance, and exposure time of the animal model to the test substance were chosen from experimental experience (Table 1). The Rotatable Central Composite Design assumes equation 1.

$$Y_i = \beta_0 + \sum_{i=1}^3 \beta_i X_i + \sum_{i=1}^3 \beta_{ii} X_i^2 + \sum_{j=i+1}^3 \beta_{ij} X_i X_j + e \dots \quad (1)$$

where Y_i is any response, β_0 = intercept, β_i = first order model coefficients, β_{ii} = quadratic coefficient for the i^{th} variable, β_{ij} = interaction coefficients for the interaction variables i and j , X_i = any independent variable. Second order coefficient was generated by regression analysis with backward elimination. Responses were first fitted for the factors by the coefficients of determination, R^2 .

Table.1: Experimental design and levels of independents variables

Variable	Units	- α	-1	0	1	+ α
Age	Days	8.6152	14	13.5	40	45.3848
Dose	mg/100g	5.8578	10	20	30	34.1421
Time of exposure	Days	-4.1421	0	10	20	24.1421

$\alpha = 1.4142$ for RCCD

III. RESULTS

Levels of the independent variables, age (X_1) 14-40 days, dose (X_2) 10-30 mg, and exposure time (X_3) 0-20 days were assigned to the experiment by experience and done according to RCCD in Table 1. Effect of the variables on packed cell volume (PCV), red blood cell (RBC), white cell count (WCC), haemoglobin content (Hb), catalase (CAT), superoxide dismutase (SOD) and glutathione (GSH) of the blood samples of the animal models are presented in Table 2. From the table, PCV ranged from 20-39%, RBC ranged from 7.92 to 9.92x10⁶/ml, WCC ranged from 7110-8221x10⁶/ml, Hb ranged from 12.01-13.17g/100g and means of 33.38%, 8.86x10⁶/ml, 7745.23

x10⁶/ml, 12.69 g/100ml respectively. CAT, SOD and GSH ranged from 328.9-954.90 IU/gHb, 5478-6917 IU/gHb and 2.35-3.99 IU/gHb and mean of 441 IU/gHb, 6164 IU/gHb, and 3.50 IU/gHb respectively. Standard deviations of the parameters were 4.59%, 0.52%, 454.20%, 12.69%, 166.90%, 416.07% and 0.493% respectively. The low standard deviation of the parameters indicated low variability in the observation as can be seen in Table 2 except SOD and GSH which showed higher values than the control. Run 14 was not included in the calculation due the extraordinarily low values.

Table.2: Experimental runs, results and standard values of the parameters

Run	Age Days	Dose mg/100g	Exposure Days	PCV %	RBC x10 ⁶ /ml	WCC x10 ⁶ /ml	Hb g/100g	CAT IU/gHb	SOD IU/gHb	GSH μ M/gHb
12	27	20	10	39	9.52	7830	13.17	394.1	5981	3.53
5	8	20	10	38	9.21	8001	12.01	593.5	6098	2.35
10	27	24	24	37	9.51	7911	13.21	394.9	6182	3.75
6	45	20	10	35	9.22	7851	12.91	374.7	5519	3.11
2	40	10	20	37	9.11	7907	13.22	392.3	6917	3.99

4	14	10	0	30	9.25	8110	12.10	389.9	6391	3.34
8	27	34	10	32	9.22	7110	13.00	389.6	6551	3.91
13	27	34	10	35	9.17	8113	12.21	954.9	6276	3.56
7	27	6	10	32	8.97	7917	13.11	377.8	6088	3.98
3	14	30	30	20	9.04	8211	12.01	328.9	5617	3.90
14	27	20	34	9	8.90	8221	12.01	328.9	5617	3.21
1	40	30	0	35	7.92	7117	12.22	389.7	6315	2.66
9	27	20	-4	35	8.21	8220	13.00	399.2	6277	3.77
11	27	20	10	34	8.22	7110	13.00	349.9	6616	3.65
15	27	20	10	34	8.22	7110	13.00	398.3	5478	3.55

PVC = packed cell volume, RBC = red blood count, WCC = white cell count, Hb = haemoglobin, CAT = catalase value, GSH = superoxide dismutase, GSH = glutathione level.

Table.3: Analysis of variance (ANOVA) of the experimental data

	PVC	RBC	WCC	Hb	CAT	SOD	GSH
Model	0.5112	0.4634	0.5540	0.1780	N/A	0.3382	0.0050
Linear effect							
Age (X ₁)	0.4299	0.9882	0.8074	0.0684	N/A	0.3060	0.0198
Dose (X ₂)	1.0000	0.7135	0.2253	0.3545	N/A	0.4073	0.7687
Exposure time (X ₃)	0.5919	0.0992	0.6191	0.3691	N/A	0.8621	0.9327
Interaction							
Age x Dose (X ₁ X ₂)	0.6690	0.5341	0.3068	N/A	N/A	0.9725	0.0080
Age x Exposure (X ₁ X ₃)	0.7863	0.2310	0.7143	N/A	N/A	0.0914	0.1968
Dose x Exposure (X ₂ X ₃)	0.1403	0.3674	0.3953	N/A	N/A	0.0898	0.0141
Quadratic							
Age ² (X ₁₁ ²)	0.4649	0.7105	0.8688	N/A	N/A	N/A	0.0011
Dose ² (X ₂₂ ²)	0.1427	0.9798	0.2788	N/A	N/A	N/A	0.0120
Exposure ² (X ₃₃ ²)	0.6318	0.5203	0.5484	N/A	N/A	N/A	0.0755
R²	0.6519	0.6741	0.6317	0.3488	0.0000	0.5030	0.9611
Adjusted R²	0.0254	0.0874	-0.0313	0.1712	0.0000	0.1302	0.8911
Mean	33.38	8.86	7745.23	12.69	441.04	6164	3.50
Lack of Fit	0.6907	0.5066	0.9441	0.6786	0.9996	0.6797	0.9539

Table 2 shows random order of the experimental runs, coefficient of estimate, coefficient of determination, and control values of the parameters. Table 3 shows statistic of analysis of variance (ANOVA) of the experimental data. The model for PCV was not significant ($p > 0.05$). The coefficient of determination, $R^2 = 0.6519$, and the Adjusted coefficient of determination, $Adj. R^2 = 0.0524$ confirmed a weak statistical confidence of the model. Linear, interaction and quadratic terms of age, dose, and

exposure time were not significant ($p > 0.05$). Variation of the parameter is represented by Fig. 1 and equation 1. According to Fig. 1 and Equation 2, the value of the parameters deviated very slightly with age, dose and exposure time. According to the equation, terms of age, dose and exposure time were synergistic, while interactive terms showed antagonistic effects, quadratic terms were very low but positive.




Fig.1: Response surface-contour plot of dose of test substance and age of model animal on pack cell volume of model animal blood

$$PVC = 23.10 - 0.146(X_1) + 1.09(X_2) + 0.70(X_3) - 6.099(X_1 X_2) - 0.03(X_2 X_3) \dots \quad (2)$$

The model of RBC was not significant ($p > 0.05$), the coefficient of determination, $R^2 = 0.6741$ indicated weak confidence on the model. The linear, interaction, and quadratic terms of the model were not significant ($p > 0.05$) except time (X_3) with which appeared to be

significant $p = 0.0992$. The model is represented by Fig. 2 which shows that red blood cells increased with age of the animal model and not as a function of dose and time (not shown here) and was corroborated by Equation 3.




Fig.2: Response surface-contour plot of dose of test substance and age of model animals on red blood cell content

$$RBC = 10.10 - 0.09(X_1) - 0.07(X_2) - 0.07(X_3) \dots \quad (3)$$

The model for WCC was not significant ($p > 0.05$). The coefficient of determination ($R^2 = 0.6317$), the linear, interaction, and quadratic terms of the model were not significant ($p > 0.05$), with mean value of 34.53×10^6 ml. The model is represented by Fig. 3 and Equation 3.

$$WCC = 7088 + 47.44(X_1) + 85.47(X_2) - 60.79(X_3) - 2.55(X_1 X_2) - 0.87(X_1 X_3) + 2.71(X_2 X_3) + 0.15(X_{11}^2) - 1.80(X_{22}^2) + 0.95(X_{33}^2) \dots \quad (4)$$




Fig.3: Response surface-contour plot of dose of test substance and age of model animals on white cell count.

$$\begin{aligned} \text{WCC} = & 70.88 - 64.150(X_1) + 85.47(X_2) \\ & - 60.80(X_3) - 2.55(X_1X_2) - 0.87(X_1X_3) \\ & + 2.71(X_2X_3) + 0.16(X_{11}^2) - 1.80(X_{22}^2) \\ & + 0.96(X_{33}^2) \quad \dots \end{aligned} \quad (4)$$

The model of Hb was not significant ($p > 0.05$), the coefficient of determination ($R^2 = 0.3488$) and Adj. $R^2 = 0.1712$ confirmed weakness of prediction of the model. The linear, interaction, and quadratic terms of the model were not significant ($p > 0.05$), except age which appeared to be significant, $p = 0.0684$. The model for Hb estimation is given in Equation 5. The model of CAT was not significant and could not be calculated, the linear, interaction, and quadratic terms of the model could not be calculated due to low range of the data (N/A), while the mean value was (420.37 IU/gHb). The model is represented by Equation 5. The model for SOD was not significant ($p > 0.05$), the coefficient of determination $R^2 = 0.5030$ confirmed lack of confidence of the model. The linear, interaction and quadratic terms of the model were not significant ($p > 0.05$). The model was represented by Equation 6.

$$\begin{aligned} \text{SOD} = & 6363 - 1.56(X_1) - 56.24 + 65.49(X_2) - 6.661(X_3) \\ & 0.0724(X_1X_2) + 3.9053(X_1X_2) - 5.10(X_1X_3) \end{aligned}$$

The model for GSH was not significant ($p > 0.05$). The coefficient of determination ($R^2 = 0.9611$, Adj. $R^2 = 0.8911$) showed some confidence on the model. The linear, interaction and quadratic terms of age, dose and exposure time were not significant ($p > 0.05$). The model is represented by equation 7 and a response surface plot (not shown).

$$\text{GSH} = 1.54 + 0.205(X_1) - 0.032(X_2) - 0.145(X_3) \quad \dots$$

IV. DISCUSSION

Many people in the local community where the study was undertaken believed that the oil of *A. indica*

seed at the level they apply to stored grains and legumes could retain the oil residues at a levels high enough to induce toxicity in mammalian system especially if consumed at high doses at an extended period of time. It is well known that the oil contains some bioactive components like alkaloids, polyphenols and flavonoids which have convinced proofs of influence on good biochemical processes in mammals. This is why they are used for as 'drugs' and nutrients (Wararut *et al.*, 2012). In spite of proven assurances of safety of some botanical oils in nutrition and health, information on every plant material to be introduced into the food chain must be made available (Ojimelukwe *et al.*, 2008) to the consumer population. Fear of toxicity of *A. indica* seed oil could originate from previous reports of toxicity of *Galega officinalis*, *Agentum conyzoles* L., *Calendula officinalis*, *Cedrus deodara* (Adebayo, *et al.*, 2010). In our work, the major haematological parameters and organic enzymes were provided on the health status of the animal models, this method is used as a preliminary or decisive step in medical diagnosis before treatment (Wasser *et al.*, 2004). Mean percentage PCV of the blood samples of 15 Albino rats treated with oil of *A. indica* seed was within the control value. The value obtained was in agreement with that reported by Wasser *et al.*, (2004) on the effect of *Gonderma spp* on haematological parameter of mice model, but different from that reported by Shamaki *et al.*, (2014) but using the same plant material on Albino rat. The variation in the values of the parameter reported could be attributed to the high dosage of the plant material which was administered, time of exposure, sex, age of the animal models and other experimental conditions. In real life situation, high residues of the *A. indica* seed oil are rare in processed food after washing and boiling, therefore treated foods may not contain residual concentration high enough to alter blood parameters above the control values. Therefore results of our study

showed no significant variations in pack cell volume (PCV), red blood cell (RBC), white cell count (WCC), and haemoglobin content (Hb), very slight variations were noticed with increasing age of the animals except in run 14 which did not show reasonable values, this was in agreement with the observation of Adel *et al.*, (2015). The observation suggest that the *A. indica* seed oil failed to alter most biochemical processes in the animal models to detectable level at the levels and time administered. Catalase (CAT), superoxide dismutase (SOD) and glutathione (GSH) are some of the important endogenous antioxidants (Geta *et al.*, 2002) which counteract the damaging effects of free radicals in living system, the marginal variation among the animal models did not suggest toxic effect of the plant material. This is not surprising because *A. indica* plant has a good history of remedying liver cirrhosis, inflammation and cancer (el-Shazy *et al.*, 2000), and is expected to control oxidative stress at the level of administration (Tables 2, 3).

In the study, it could be concluded that the of *A. indica* seed oil which was administered orally to the animal models did not alter haematological indices and antioxidant values significantly from the control. The slight changes observed did not mean that the animals were sick but could be attributed to experimental errors. Therefore, the use of *A. indica* oil as an alternative to some chemical synthetic insecticides should be considered.

V. CONCLUSION

Azadirachta indica seed oil may be a safe alternative to the synthetic chemical insecticides. Because the plant material failed to elicit appreciable alteration on the haematological indices and antioxidant levels in Albino rats at the conditions of administration. The use of the plant material as an alternative to the synthetic chemical insecticides should be encouraged, but more work should be carried out on higher levels of other independent variables.

Conflicts of interest: The authors declare no conflict of interest.

REFERENCES

- [1] Adebayo, J. O., Adesokan, L. A., Olatunji, D. O., Burro, D., Soladoye, A. O. (2005) Effect of ethanolic extract of *Bougainvillea spectabilis* leaves on haematological and serum lipid variables in rats. *Biochemistry*, (17): 45-48.
- [2] Adel, K., Raizman, J., Chen, Y., (2015) Complex biological profile of hematologic markers across pediatric, adult, and geriatric ages: establishment of robust pediatric and adult reference intervals on the basis of the Canadian Health Measures Survey. *Clin Chem* 61: 8.
- [3] Aebi, H. E. (1983) Catalase. In *Methods of enzymatic analysis*, 3rd ed., Bergmeyer, H. U., Verlag Chemic. Weinheim, FL, USA., pp 273-286.
- [4] Adewale O. O., Oduyemi, O. I., Ayokunle, O. (2014) Oral administration of leaf extracts of *Momordica charantia* affect reproductive hormones of adult female Wistar rats. *Asian Pac J Trop Biomed.* 4(Suppl 1):S521-4.
- [5] Alle, A., Dembele, A., Yao, B and Ado, G. (2009) Distribution of organochlorine pesticides in human breast milk and adipose tissue from two locations in Cote d'ivoire. *Asian J. Applied Sci.*, 2: 456-463.
- [6] Chaudhary, S., Kanwar, R. K., Sehgal, A., Cahill, D. M., Barrow, C. J., Sehgal, R., & Kanwar, J. R. (2017). Progress on *Azadirachta indica* based biopesticides in replacing synthetic toxic pesticides. *Frontiers in Plant Science*, 8, 610.
- [7] CLSI (Clinical and Laboratory Standards Institute) (2000) How to define and determine reference intervals in the clinical laboratory: approved guideline. Wayne, PA, USA (CLSI document C28-A2).
- [8] Coats, J. R. (1999) Mechanisms of toxic and structure-activity relationships for organochlorine and synthetic pyrethroid insecticides. *Environmental health perspectives* (87): 255-262.
- [9] El-Shzygy, M. M. and el-sharnoubi, E. D. (2000) toxicity of a *Neem* (*A. indica*) Insecticide to certain aquatic organisms *Arch. Envir. Contam Toxicol.* 39(3): 329-36.
- [10] El-Wakeil, N. E. (2013) Botanical pesticides and their mode of action. *Gesunde Pflanzen* 65:125-149.
- [11] Gaeta L. M.; Tozzi, G., Pastore, A.; Federici, G.; Bertini, E.; Piemonte, F. (2002) Determination of superoxide dismutase and glutathione peroxidase activities in blood of healthy paediatric subjects. *Clin Chim Acta.* 322(1-2):117-20.
- [12] Gerald, R. S., Ian, G. F., Patrick, T. H., and Monica, N (2006) Glossary of Terms Relating Pesticides (IUPAC Recommendations 2006). *Pure Appl. Chem.*, 78 (11): 2075–2154.
- [13] Han Sub Kwak, Misook Kim and Yoonhwa Jeong (2015) Developing water -extracted cacao-coffee beverage: effects of temperature and time on cacao roasting and the beverage acceptability. *CyTA - Journal of Food.*14(2): 248-254.
- [14] Huang, Y. and Ho, S.H. (1998). Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Journal of Stored Products Research* 34: 11–17.

- [15] Igbokwe, C. O., Agina, O. A., Okoye, C. N., Onoja, R. I (2017) Haematological and serum biochemistry profile of the juvenile wild African giant rat (*Cricetomys gambianus*, Waterhouse – 1840) in Nsukka, South-Eastern Nigeria – a preliminary investigation. *Journal of Applied Medical Research*. 45(1): 190-194.
- [16] Kanu K. C., Ijioma, S. N. and Atiata, O. (2016) Haematological, biochemical and antioxidant Changes in Wistar Rats Exposed to Dichlorvos based insecticide formulation used in southeast Nigeria. *Toxics* 2016, 4, 28.
- [17] Kwak, H.S., Kim, Misook, Jeong, Yoonhwa (2016) Development water - extracted cocoa - coffee beverage: effects of temperature and time on cocoa roasting and beverage acceptability. *CyTA - Journal of Food* 14 (2): 248 - 254.
- [18] Lars, H.; Johansson, L. A.; Håkan, B. (1987) A spectrophotometric method for determination of catalase activity in small tissue samples. *Journal of Analytical Biochemistry*. 174(1): 331-336.
- [19] Meyers, R. H., Montgomery, C. (2002) Response surface methodology: process and product optimization using designed experiments. New York: John Wiley and Sons. p 798.
- [20] Mikhael, A. A. (2011). Potential of some volatile oils in protecting packages of irradiated wheat flour against *Ephestia kuheniella* and *Tribolium castaneum*. *Journal of Stored Products Research*, 47(4): 357–364.
- [21] Mitra, A., Chatterjee, C., Mandal, F. B. (2011) Synthetic pesticide and their effects on birds. *Research Journal of Environmental Toxicology* (5(2):81-96.
- [22] Ndodo, N. D., Anuka, J. A., Esomonu, U. G., Onu, J. E., Okolo, R. U. and Onwuchekwa, C. (2013) *IOSR Journal of Pharmacy*, 3(8): 24-28.
- [23] Negahdar, M., Djalali, M., Abtahi, H., Sadeghi, M. R., Aghvami, T., Javadi, E., Layegh, H. (2005) Blood Superoxide Dismutase and Catalase Activities in Women Affected with Breast Cancer. *Iranian J Publ Health*, 34(3): 39-43.
- [24] Ojmelukwe, P. C. (2000). Potentials of *Xylopiia Aethiopica* for short-term protection of cowpea (VAR IT-81D-975) seeds in Nigeria. *Nigerian Agricultural Journal*. 31(2000)
- [25] Ojmelukwe, P. C. and Adler, C. (1999). Potential of Zimtaldehyde, 4-Allyl-anisol, Linalool, Terpinol and other phytochemicals for the control of the confused flour beetle (*Tribolium confusum* Duval) (Col., Tenebrionidae). *J. Pest Sci*. 72: 81-86.
- [26] Ojmelukwe, P. C. and Ogwumike, F. C. (1999). Effects of infestation by Bruchid Beetles (*Callosobruchus maculatus*) on the nutritional quality and sensory properties of cowpeas (*Vigna unguiculata*) *Journal of Food Biochemistry* 23(6): 637-645.
- [27] Ojmelukwe, P. C. (2008). Biosafety of potential plant products for the protection of selected grains against insect pest. XXIII International Congress of Entomology. July, 2008. International Convention Center, Durban.
- [28] Ojmelukwe, P.C. and Adler, C. (2000). Toxicity and repellent effects of eugenol, thymol, linalool, menthol and other pure compounds on *Dinoderus bifoveatus* (Coleoptera: Bostrichidae). *J. Sustain. Agric. Environ*. 2:47-54.
- [29] Okigbo, (2008) Antimicrobial effects of *Piper Guineense* 'Uziza' and *Phyllanthus amarus* 'Ebe-Benizo' on *Candida albicans* and *Streptococcus faecalis*. *Acta Microbiologica et Immunologica Hungarica*, 54(4):353-66.
- [30] Oparaeke, A. M. (2007). Toxicity and spraying schedules of biopesticide prepared from *Piper guineense* against two cowpea pests. *Plant Protect. Sci*. 43(3):103-108.
- [31] Oparaeke, A. M. and Bunni, J. O. (2006). Bioactivity of two powdered spices (*Piper guineense* Honn and Schum and *Xylopiia aethiopica* (Dunal) A. Richard) as home made insecticides against *Callosobruchus subinnotatus* (Pic.) on stored Barbara groundnut. *Agricultura Tropica et Subtropica*, 39(2): 132-134.
- [32] Pallav, K. D., Ragini, G., Anupam, Kumar, P. (2014) Phytochemical analysis and evaluation of antimalarial activity of *Azadirachta indica* *The Pharma Innovation Journal*; 3(9): 12-16.
- [33] Shamaki, B. U., Ogbe, A. O., Abdulrahman, F. I. and Sandabe, U. K. (2014) Haematological parameters in normal albino rats following oral administration of methanolic extract of wild *Ganodema sp.* *International Journal of pharmaceutical screening methods*. 4(2):94-99.
- [34] Schmitterer, H. and Sscher, K. R. S. (1987) Natural insecticides from the neem trees, *A.indica*, A. Juss. and other tropical plants. Proc. 2nd Int. Neem Conf., Rauschholzhausen, 25-28 May, 1983.
- [35] Wararut, B., Supap, S., Siriwadee, C. and Kanokporn, S. (2012) Evaluation of biochemical, hematological and histopathological parameters of albino rats treated with *Stemona aphylla* Craib. Extract. *Journal of Medicinal Plants research* 6(27): 4429-4435.
- [36] Yin, H., Xu, L., and Porter, N. A. (2011) Free radical lipid peroxidation: mechanisms and analysis. *Chemical reviews*. 111: 5944-5972.