



PHARMACOLOGICAL ACTIVITIES OF PYRAZOLONE DERIVATIVES

Rishi Ram Parajuli, Priyanka Pokhrel, Amish Kumar Tiwari, Janmajoy Banerjee*

Department of Pharmacy, Sunsari Technical College, Dharan, Nepal

Pyrazoline is a five member heterocyclic ring which is a versatile lead compound for designing potent bioactive agents. The review of the literature shows that the pyrazoline derivatives are quite stable and has inspired the chemists to synthesize the new pyrazoline derivatives. The past studies of pyrazoline derivative revealed that they are useful in pharmaceutical and agrochemical research. Pyrazoline derivatives display various pharmacological activities such as antitumor, antitubercular, antimicrobial, antibacterial, anti-inflammatory and antioxidant etc. and the pharmacological activities of different synthesized compound are reviewed in the present article.

Key words: Pyrazoline, antimicrobial, anti-inflammatory, antitumor activity


INTRODUCTION

Pyrazoles are five member ring heterocyclic compounds, have some structural features with two nitrogen atoms in adjacent position and are also called as Azoles [1]. It has only one endocyclic double bond and is basic in nature. The best described property of almost every group of pyrazoles is in the treatment of inflammation and inflammation associated disorders, such as arthritis [2]. Pyrazole derivatives are the subject of many research studies due to their widespread potential biological activities such as antimicrobial [3], antiviral [4], antitumor [5, 6], antihistaminic [7], antidepressant [8], anti-inflammatory [9], tranquilizing [10], anticancer [11, 12], anti-hypertensive [13], anti-arrhythmic [14], psychoanaleptic [15], anticonvulsant [16] & anti-diabetic activities [17]. Mostly pyrazolone are synthesized taking ethylacetacetate and substituted hydrazine as starting chemicals. Pyrazolone was first prepared by Knorr in 1883 when he was trying to synthesis quinoline derivatives, but he obtained pyrazolone derivative called antipyrin, and also called phenazone. Pyrazolone can be considered as intermediate compound for synthesis of various cyclic compounds of high biological activity.

Pyrazoline is basic in nature. An intra molecular conjugated charge transfer process has been reported to exist in it in the excited state.

The conjugated part (–N1–N2–C3–) of the ring, the


nitrogen atom at the 1-position and the carbon atom at the 3-position are, respectively, electron donating and withdrawing moieties. The carbon atoms at 4- and 5-positions do not conjugate with the remaining part of the ring. In general pyrazolones are yellow solids with high melting points and moderate solubility in water, alkaline and acidic solutions. They react with aldehydes in alcoholic basic medium and with the bromine to give high biologically active compounds. It is reported that pyrazolones undergo tautomerism to certain extent which are affected by substituent and solvents.



Pharmacological Activities

[A] Antibacterial activity

Mogilaiah et al [18] synthesized & found antibacterial activities of 1, 3, 4-Oxadiazole and pyrazoline derivatives containing 1, 8-Naphthyridine moiety. All the compounds were far less active than the standard drug (gentamycin) taken.




Vergiya et al [19] synthesized some new 3, 5-Diaryl-1-phenyl/isonicotinoyl-2-pyrazolines and evaluated its


For Correspondence

jj.banderjee983@gmail.com


biological activity. All the synthesized compounds showed antibacterial activity against Gram +ve bacteria *S. Aureus*, *S. albus*, *S. pyogenes*, *S. Viridians* and Gram -ve bacteria *E. coli*, *S. Typhosa*



Waheed et al [20] synthesized certain substituted 1, 2-Pyrazolines from nalidixic acid as antibacterial and analgesic agents. They were found to have significant antibacterial activity against Gram -ve bacteria and possessed appreciable analgesic activity.




Abdullah et al [21] Synthesized of new 3-(5-aryl-4,5-dihydro-1H-pyrazol-3-yl)-4-hydroxy-2H-chromene-2-one derivatives and evaluation of their antibacterial activity.




[B] Antitubercular activity


Kini et al [22] synthesized a novel series of heterocyclic o/m/psubstituteddiphenyl ether derivatives and determined their activity against H37Rv strain of Mycobacterium. All 10 compounds inhibited the growth at concentrations as low as 1 $\mu\text{g.ml}^{-1}$. This level of activity was found comparable to the reference drugs rifampicin and isoniazid at the same concentration.




Ali et al [23] synthesized a series of 5-(4-(Substituted)phenyl)-3-(4-hydroxy-3-methylphenyl)-4,5-dihydro-1H-1-pyrazolyl-2-toluidino methanethione and 5-(Substituted)phenyl-3- (4-hydroxy-3-methylphenyl)- 4, 5 – dihydro – 1H – 1 – pyrazolyl – 2 – methoxyanilino methanethione and tested for their *in vitro* antitubercular activity against *M. tuberculosis* H37Rv. Among the synthesized compounds, compound Anilino – 3 - (4 – hydroxyl – 3 – methylphenyl) – 5 - (2, 6-dichloro phenyl)-4, 5-dihydro- 1H- 1-pyrazolyl methanethione was found to be more active agent against *M. tuberculosis* H37Rv with minimum inhibitory concentration of 0.0034 μM .



Babu et al [24] synthesized and evaluated biological activity of 1, 3, 5 – Trisubstituted pyrazolines bearing benzofuran. They were found to be antitubercular, antimicrobial and anti-inflammatory in nature.




Joshi et al [25] Synthesized a series of 1-acetyl-3,5-diphenyl-4,5-dihydro-(1H)-pyrazole derivatives and evaluation of their antitubercular activity.




[C] Antimicrobial activity


Ozdemir et al [26] synthesized several 1-(4-Aryl-2-thiazolyl)-3-(2-thienyl)-5-aryl-2-pyrazoline derivatives and investigated their antimicrobial activities against *Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhimurium*, *Bacillus cereus*, *Streptococcus faecalis*, *Aeromonas hydrophila*, *Candida albicans* and *Candida glabrata*. A significant level of activity was observed.




Abdelwahab et al [27] synthesized 1-(Benzofuran-2-yl)-4-nitro-3-arylbutan-1-ones and 3(Benzofuran-2-yl)-4,5-dihydro-5-aryl-1-[4-(aryl)-1,3-thiazol-2-yl]-1H-pyrazoles and evaluated their antibacterial and antifungal activities at 100 µg concentration. Some of the compounds showed excellent antimicrobial activities than control drugs.




Stirrett et al [28] synthesized small molecules with structural similarities to siderophores and evaluated as novel antimicrobials against *Mycobacterium tuberculosis* and *Yersinia pestis*.



Agrawal et al [29] synthesized 1,3,5-trisubstituted pyrazoline derivatives and screening for their antimicrobial activity.




Hassan [30] synthesized new pyrazoline derivatives and evaluated their anti microbial activity.



[D]Anti-inflammatory activity


Amir et al [31] synthesized a series of 3-(4-Biphenyl)-5-substituted phenyl-2-pyrazolines and 1-Benzoyl-3-(4-biphenyl)-5-substituted phenyl-2- pyrazolines and screened for their anti-inflammatory and analgesic activity. Among the compounds studied, compound-17 showed more potent anti-inflammatory and analgesic activity than the standard drug, along with minimum ulcerogenic index.




Barsoum et al [32] synthesized a variety of Bis(3 – aryl – 4, 5 – dihydro – 1H – pyrazole – 1 – carboxamides) and screened for their anti-inflammatory properties and

PGE2 inhibitory properties (at a dose level of 50 mg.kg⁻¹) utilizing *in vivo* acute carrageenan – induced paw oedema standard method in rats. They exhibited that many of the tested compounds reveal considerable anti-inflammatory properties, especially which reveal remarkable activities related to indomethacin (which was used as a reference standard at a dose of 10 mg.kg⁻¹ of body weight). They exhibited lower ulcer index

values than the used reference standard (indomethacin).




Khalil [33] synthesized 1 – acetyl/propanoyl – 5 – aryl – 3 – (4 – morpholinophenyl) – 4, 5 – dihydro – 1H – pyrazole derivatives and evaluated their anti-inflammatory activity.




Mahesh et al [34] synthesized a new series of 3, 5-diaryl-1-phenyl-2-pyrazoline by cyclocondensation of chalcones with phenyl hydrazine. The compounds were screened *in vivo* for their anti – inflammatory activity by using carrageenan induced rat paw edema method. Compounds P4, P7, P8 , P10 have shown maximum anti-inflammatory activity.

[E] ANTITUMOR ACTIVITY


Braulio et al [35] synthesized a series of novel 3- aryl- 4-(3- aryl -4, 5-dihydro-1H-pyrazol-5-yl)-1-phenyl -1H-pyrazoles and 1-substituted 3-aryl-5-aryl (hetaryl)-2-pyrazolines and screened their antitumor activity.



Zhu et al [36] Synthesized a series of 3-(substituted phenyl) - 5-(substituted phenyl) - 4, 5-dihydro-1H-pyrazole- 1-carbothioamide and 1-(5-(substituted phenyl)-3-(substituted phenyl)-4, 5-dihydro-1H-pyrazol-1-yl) ethanone as anticancer agents.




Havrylyuk et al [37] synthesized several novel thiazolone based compounds containing 5-Aryl-3-phenyl-4, 5-dihydro-1Hpyrazol-1-yl framework and tested for in vitro anticancer activity. Most of them displayed anticancer activity on leukemia, melanoma, lung, colon, CNS, ovarian, renal, and prostate and breast cancer cell lines. The most efficient anticancer compound-39 was found to be active with selective influence on colon cancer cell lines, especially on HT 29 ($\log GI_{50} = -6.37$).




[F] ANALGESIC ACTIVITY:

Shau et al [38] synthesized a series of novel 4-(5-substituted aryl-4, 5-dihydropyrazole-3-yl-amino) phenols 2a-f where introduction of p-nitro and phdroxy group in aryl moiety of the pyrazoleanalogs 2c and 2e produce compounds with potent analgesic, anti-inflammatory and, in a few cases, antimicrobial properties.



Sivakumar et al [39], synthesized a series of (4Z)-3-methyl-1-[(2-oxo-2H-chromen-4-yl) carbonyl]-1H-pyrazole- 4, 5- dione 4- [(4- substituted phenyl) hydrazone] (5a-i). The titled compounds were screened for their anti-inflammatory and analgesic activity. Among the synthesized compounds, compound 5a, 5c, 5g and 5h exhibited significant anti-microbial activity

and compound 5a, 5b, 5d, 5h and 5i exhibited significant analgesic activity compared with the standard drug (indomethacin 5mg/kg) at the dose level of 50mg/kg on oral administration.




[G] ANTIFUNGAL ACTIVITY:

Baseer et al [40] synthesized fourteen new N-acetylated and non-acetylated pyrazoline derivatives by reacting chalcones with hydrazine in the presence of absolute ethanol however reaction was carried out in the presence of glacial acetic acid to afford N-acetylated pyrazolines. The pyrazolines (1-14) were screened for antifungal activity against *Aspergillus flavus*, *Aspergillus niger* and *Aspergillus pterus*. Pyrazolines (1-14) found to exhibit well to excellent anti fungal properties


Ramiz et al [41] synthesized a series of new pyrazolone and pyrazole derivatives with expected anti fungicidal activity through the reactions 3-phenyl-1-H-pyrazol-5(4H)- one (3) and 4-(dimethyl amino methylene)-3-phenyl-1H-pyrazol-5(4H)-one (5) with a variety of electrophilic reagents and nucleophilic reagents.

[H] ANTICONVULSANT ACTIVITY:

Singh et al [42] synthesized a series of 1-[(4,5-dihydro-5-phenyl-3-(phenyl amino) pyrazol-1yl)] ethanone derivatives I-VI were synthesized and evaluated for their anticonvulsant activity against electric shock induced convulsion method. Compounds III and V are found to be the most potent compounds of all synthesized compounds.




Singh et al [43] synthesized several 3-(3-Acetoamino)phenyl-1, 5-substituted phenyl-2-pyrazolines [21] which were evaluated for their anticonvulsant activity. All the substituted pyrazolines exhibited anticonvulsant activity, which was reflected by 30-80% protection observed against PTZ-induced seizures.




[1] ANTIHELMINTIC ACTIVITY:

Sreenivasa et al [44] synthesized a series of pyrazole derivatives and evaluated for their anthelmintic activity. Synthesized compounds of pyrazole derivatives were tested for anthelmintic activity against earthworms, *Peritumaposthuma* compared to standard Albendazole. VII P8, VIII P6, VIII P7, VIII P8, VIII P9, VIII P10, VIII P11, VIII P12 showed significant activity compared to standard Albendazole.




[J] ANTIOXIDANT ACTIVITY


Pasin et al [45] synthesized a series of pyrazole derivatives and screened for their antioxidant activity. All compound showing good activity.



Umesh et al [46] Synthesized of 5-methyl-2-(5-methyl-1,3-diphenyl-1H-pyrazole-4-carbonyl)-2,4-dihydro-pyrazol-3-one and evaluation of their antioxidant activity.




Babu et al [47] synthesized a series of pyrazoline derivatives and evaluated antioxidant activity at 1000, 500, 250, 100, 50, 25 and 10 mg.ml⁻¹ concentrations against standard drug ascorbic acid. Six of the synthesized compounds showed interesting antioxidant activity as compared with ascorbic acid.




[K] CYTOTOXIC ACTIVITY

Zahar et al [48] synthesized a series of pyrazole derivatives which show good activity




[L] HERBICIDAL

Noriaki et al [49] synthesized a series of 1,5-Diarylpyrazole Derivatives Some of these compounds showed noticeable pre-emergent herbicidal activities against various kinds of weeds. Among the synthesized compounds, methyl 4-chloro-1-(2,5-difluorophenyl)-5-(4-fluorophenyl)-pyrazole-3-carboxylate 19t exhibited good activity. Diarylimidazole carboxylates and carboxamides were also synthesized, but they did not show any herbicidal activities.




[M] ACE INHIBITORY ACTIVITY

Bonesi et al [50] synthesized a series of pyrazole derivatives and investigated their angiotensin I converting enzyme inhibitory activity by performing assay. The following pyrazole derivative-25 showed effective ACE-inhibitory activity with 0.123m MIC value.




[N] ANTIHYPERTENSIVE ACTIVITY

Zitouni et al [51] synthesized some 1-(4-Arylthiazol-2-yl)-3, 5-diaryl-2- pyrazoline derivatives and investigated their hypotensive activity by the tail-cut method using clonidine as reference compound. All examined compounds showed appreciable hypotensive activities. One of the active compounds is given below



[O] CHOLESTEROL INHIBITORY ACTIVITY:

Jeong et al [52] synthesized a series of 3-(3, 5-Di-tert-butyl-4-hydroxyphenyl)-5-(multi-substituted 4-hydroxy phenyl)-2-pyrazolines and carried out their inhibitory activity on acyl-CoA: cholesterol acyltransferase. They showed invitroinhibitory activity on hACAT-1 and -2. One of the active compounds is shown below



CONCLUSION:

Pyrazolone moiety and its various derivatives was studied frequently in the past time and found potent in various pharmacological and pathological conditions, which are discussed in brief in this article. In conclusion, we have described the biological activities of pyrazoline derivatives such as antitumor, antitubercular, antimicrobial, antibacterial, anti-inflammatory and antioxidant etc. Various other activities are also been studied like anticonvulsant,

anticancer, antihelmintic etc. Thus by studying all the derivatives showing variety of activities we can say that pyrazole ring have been explored in past years and is still be used for future development of new drugs against many more pathological conditions.

ACKNOWLEDGMENT

The corresponding author expresses deep gratitude to Mr. Janmajoy Banerjee and friends for their co-operation and guidance in searching various articles and journals for completion of this review.

REFERENCE:

1. T, Eicher, S.Hauptmann, (2003), Edition IIInd, The Chemistry of Heterocycles: Structure, Reactions, Syntheses, and Applications, Wiley-VCH, ISBN 3527307206.
2. J. John. Talley, J. Donald, Jr. Rogier, both of St. Louis, Mo. G.D. Searle & Co., Skokie, 1995, Pt No: 5, 434,178.
3. E.V. Pimerova, E.V. Voronina., Pharm. Chem. J., 2001, 35, 18-20.
4. S.L. Janus; A.Z. Magdif ; B.P. Erik; N.Claus; Chem., 1999, 130, 1167-1174
5. H.J. Park, K.Lee, S.Park, B. Ahn, J.C.Lee, H.Y. Cho, K.I. Lee. Bioorg. Med. Chem. Lett., 2005, 15, 3307- 3312.
6. I. Bouabdallah, L.A M'barek, A. Zyad, A. Ramadan, I. Zidane, A.Melhaoui, Nat. Prod. Res., 2006, 20, 1024-1030.
7. I. Yildirim, N. Ozdemir, Y.Akçamur, M. Dinçer, O.Andağ, ActaCryst., 2005, E61, 256-258.
8. D.M. Bailey, P.E. Hansen, A.G. Hlavac, E.R. Baizman, J.Pearl, A.F. Defelice, M.E Feigenson, J. Med. Chem.,1985, 28, 256-260.
9. M. Amir, H. Kumar, S. A. Khan, Bioorg." Synthesis and pharmacological evaluation of pyrazoline derivatives as new anti-inflammatory and analgesic agents". Med. Chem. Lett. 18, 918- 922, 2008.
10. E. Palaska, M. Aytemir, I. T. Uzbay, D. Erol," Synthesis and antidepressant activities of some 3,5-diphenyl-2- pyrazolines. Eur. J. Med. Chem. 36, 539-543, 2001
11. E. Nassar, "Synthesis, (in vitro) antitumor and antimicrobial activity of some pyrazoline, pyridine and pyrimidine derivatives linked to indolesmoiety" J. Am. Sci, 6, 463-471, 2010.
12. D. Azarifar, M. Shaebanzadeh, "Synthesis and characterization of new 3,5-dinaphthyl substituted 2-pyrazolines an study of their antimicrobial activity" Molecules 7, 885-895, 2002 .
13. H. Dmytro, Z. Borys, V. Olexandr, Z. Lucjusz, G. Andrzej, L. Roman, "Synthesis of novel thiazolone based compound containing pyrazoline moiety and evaluation of their anticancer activity". Eur. J. Med. Chem. 44, 1396-1404, 2009.
14. M. S. Mui, B. N. Siew, A. D. Buss, S. C. Crasta, L. G. Kah, K. L. Sue, "Synthesis of N-1 acidic functionality affording analogues with enhanced antiviral activity against HIV" Bioorg. Med. Chem. Lett. 12, 679-699, 2002.
15. S. S. Parmar, B. R. Pandey, C. Dwivedi, R. D. Harbison, "An-ticonvulsant activity and monoamine oxidase inhibitory properties of 1,3,5-trisubstituted pyrazolines" J. Pharm. Sci. 63, 1152- 1155, 1974.
16. A. A. Siddiqui, M. A. Rahman, M. Shaharyar, R. Mishra, "Synthesis and anticonvulsant activity of some 3,5-diphenyl- 2-pyrazoline-1-carboxamide derivatives" Chem. Sci. J. 8, 1-10, 2010 .
17. N. Soni, K. Pande, R. Kalsi, T. K. Gupta, S. S. Parmar, J. B. Barthwal, Inhibition of rat brain monoamine oxidase and suc-cinic dehydrogenase by anticonvulsant pyrazolines "Res. Commun. Pathol. Pharmacol. 56, 129-132, 1987
18. Mogilaiah K, Sakram B. Synthesis and antibacterial activities of 1,3, 4-oxadiazole and pyrazoline derivatives containing 1, 8-naphthyridine moiety. Ind J Het Chem. 2004; 13:289-292.
19. Vijayvergiya D, Kothari S, Verma BL. Synthesis and biological activity of some new 3,5-diaryl-1-

- phenyl/ isonicotinoyl-2-pyrazolines. *Ind J Het Chem.* 2003; 13:105-110.
20. Waheed A, Khan SA. Synthesis of certain substituted 1, 2-pyrazolines from nalidixic acid as antibacterial and analgesic agents. *Ind J Het Chem.* 2001; 11:59-62.
21. Abdulla Sulaiman Al-Ayed, Synthesis, spectroscopy and electrochemistry of new 3-(5-aryl- 4,5-dihydro-1H-pyrazol-3-yl)-4-hydroxy-2H-chromene-2-one as a novel class of potential antibacterial and antioxidant derivatives, *Inter. J. Org.Chem.*, 1: 87-96, (2011)
22. Kini SG, Bhat AR, Bryant B, Williamson JS, Dayan FE. Synthesis, antitubercular activity and docking study of novel cyclic azole substituted diphenyl ether derivatives. *Eur J Med Chem.* 2008; 41: 1-9.
23. Ali MA, Shaharyar M, Siddiqui AA. Synthesis, structural activity relationship and anti-tubercular activity of novel pyrazoline derivatives. *Eur J Med Chem.* 2007; 42:268-275.
24. Babu VH, Manna SK, Sneha, Srinivasan KK, Bhatt GV. Synthesis and biological evaluation of 1, 3, 5-trisubstituted pyrazolines bearing benzofuran. *Ind J Het Chem.* 2004; 13:253-256.
25. Chovatia PT, Akabarij JD, KachhadiaPK, Zalavadia PD and Joshi HS, Synthesis and selective antitubercular and antimicrobial inhibitory activity of 1-acetyl-3,5-diphenyl-4,5-dihydro-(1H)-pyrazole derivatives, *J. Serb. Chem. Soc.*, 71(7):713–720, (2007)
26. Ozdemir A, TuranZitouni G, Kaplancıklı ZA, Revial G, GuvenK. Synthesis and antimicrobial activity of 1-(4-aryl-2-thiazolyl)-3-(2-thienyl)-5-aryl-2-pyrazoline derivatives. *Eur J Med Chem.* 2007; 42: 403-409.
27. AbdelWahab BA, AbdelAziz HA, Ahmed EM. Synthesis and antimicrobial evaluation of 1-(benzofuran-2-yl)-4-nitro-3-arylbutan-1-ones and 3-(benzofuran-2-yl)-4,5-dihydro-5-aryl-1-[4-(aryl)-1,3-thiazol-2-yl]-1H-pyrazoles. *Eur J Med Chem.* 2008; 43:1-4.
28. Stirrett KL, Ferreras JA, Jayaprakash V, Sinha BN, Renc T, Quadri LEN. Small molecules with structural similarities to siderophores as novel antimicrobials against *Mycobacterium tuberculosis* and *Yersinia pestis*. *Bioorg Med ChemLett.* 2008; 18:2662-2668.
29. Manish Agrawal, Pankaj Kumar Sonar, Shailendra Saraf K, Synthesis of 1,3,5-trisubstituted pyrazoline nucleus containing compounds and screening for antimicrobial activity, *Med. Chem. Res.*, 21: 3376–3381(2012)
30. Seham Hassan Y, Synthesis and biological activity of some new pyrazoline and pyrimidine derivatives, *J. Braz. Chem.Soc.*, 22(7): 1286-1298, (2011)
31. Amir M, Kumar H, Khan SA. Synthesis and pharmacological evaluation of pyrazoline derivatives as new anti-inflammatory and analgesic agents. *Bioorg Med ChemLett.* 2008; 18:918-922.
32. Barsoum FF, Grgis AS. Facile synthesis of bis (4,5-dihydro-1Hpyrazole-1-carboxamides) and their thio-analogues of potential PGE2 inhibitory properties. *Eur J Med Chem.* 2008; 15:1-6.
33. Omneya Khalil M, Synthesis and anti-inflammatory activity of 1-acetyl/propanoyl-5-aryl-3-(4-morpholinophenyl)-4,5-dihydro-1Hpyrazole derivatives, *Med. Chem. Res.*, 21: 3240–3245, (2012)
34. Mahesh. D. Bhanushali, Shivkumar. M. Hipparagi, Suhas. A. Patil, Sudhir. N. Sambrekar, Synthesis and biological screening of 3, 5-diaryl-1-phenyl-2-pyrazolines as anti-inflammatory agents.
35. BraulioInsuasty, Alexis Tigreros, Fabian Orozco, JairoQuiroga, Rodrigo Abonia, ManuelNogueras, Adolfo Sanchez, and Justo Cobo. Synthesis of novel pyrazolic analogues of chalcones and their 3-aryl-4-(3-aryl-4,5-dihydro-1H-pyrazol-5-yl)-1-phenyl-1H-pyrazole derivatives as potential antitumor agents, *Bioorg. Med.Chem.*, 2010, 18(14): 4965–4974

36. Peng-Cheng Lv, Huan-Qiu Li, Juan Sun, Yang Zhou, and Hai-Liang Zhu, Synthesis and biological evaluation of pyrazole derivatives containing thiourea skeleton as anticancer agents, *Bioorg. Med. Chem.*, 18(13): 4606–4614, (2010)
37. Havrylyuk D, Zimenkovsky B, Vasylenko O, Zaprutko L, Gzella A, Lesyk R. Synthesis of novel thiazolone based compounds containing pyrazoline moiety and evaluation of their anticancer activity. *Eur J Med Chem.* 2008; 42:1–9.
38. SK Sahu, M Banerjee, A Samantray, C Behera and MA. Synthesis, Analgesic, Anti-inflammatory and Antimicrobial Activities of Some Novel Pyrazoline Derivatives
39. K. K. Sivakumar, A. Rajasekaran, I. Ponnilarasan, A. Somasundaram, R. Sivasakthi, S. Kamalaveni. *Der Pharmacia Lettre*, 2010, 2, 211-219.
40. Muhammad Baseer, FarzanaLatif Ansari1 and Zaman Ashraf, Synthesis and in vitro antibacterial and antifungal activity of some n-acetylated and non-acetylated pyrazolines
41. Mahmoud. M. M. Ramiz, Ibrahim. S. Abdel Hafiz, Mohamed. A. M. Abdel Reheimb and Hatem M. Gaber. Pyrazolones as Building Blocks in Heterocyclic Synthesis: Synthesis of New Pyrazolopyran, Pyrazolopyridazine and Pyrazole Derivatives of Expected Antifungal Activity *Journal of the Chinese Chemical Society*.
42. Anoop Singh and A. C. Rana, *J. Chem. Pharm. Res.* 2011, 2, 505-511.
43. Singh SP, Chaudhari A, Barthwal JP, Parmar SS. Anticonvulsant activity and selective inhibition of nicotinamide adenine dinucleotide-dependent oxidations by 1, 3, 5-trisubstituted pyrazolines. Wiley interscience, 1974.
44. Sreenivasa G.M., Jayachandran E., Shivakumar, B., Jayaraj Kumar K, Vijay Kumar M.M.J. *Arch Pharm Sci& Res.* 2009, 1, 150 – 157.
45. J.S.M. Pasin, A.P.O. Ferreira, A.L.L. Saraiva, V. Ratzlaff, R. Andriguetto, P. Machado, S. Marchesan, R.A. Zanette, H.G. Bonacorso, N. Zanatta, M.A.P. Martins, J. Ferreira and C.F. Mello, *Braz J Med Biol Res*, 2010, 43, 1193-1202.
46. Umesha KB, Rai KML, Harish NayakaMA, Antioxidant and antimicrobial activity of 5-methyl -2-(5-methyl-1,3-diphenyl- 1Hpyrazole -4-carbonyl)- 2,4- dihydro-pyrazol-3-one, *Int. J. Biomed. Sci.*, 5(4): 359-368,(2009)
47. Babu VH, Sridevi C, Joseph A, Srinivasan KK. Synthesis and biologicalevaluation of some novel pyrazolines. *Ind. J. Pharm. Sci.*, 2008; 69:470-473.
48. Magdy I. El-Zahar, *World J of Chemistry*, 2009, 4, 182-194.
49. Noriaki Kudo, Satoru Furuta, *Chem. Pharm. Bull.* , 1999, 47, 857—868.
50. Babu VH, Sridevi C, Joseph A, Srinivasan KK. Synthesis and biological evaluation of some novel pyrazolines. *Ind. J. Pharm. Sci.*, 2008; 69:470-473.
51. Bonesi M, Loizzo M R, Statti G A, Michel S, Tillequin F, Menichini. The synthesis and ACE inhibitory activity of chalcones and their pyrazole derivatives. *Bioorg. Med. Chem. Lett*, 2010; 20: 1990–1993
52. Srivastava BK, Joharapurkar A, Raval S, et al. Diaryl dihydropyrazole-3-carboxamides with significant in vivo antiobesity activity related to CB1 receptor antagonism: Synthesis, biological evaluation, and molecular modeling in the homology model. *J. Med. Chem.* 2007; 50: 5951-5966.

Received 9th September 2012

Revised 11th October 2013

Accepted 18th October 2013

J. App. Pharm. Res., 1 (1); 2013: 5 – 13