

FEATURES OF CLINICAL COURSE AND IMMUNE REGULATION OF THE HEMOSTATIC SYSTEM IN CHILDREN WITH ACUTE BRONCHOPULMONARY DISEASES WITH VARIOUS ECOLOGICAL RESIDENCE CONDITIONS

**T.A. Bobomuratov¹,
U.U. Yusupova¹,**

¹Tashkent Medical Academy, Tashkent, Uzbekistan

G.M.Djumaniyazova².

² Urgench branch of Tashkent Medical Academy, Urgench, Uzbekistan

Article history:		Abstract:
Received:	April 12 th 2021	To learn the features of clinical course, condition of homeostatic system and the interleukin in children with acute bronchopulmonary diseases with different environmental living conditions
Accepted:	April 26 th 2021	
Published:	May 21 th 2021	
Keywords: Acute pneumonia, bronchitis, interleukins, hemostatic system, correlation relationship.		

MATERIAL AND METHODS:

We examined 50 children with acute community-acquired pneumonia and 44 children with acute bronchitis (group I) living in the Aral Sea region (within the territory Khorezm) and 40 children with acute community-acquired pneumonia and 30 children with acute bronchitis aged 1 to 3 years living in the Samarkand region (group II). The diagnosis is made on the basis of an anamnesis, clinical signs, the results of instrumental and laboratory studies. We were determined concentrations of the cytokines TNF- α , IL-1 β , IL-6, IL-8, IFN γ in the serum, and studied the hemostatic system.

RESULT:

Patients living in ecologically unfavorable regions like khorezm are often accompanied by a complicated course of acute pneumonia and bronchitis, complications are background diseases, in 55,5% of cases two or more background diseases occur. The level of cytokines and indicators of the hemostatic system depended on the severity of the disease. In patients of group I in the acute period of the disease, the levels of pro-inflammatory cytokines IL-1 β , IL-6, IL-8 and TNF- α were higher than in patients of group II. The decrease in the IFN γ content mainly occurred in patients of group I. The level of interleukins in acute pneumonia was higher than bronchitis, 4.1 times in acute pneumonia, 1.5 times higher in acute bronchitis than in control. Homeostasis indicators in patients with acute complicated pneumonia in an adverse environment In adverse conditions in acute pneumonia, patients showed higher rates of fibrinogen, indicating that the hemostasis system had a high degree of readiness for intravenous blood clotting, hypo-hypercoagulation, thrombinemia, and fibrinolysis.

Younger children are characterized by severe and complicated complications of acute pneumonia and bronchitis, as well as compensatory hypercoagulation in the hemostasis system, subcompensative hypercoagulation, decompensation hypercoagulation (hypercoagulation hypocoagulation, various hemorrhages).

To identify the relationship between cytokines and the hemostatic system, a correlation analysis was performed: the average positive correlation between IL-6, TNF- α and RFMC ($r = 0.32$; $r = 0.49$) and between IL-6, IL-1 β and fibrinogen concentration ($r = 0,42$). These changes in acute pneumonia in young children indicate the presence of thrombogenic risk, which in turn requires effective correction and prevention of hemostasis.

Despite significant progress in the study of etiology, epidemiology, diagnosis and treatment, acute respiratory diseases occupy first place in the structure of morbidity in children and third in the structure of infant mortality. And today, complications, relapses, long duration and chronic respiratory diseases are common [O.V. Zaitseva, 2015; G.A. Samsygina 2015].

Recently, due to significant environmental pollution, the problem of the environment and its impact on the health of the child has become one of the most relevant. In children living in ecologically unfavorable regions, acute bronchopulmonary diseases have their own peculiar course. These are those children who are accompanied by frequent relapses of the disease and a complicated course of respiratory diseases. Such children, constituting a risk group, attract attention as not only a medical, but also a socio-economic problem [V.F. Uchaykin, 2009, A.A. Baranov 2003].

Researchers draw attention to the study of the immune regulation of physiological functions. In this regard, the hemostatic system is no exception. It is known that immunocompetent cells carrying receptors for the active factor of

blood coagulation are able to realize afferent. In the literature there are data on the effect of cytokines on vascular-platelet hemostasis, blood coagulation and fibrinolysis [Balud V.P. 1995]. It is also known that coagulation factors are able to affect the production of certain cytokines (Jones A., Greczy C. L., 1990; Vase M. et al., 1996). However, the study of the mechanism of action of cytokines in the field of medicine practically did not affect the underlying problems of the study of the hemostatic system. Simultaneously, the study of the links between immunogenesis and hemostasis with the direct participation of interleukins allows us to understand at the molecular-cellular level the mechanisms of regulation of blood coagulation and fibrinolysis in the development of DIC and other disorders of the hemostasis system in acute respiratory diseases.

In addition, a new direction in the study of this pathology is the study of clinical features and the role of cytokines in the regulation of the hemostatic system of the ADBL system in children with different environmental living conditions. Knowledge of these features would expand the information on the pathogenesis of acute bronchopulmonary diseases. And they would also allow us to develop new ways of predicting, diagnosing, treating disorders in the hemostatic system, which often occur in acute bronchopulmonary diseases in ecologically unfavorable regions.

PURPOSE OF THE STUDY:

To evaluate the features of the clinical course and hemostasis system in children with acute bronchopulmonary diseases with different environmental living conditions.

MATERIALS AND METHODS:

We examined 42 children with acute community-acquired pneumonia and 34 children with acute community-acquired pneumonia and 40 children with acute community-acquired pneumonia and 30 children with acute bronchitis aged 1 to 3 years living in the Samarkand region (group II). The examined children were at a stationary distance.

The diagnostic algorithm for verifying pneumonia and bronchitis is based on generally accepted clinical symptoms and according to the Protocol for the treatment of pneumonia and bronchitis in children. Pneumonia was diagnosed in the presence of a complex of symptoms of general intoxication, fever, catarrhal manifestations (productive cough), respiratory failure (shortness of breath, assisted muscles in breathing, acrocyanosis, etc.), percussion (local shortening of percussion tone) and auscultatory (hard or bronchial crepitus, asymmetric moist finely bubbling rales) changes in the lungs. Also confirmed by the presence of pulmonary infiltrates according to x-ray studies of the chest in a straight line, if necessary - in a lateral projection.

Hemostasis systems included determination of prothrombin time (RT), fibrinogen concentration (FIB), partial thromboplastin time activity (APTT), prothrombin index (RTI) on an apparatus (HUMAN CLOT DUO plus), platelet count using an analyzer - "Couiter MD", coagulability blood by Fonio.

The concentration of cytokines TNF- α , IL-1 β , IL-6, IL-8 in blood serum was carried out by the method of enzyme-linked immunosorbent assay using the ELISA test system IFA-TNF- α , IL-1 β , IL-6 (Vector -Best ", Russia, 2009), the level of interferon γ (IFNu) - by enzyme-linked immunosorbent assay. The study of various parts of the hemostatic system and the concentration of cytokines was carried out upon admission and before discharge of the patient (if necessary, more often).

Statistical processing of the research results was carried out using modern computing systems such as IBM using the standard software package "Excel". To identify the relationships between the analyzed indicators, a correlation analysis was carried out using the correlation coefficient r and checking its significance using the t-student and Pearson χ^2 criteria.

THE RESULTS OF THE STUDY:

The results of the study showed that the high incidence of complicated forms of pneumonia was observed in children of group I (81%), compared with group II (25%) living in the Samarkand region. In patients of group I, the most frequent complications revealed carditis, which was determined in 45% (19), broncho-obstructive syndrome was detected in 26% (11) and DIC syndrome 9.5% (4). Whereas in patients of group II carditis was detected 3 (7.5%), bronchial obstructive syndrome was detected in 6 (15%). When analyzing cases of bronchitis, acute obstructive bronchitis was often detected in 24 (71%) patients in groups I, recurrent course was detected in 14 (41.2%), while acute obstructive bronchitis was found in patients in group II patients (6 (20%), recurrent 4 course (13.3%).

The main risk factors for the development of community-acquired pneumonia and acute bronchitis in patients living in averse zones was premorbid background. The data obtained showed that in the stationary group of children with 58% (76.3%). At the same time, 53 (69.7%) children were diagnosed with rickets, 43 (56.6%) children had protein-energy deficiency of BEN, anemia of 68 (89.5%) children, 2 and 3 concomitant diseases occurred in 42 (55, 3%) children of the first group. In the second group, the presence of concomitant diseases was detected in 42 (60%) children. The diagnosis of rickets was determined in 28 (40%) children, BEN, in 32 (45.7%) children, the diagnosis of anemia was detected in 57 (81.4%) children. In this case, 2 or more concomitant diseases in children of the second group were detected in 34 (48.6%) children.

The results of the study showed that both in group I and group II in the acute period of the disease in the cytokine status, as well as in the hemostatic system, significant changes occur. Of greatest interest are the results of

studies of the cytokine system. Changes in cytokine levels characteristic of varying severity corresponded to the volume of lung tissue lesion.

Tab. 1. Indicators of cytokine status in patients of group I and II ($M \pm m$)

Indicators	Acute pneumonia I group (n=42)	Acute pneumonia II group (n=40)	Bronchitis I group (n=34)	Bronchitis II group (n=30)	Healthy children (n=30)
IL-1 β , ng/ml	186,0 \pm 6,9***	141,9 \pm 5,6***	72,6 \pm 3,8*	54,6 \pm 3,4*	44,8 \pm 2,8
FNO- α , ng/ml	204,8 \pm 9,0***	159,8 \pm 8,9***	88,5 \pm 2,7***	59,8 \pm 2,2***	54,5 \pm 2,4
IL-6, ng/ml	24,1 \pm 2,1***	15,2 \pm 2,2**	13,8 \pm 0,5*	8,2 \pm 0,6	6,5 \pm 0,2
IL-8, ng/ml	109,6 \pm 4,2***	84,3 \pm 2,2***	79,3 \pm 2,4**	63,3 \pm 2,2**	54,3 \pm 3,3
INF γ , ng/ml	21,2 \pm 1,3***	44,0 \pm 1,7***	32,1 \pm 2,4***	64,1 \pm 2,8**	86,2 \pm 4,6

Note. The statistical significance of the difference with the indices of the control group: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

As can be seen from the table. 1, in patients of group I in the acute period of the disease, the levels of pro-inflammatory cytokines IL-1 β , IL-6, IL-8 and TNF- α were higher than in patients of group II. The decrease in the INF γ content mainly occurred in patients of group I. Mostly occurred in patients of group I. The level of damage in the bronchopulmonary system, 4.1 times with acute pneumonia, 1.5 times higher with acute bronchitis. Normalization during the period of clinical remission in patients of group I does not occur. The high level of pro-inflammatory cytokines observed in the period of clinical remission causes less adaptive capabilities of these children. At the beginning and in the midst of acute pneumonia, a normalization of the inflammatory activity of IL-8 was detected in both groups. However, in patients with a severe form of the disease, the value of regulatory IL-6 remained significantly high. In patients of group I, who have frequent complications, more pronounced changes occurred (Table 1).

Thus, a change in the cytokine status, a severe, often complicated course of acute pneumonia and bronchitis in the period of clinical remission in group I patients, causes inflammation to persist, preservation of the infectious agent, and during clinical remission.

Laboratory indicators of hemostasis upon admission to the hospital in patients of groups I and II are presented in table 2.

Tab. 2. Indicators of hemostasis in patients of group I ($M \pm m$)

Criterion	Acute pneumonia I group (n = 32)	Acute pneumonia I group (n = 32)	Bronchitis I group (n = 24)	Bronchitis II group (n = 30)	Healthy children group (n = 30)
Platelets ($\times 10^9/l$)	379 \pm 15,3	330 \pm 15,9	234 \pm 8,3	239 \pm 11,6	205,0 \pm 16. 9
AChTV (sec)	47,7 \pm 1,9*	33,9 \pm 2,1	33,8 \pm 1,9	33,4 \pm 1,9	33 \pm 0,7
TV (sec)	9,7 \pm 1,1*	9,8 \pm 1,1	9,3 \pm 1,2	9,5 \pm 1,4	17,2 \pm 1,2
PTV (sec)	15,7 \pm 1,2*	14,7 \pm 1,2	14,4 \pm 1,3	14,4 \pm 1,23	16,8 \pm 2,5
Fibrinogen (g / l)	4,49 \pm 0,32*	3,9 \pm 0,32	2,9 \pm 0,37	3,29 \pm 0,34	2-4 г/л

Note. M - arithmetic mean; m is the error of the average value; n is the number of patients; Statistical significance of the difference with the indices of the control group: * - significant intergroup differences ($p \leq 0.05$; $p \leq 0.01$).

In acute pneumonia, the inflammatory process cannot be described as separate, without changes in the hemostasis, since the effects of the clotting and coagulation system are of particular importance in the pathogenesis of the disease. Analysis of laboratory observations of our observations shows that, especially in group 1, patients had increased fibrinogen content (4.49 + 0.32g/l), increased platelet time (AChTV 47.7 + 1.9 sec), increased platelet count, prothrombin index, and time was reduced. Normal or significant changes were not detected in 9% of the patients analyzed, 22% had compensatory hypercoagulation, 55% had subcompensated hypercoagulation, 14% had decompensated hypercoagulation (hypercoagulation hypocoagulation, various hemorrhages). Changes in the system of hemostasis in acute pneumonia are more pronounced than in uncomplicated forms, and a dramatic increase in platelet count is the compensatory response of platelets to the occurrence of intracranial clotting. It has also been shown that a number of coagulation joints of hemostasis in complicated pneumonia are at high thrombotic readiness

for intravenous coagulation. Failure of timely correction of hypercoagulation syndrome in sick children leads to the worsening of the general condition of patients and the decompensation of hypercoagulation.

In acute armor in different groups of changes in coagulation hemostasis, significant hypercoagulation (AChTV 34 ± 2.1 ; 33.8 ± 1.9) are distinguished against the background of inhibition of the final stage of coagulation (TB 9.9 ± 1.1 ; 9.3 ± 1 , 2), fibrinogenemia (fibrinogen 3.2 ± 0.34 ; 2.9 ± 0.37), fibrinemia (127.2 ± 5.3 ; 118.2 ± 4.3), inhibition of fibrinolysis and an increase in blood levels of products fibrin degradation. The revealed changes are due to severe thrombinemia (199.2 ± 12.3 ; 134.9 ± 4.7) and activation of intravascular coagulation to a greater extent. decompensation of the current intravascular coagulation. All this is an indication for the correction of the blood coagulation system.

To make the relationship between cytokines and the hemostatic system, we conducted a correlation analysis of their indicators. The greatest effect in the pathogenesis of acute bronchopulmonary diseases is characteristic of IL-6 and IL-8, which are important triggers of hypercoagulation [Kuznik BI, 2005]. In children of group I (vulnerable zone), a high positive correlation was observed between the levels of IL-6, IL-1 and fibrinogen concentration ($r = 0.42$), compared with group II ($r = 0.48$). This can be explained by the fact that a high level of pro-inflammatory cytokines, especially IL 6, acting on hepatocytes, stimulate the cleavage of proteins of the acute phase (α_1 – antitrypsin, α_2 – macroglobulin and fibrinogen) [Gusev E.Yu., 2008; Ketlinsky S.A. 2008.]. An average positive correlation was observed between the content of IL-6, FNO- α and RFMC ($r = 0.32$; $r = 0.49$), which reflects hypercoagulation and a decrease in fibrinolytic activity in the examined patients.

Between IFNu and indicators of the hemostatic system revealed a close negative correlation. This can be explained by the "immune mechanisms of hemostasis regulation." According to this theory, how low the number of immunocompetent cells will be, so the coagulation ability of the blood will be prone to increase [Kuznik BI, 2005].

In the body there is a single integrated cellular-humoral defense system, including immunity, hemostasis and nonspecific resistance of the body (A.Yu. Vitkovsky, 1997). Changes in one of the functional links of this system entail shifts in others.

Thus, in acute bronchopulmonary diseases in children with complicated forms of acute bronchopulmonary diseases, there was a hemostasis reaction in the form of thrombinemia, inhibition of fibrinolysis. A change in the cytokine status and hemostasis system may be one of the factors that determines the complicated and recurrent course of the pathological process. These changes were more pronounced in children, patients living in the Aral Sea region (in the Khorezm region) compared with those living in the Samarkand region.

CONCLUSIONS:

1. Background diseases play a significant role in the development and progression of acute bronchial lung diseases in children, and children living in the Khorezm region, where environmental conditions are most common (76.3%), have two or more background diseases in 55.3%. These background diseases cause severe and prolonged development of acute pneumonia and bronchitis in children, complications, profound changes in the immune system and hemostasis.
2. In the acute period of bronchopulmonary diseases, the level of cytokines and indicators of the hemostatic system depended on the severity of the disease:- in all patients, the level of pro-inflammatory cytokines IL-1 β , IL-6, IL-8 and FNO- α increases and the concentration of interferon in both groups decreases, but to a greater extent in children more living in the Khorezm region compared to healthy children.
3. Younger children are characterized by severe and complicated complications of acute pneumonia and bronchitis, as well as compensatory hypercoagulation in the hemostasis system, subcompensative hypercoagulation, decompensation hypercoagulation (hypercoagulation hypocoagulation, various hemorrhages). In adverse conditions in acute pneumonia, patients showed increased hemostasis, mainly with increased readiness for intravenous blood clotting, increased fibrinogen levels, decreased hypercoagulation, thrombinemia, and fibrinolysis on AChTV. These changes in acute pneumonia in young children indicate the presence of thrombogenic risk, which in turn requires effective correction and prevention of hemostasis.
4. The established correlation between cytokines and hemostasis indicates immune regulation of blood coagulation. Changes in one of the functional links of this system entail shifts in others. At the same time, cytokines are information molecules that adapt protective mechanisms to maintain homeostasis of the body.

LITERATURES (List of published works)

1. Baranov A.A., Albitsky V.Yu. Social and organizational problems of pediatrics. M .: Dynasty. 2003; 233-252. [Baranov A.A., Al'bitskiy V.Yu. Sotsial'nye i organizatsionnye problemy pediatrii. (Social and organizational problems of Pediatrics.) Moscow: Dinastiya. 2003; 233-252. (In Russ.)]
2. Bobomuratov T.A. Studying the hemostatic system and evaluating the effectiveness of lymphotropic anticoagulant therapy in young children with acute pneumonia. Lymphology Andijan. 2000; 1-2: 9-13.
3. Gusev E.Yu., Yurchenko L.N., Chereshev V.A. et al. Options for the development of acute systemic inflammation. Cytokines and inflammation. 2008; 7 (2): 9–17. [Gusev E.Y., Yurchenko L.N., Chereshev V.A. et al. The variants of acute systemic inflammation evolution. Tsitokiny i vospalenie. 2008; 7 (2): 9–17. (In Russ.)]

4. Zaitseva OV Recurrent respiratory infections: is it possible to prevent? Pediatrics. 2015; 94 (2): 185-192. [Zaytseva O.V. Recurrent respiratory infections: is it possible to prevent them? Pediatriya. Zhurnal im. G.N. Speranskogo. 2015; 94 (2): 185-192. (In Russ.)]
5. Ketlinsky S.A., Simbirtsev A.S. Cytokines. SPb .: Tome. 2008; 552 p. [Ketlinskiy S.A., Simbirtsev A.S. Tsitokiny. (Cytokines.) St. Petersburg: Foliant. 2008; 552 p. (In Russ.)]
6. Kuznik B. I., Tsybikov N. N. Autoimmune mechanisms of regulation of the hemostatic system. Siberia onkol. g. 2005; 13 (1): 88-95. [Kuznik B.I., Tsybikov N.N. Autoimmune mechanisms of the hemostatic system regulation. Siberian onkologicheskij zhurnal. 2005; 13 (1): 88-95. (In Russ.)]
7. Mamatkulova D.Kh., Bobomuratov T.A. Clinical and pathogenetic significance of changes in hemostatic disorders and their correction in chronic pneumonia in children. Medical sciences. 2004; 2: 29-30.
8. Samsygina G.A. The problem of often ill children in pediatrics. Pediatrics. 2015; 94 (1): 167-169. [Samsygina G.A. The issues of frequently ill children in pediatrics. Pediatriya. Zhurnal im. G.N. Speranskogo. 2015; 94 (1): 167-169. (In Russ.)]
9. Uchaykin V.F. Recurrent respiratory infections in children: the use of immunomodulators for treatment and prevention. Pediatrics. 2009; 87 (1): 127-132. [Uchaykin V.F. Recurrent respiratory infections in children: the use of immunomodulators for the treatment and prevention. Pediatriya. Zhurnal im. G.N. Speranskogo. 2009; 87 (1): 127-132. (In Russ.)]
10. Arai K. Cytokines: coordinators of immune and inflammatory responses. Ann. Rev. Biochem. 2006; 59: 783.
11. Bellanti J. Recurrent respiratory tract infection in pediatric patient. Drugs 2006; 54 (suppl. 1): 1-4.