

# Effects of Temperature, Relative Humidity, and DEN-2 Virus Transovarial Infection on Viability of *Aedes aegypti*

## Pengaruh Temperatur, Kelembaban Relatif, dan Infeksi Transovarial Virus DEN-2 terhadap Viabilitas *Aedes aegypti*

Tri Baskoro T. Satoto\* Sitti R. Umniyati\* Adi Suardipa\*\* Margareta M. Sintorini\*\*\*

\*Pusat Kedokteran Tropis Fakultas Kedokteran Universitas Gadjah Mada, \*\*Puskesmas Naioni Kupang, \*\*\*Departemen Teknik Lingkungan Universitas Trisakti

### Abstract

Environmental changes influenced survival life and virus transmission of dengue virus (DEN) in a mosquito. The purpose of the present study was to define DEN-2 virus transmission dynamic and effect of temperature, relative humidity (RH), and DEN-2 virus infection on viability of *Aedes aegypti* (*Ae. aegypti*). This experimental study with pretest-posttest control group design was conducted at the Laboratory of Center for Tropical Medicine, Faculty of Medicine, Gadjah Mada University (UGM), Yogyakarta. Seventh-days old female *Ae. aegypti* (F0) were infected DEN-2 via oral membrane and kept until F2 generation by transovarial transmission, number of eggs produced and hatched was recorded. After 14-day incubation was found that transovarial transmission rate of DEN-2 virus infection in F0 and F1 were 93.3% and 82.2%, respectively. Egg production, hatching rates from infected and uninfected mosquitoes F0 were 68% and 85%; and F1 were 72.6% and 76%, respectively. At defined room condition tests, 7 day adult mosquitoes in dark and humid environment produced highest number of eggs, compared normal environment and in incubated without CO<sub>2</sub>. In fourteenth day old mosquitoes at dark and humid produced highest number of eggs, compare normal environment condition, and in incubated without CO<sub>2</sub>. DEN-2 virus was able to infect *Ae. aegypti* by transovarial transmission where the infection rate in F0 was higher than F1 generation. Temperature and humidity affected the ability of *Ae. aegypti* eggs to live and grow to adulthood.

**Keywords:** *Aedes aegypti*, DEN-2 virus, humidity, temperature, transovarial

### Abstrak

Perubahan lingkungan memengaruhi hidup dan transmisi virus dengue dalam tubuh nyamuk. Tujuan penelitian ini adalah mengetahui pengaruh suhu, kelembaban udara (RH), terhadap transmisi virus DEN-2 pada nyamuk *Aedes aegypti*. Studi eksperimental dengan desain *pre* dan *post test control* group dilakukan di laboratorium pusat kedokteran tropis, Fakultas Kedokteran, Universitas Gadjah Mada, Yogyakarta. Penelitian ini dilakukan

pada kelompok *Ae. aegypti* betina umur 7 hari (F0). Virus DEN-2 diinfeksi secara transovarial cara membran oral sampai generasi F2. Kelompok lain sebagai kontrol di inkubator temperatur dan suhu tertentu, waktu tertentu, jumlah telur yang dihasilkan, yang menetas dan mengandung virus dicatat. Hasil penelitian menemukan indeks transmisi transovarial generasi F0 dan F1 selama 14 hari masa inkubasi adalah 93,3% dan 82,2%, laju tetas telur dari nyamuk F0 yang terinfeksi dan tidak terinfeksi masing-masing 68% dan 85%, sedangkan laju tetas telur dari nyamuk F1 yang terinfeksi dan tidak terinfeksi masing-masing 72,6% dan 76%. Pada tiga kondisi ruang uji, nyamuk berumur 7 hari dalam ruang gelap dan lembab menghasilkan telur paling banyak dibandingkan pada kondisi normal dan pada inkubasi tanpa CO<sub>2</sub>. Nyamuk umur 14 hari menghasilkan telur tertinggi dalam ruang gelap dan lembab, dibandingkan pada kondisi ruang normal dan dalam inkubasi tanpa CO<sub>2</sub>. Virus DEN-2 dapat menginfeksi *Ae. aegypti* secara transovarial dengan laju infeksi lebih tinggi pada F0 daripada F1. Suhu dan kelembaban mempengaruhi kemampuan produksi telur *Ae. aegypti* untuk hidup dan tumbuh.

**Kata kunci:** *Aedes aegypti*, virus DEN-2, kelembaban, suhu, transovarial

### Introduction

Dengue has become a global health problem, particularly in tropical and subtropical countries. As such, global warming along with climate changes significantly affects the mechanism of dengue transmission. Changes in air temperature and relative humidity may reduce the viability of Dengue virus (DEN) in the mosquito's body.<sup>1</sup> To maintain its existence in nature, DEN virus has two mechanisms i.e. by horizontal transmission between ver-

Alamat Korespondensi: Tri Baskoro T. Satoto, Pusat Kedokteran Tropis Fakultas Kedokteran Universitas Gadjah Mada, Jl. Farmako Sekip Utara, Yogyakarta 55281, Hp. 0816680974, e-mail: tribaskorots@yahoo.com

tebrate viremia which is transmitted by *Aedes* mosquitoes and by vertical transmission (transovarial) from the next generation of infective female.<sup>2</sup> Of these transmissions, transovarial mechanism is an important aspect in maintaining the viability of DEN virus during the inter-epidemic period in nature.

There are four types dengue virus namely DEN-1, DEN-2, DEN-3, and DEN-4, one of most potent is DEN-2 virus has been isolated from larvae of *Ae. aegypti* in Yangon, Myanmar, and was expected that transmission occurs by vertical mechanism.<sup>3</sup> Transovarial transmission can take place until the 7<sup>th</sup> generation of mosquitoes that have been infected by DEN virus parentally.<sup>4</sup> Eggs from infective female mosquitoes, which hatched after an incubation period of several weeks at room temperatures, increased the percentage of transovarial transmission.<sup>5</sup> At temperature ranges from 12 to 36°C, with maximum and minimum limits of 35°C and < 12°C, respectively, egg viability was greater than 80%.<sup>6</sup>

The existence of this nature phenomenon is important for *Aedes* mosquitoes in supporting and maintaining the vector endemic capacity of a region. This indirectly affects the incidence of dengue in the community which shows a complete Dengue transmission mechanism. So, for better understanding of the transmission mechanism, it is necessary to know the contributing components of the dynamic transmission. The purposes of this study was to prove the existence of transovarial transmission of DEN-2 virus in *Ae. aegypti*, to know the number of eggs produced by mosquitoes *Ae. aegypti* infected with DEN-2 virus, and to investigate the effect of temperature and humidity on the hatchability of eggs of transovarial DEN-2 virus-infected *Ae. aegypti* after the eggs are stored for 7 days and 14 days.

## Methods

This experimental study with pre-posttest control group design was conducted at the laboratory of Center for Tropical Medicine, Faculty of Medicine, Gadjah Mada University (UGM), Yogyakarta. Adult female *Ae. aegypti* mosquitoes aged 7 days were used as subject. The mosquitoes were infected by DEN-2 virus through oral membrane feeding (fresh rat skin that had been sheared fur).<sup>7</sup> The DEN-2 viruses used in this experiment have been colonized in *Ae. aegypti* since 2004 at the Insectariums of the Center for Tropical Medicine, UGM. DEN virus suspension consisting of C6/36 cell culture supernatant of infected mosquitoes, which was previously infected by DEN-2 virus (Hawaiian isolate) for 6 days incubation, was mixed with erythrocytes of healthy persons containing anti-coagulant heparin and sucrose 10% with a ratio of 1 : 1 : 1.12. The mixture was placed in the room at 25°C ± 3°C and rela-

tive humidity (RH) 70% ± 5%. On the third day, the mosquitoes were blood-fed on healthy rats. Further, mosquito eggs were collected from the parent (F0) of DEN-2 infected group and control group mosquitoes after 10 days incubation period (*Gonotrophic* II). Meanwhile, mosquito eggs from the progeny (F1) from the infected and control parent (F0) mosquitoes were collected after 10-day-old mosquitoes (*Gonotrophic* II). The eggs were collected by confining each individual female mosquito in paper cups covered by nylon netting on the top and bottom parts. Cotton pads, covered by wet filter paper (ovistrip) on its upper, were placed in the paper cups for egg laying. After the egg is released, the female parent was tested for the presence of DEN virus using head squash Immunocytochemical SBPC method with monoclonal DSSC7 as primary antibodies.<sup>8,9</sup>

Eggs from F1 *Ae. aegypti* with positive DEN-2 virus transmitted transovarially after 3-day embryonation were referred as group 1 (K1) and group 2 (K2) for 7-day and 14-day incubations, respectively. The K1 and K2 were then categorized into three groups of room temperature and RH: ambient condition at 25°C ± 5°C and RH 80% ± 5% (T<sub>1</sub>RH<sub>1</sub>), humid dark room at 20°C ± 5°C and RH 85% ± 5% (T<sub>2</sub>RH<sub>2</sub>), and incubated without CO<sub>2</sub> at 30°C ± 5°C and RH 70% ± 5% (T<sub>3</sub>RH<sub>3</sub>). Sample of 100 mosquitoes eggs in each group and attached to the ovistrip. The temperature and humidity were measured using thermohygrometer and were observed twice a day in the morning (07:00 to 08:00 pm), hereinafter referred to as P1, and in the afternoon (12:00 am to 1.00 pm), hereinafter referred as P2. F1 mosquito eggs, after being kept into adult mosquitoes in 3 different treatments, were colonized until F2 generation. The results were analyzed statistically with the Mann-Whitney test and ANOVA. Mean differences between treatments revealed a significant effect if the critical value  $F_0 \geq \text{critical value } F \text{ table at } \alpha = 0.05$ .<sup>10</sup>

## Results

The results of DEN-2 virus antigen test of females (F0) *Ae. aegypti*, which were infected orally by DEN-2 virus in 14 day incubation period and their progeny (F1) adult of 14 days of age are presented in Table

Table 1. Antigen Test of DEN-2 Virus in Preparations Mosquito Head Squash Parent (F0) and the Progeny (F1) Adult Stage

Mosquitoes Generation	Groups of Mosquitoes Tested	Antigen Virus DEN-2			Infection Rate (%)
		Positive	Negative	Total	
F0	Infected by DEN-2 virus	56	4	60	93.3
	Not infected	0	30	30	0
F1	Infected by DEN-2 virus	74	16	90	82.2
	Not infected	0	30	30	0

1. It reveals the DEN-2 virus infection rate in F0 and F1 generation mosquitoes. Number of eggs produced from F0 and F1 *Ae. aegypti* mosquitoes are presented in Table 2. It describes the number of hatching and not hatching eggs from infected and non infected mosquitoes.

Results of colonization of eggs from F1 mosquitoes, which were positively infected by DEN-2 virus transovarially, are presented in Table 3, whereas colonization results of uninfected F1 mosquitoes are presented in Table 4. Both tables cover 7-day and 14-day test periods for 3 different temperature and RH experimental conditions.

During laboratory colonization, the average air temperature and RH were 24.5°C and 65.5%, respectively. At this condition, eggs hatched into larvae within 1 – 3 days, larva stage lasted 8 – 12 days to become pupae, and pupal stage lasted 2 – 3 days to become adult mosquito.

## Discussion

Table 1 indicates that local strains of *Ae. aegypti*, which has been colonized since 2004 at the Center for Tropical Medicine UGM, is susceptible to DEN-2 virus infection. It was found 75% infection rates in first colony by immunocytochemistry SBPC head squash methods.<sup>9</sup> These results have good agreement with previous research that the mosquitoes were categorized as vulnerable if after infected orally the DEN virus, found positive results on brain tissue, based on head squash preparations immunofluorescence method in 2-week incubation.<sup>11</sup> Vulnerability of *Ae. aegypti* against DEN virus was associated with vertical transmission rate because higher transovarial transmission in the offspring may be caused by the high number of infected females and vulnerability.<sup>4</sup>

**Table 2. Number of Hatching and Not Hatching Eggs from DEN-2 Virus-Infected and Uninfected (Control) F0 and Progeny (F1) of *Ae. aegypti* Mosquitoes**

Mosquito Generation	Group of Test	Hatching and Not Hatching Eggs				Total	Total Eggs Produced	Average Eggs <sup>a</sup>
		Hutched (+)		Hutched (-)				
		n	(%)	n	(%)			
F0	Infected by DEN-2 virus	68	68	32	32	100	4,235	62
	Not infected	85	85	15	15	100	6,688	79
F1	Infected by DEN-2 virus	109	72.6	41	27.4	150	7,990	73
	Not infected	114	76	36	24	150	9,459	83

Notes:

<sup>a</sup>Per each mosquitos

**Table 3. Colonization Results of Eggs of Transovarially DEN-2 virus-infected F1 *Ae. aegypti* After 7-day and 14-day Treatment at 3 Different Room Temperature and Humidity Conditions.**

Test Period	Test Group	Repetition	Eggs Tested	Adult Mosquito Live		Total	Average
				Male	Female		
7-day	T <sub>1</sub> RH <sub>1</sub>	1	100	33	22	55	52
		2	100	25	28	53	
		3	100	25	23	48	
	T <sub>2</sub> RH <sub>2</sub>	1	100	40	34	74	62
		2	100	23	31	54	
		3	100	17	42	59	
	T <sub>3</sub> RH <sub>3</sub>	1	100	28	17	45	42
		2	100	19	22	41	
		3	100	18	21	39	
14-day	T <sub>1</sub> RH <sub>1</sub>	1	100	14	20	34	36
		2	100	12	27	39	
		3	100	10	25	35	
	T <sub>2</sub> RH <sub>2</sub>	1	100	23	21	44	46
		2	100	15	32	47	
		3	100	22	26	48	
	T <sub>3</sub> RH <sub>3</sub>	1	100	22	18	40	40
		2	100	23	14	37	
		3	100	19	23	42	

Notes:

T<sub>1</sub>RH<sub>1</sub> = Normal room conditions (temperature 25 ± 5°C, relative humidity (RH) 80 ± 5%); T<sub>2</sub>RH<sub>2</sub> = Dark and humid room (temperature 20 ± 5 °C, RH 85 ± 5%); T<sub>3</sub>RH<sub>3</sub> = incubated without CO<sub>2</sub> (temperature 30 °C ± 5°C, RH 70% ± 5%)

**Table 4. Colonization Results of Eggs of Uninfected F1 *Ae. aegypti* After 7-day and 14-day Treatment at 3 Different Room Temperature and Humidity Conditions**

Test Period	Test Group	Repetition	Eggs Tested	Adult Mosquito Live		Total	Average
				Male	Female		
7-day	T <sub>1</sub> RH <sub>1</sub>	1	100	51	39	90	79
		2	100	45	33	78	
		3	100	21	48	69	
	T <sub>2</sub> RH <sub>2</sub>	1	100	46	42	88	86
		2	100	47	36	83	
		3	100	52	34	86	
	T <sub>3</sub> RH <sub>3</sub>	1	100	31	37	68	72
		2	100	41	31	72	
		3	100	3	36	75	
14-day	T <sub>1</sub> RH <sub>1</sub>	1	100	32	36	68	58
		2	100	34	25	59	
		3	100	28	19	47	
	T <sub>2</sub> RH <sub>2</sub>	1	100	43	29	72	66
		2	100	41	21	62	
		3	100	27	37	64	
	T <sub>3</sub> RH <sub>3</sub>	1	100	23	34	57	59
		2	100	19	35	54	
		3	100	36	29	65	

## Notes:

T<sub>1</sub>RH<sub>1</sub> = normal room conditions (temperature 25 °C ± 5°C and relative humidity (RH) 80% ± 5%); T<sub>2</sub>RH<sub>2</sub> = dark and humid room at temperature 20 °C ± 5 °C and RH 85% ± 5%; T<sub>3</sub>RH<sub>3</sub> = incubator without CO<sub>2</sub> at temperature 30 °C ± 5°C and RH 70% ± 5%).

Mosquitoes are more vulnerable to viruses due to variation in inter (genetic) and between species in inhibiting the infection and spreading out of the virus leading to differences in the infection rate and transmission. In susceptible mosquitoes, the titer will increase, possibly due to replication in secondary target organs, while the mechanism of vertical transmission of arbovirus in the mosquito's body will produce progeny with infection rates exceeding 80%.<sup>11</sup>

Susceptibility of *Ae. aegypti* against DEN-2 virus infection provides good mechanism for the survival of the virus and mosquitoes in nature. The present study shows that the control group mosquitoes produced more eggs than the infected ones. As shown in Table 2, the uninfected parent (F0) *Ae. aegypti* produced 6,688 eggs from 85 mosquitoes (79 eggs per mosquito in average). This figure was significantly different ( $p = 0.001$ ) with infected F0 which produced only 4,235 eggs from 68 mosquitoes (62 eggs per mosquito in average). The same figure was found in the progeny (F1), that the uninfected mosquitoes significantly produced higher eggs (9,459 eggs) than the infected ones (7,990 eggs) ( $p = 0.009$ ).

In one gonotrophic cycle, a female *Ae. aegypti* is able to produce as many as 100 to 400 eggs.<sup>12</sup> Based on this figure, the number of eggs and mean number of eggs produced by F0 and F1 mosquitoes of both infected and uninfected were lower (Table 2). These possibly due to room temperature and humidity and lack of food intake (blood) which is a source of protein for growth and de-

velopment of eggs. In the life of mosquito after feed the blood to be continued produce eggs, thus the number of eggs were deposited. The difference between the control (uninfected) and infected mosquitoes probably due to the effects of DEN-2 virus infection during embryogenesis. Virus multiplication in different organs during embryogenesis or in the final stages of mosquito may vary due to the tissue tropism, descendant of the virus, and host genetic creation. These factors, either alone or together, can contribute to the duration of larvae, productivity, fertility, and mortality is different in the infected mosquito transovarium.<sup>4</sup>

Temperature and humidity have significant effect on *Ae. aegypti* in producing eggs. As shown in Table 3 and Table 4, infected and uninfected F1 mosquitoes kept 7 days in dark and humid room (T<sub>2</sub>RH<sub>2</sub>: 22 – 26°C, RH 87 – 92%) produced in average 62 and 86 eggs, respectively, contrast to 52 and 79 eggs at ambient condition (T<sub>1</sub>RH<sub>1</sub>: 25.5 – 28°C, RH 79 – 87%) and 42 and 72 eggs in incubator without CO<sub>2</sub> (T<sub>3</sub>RH<sub>3</sub>: 33 – 35°C, RH 68 – 72%). The same pattern was observed in 14-day period. These results show a decline number of eggs hatching and growing into adult mosquitoes live where 14-day treatment was worse than 7-day treatment.

The effect of temperature, humidity, and DEN-2 virus infection was observed in the offspring of *Ae. aegypti*. As shown in Table 3 and Table 4, the number of survival eggs hatching into adult mosquitoes was different among temperature and humidity conditions.



The low hatchability of eggs incubated without CO<sub>2</sub> (T<sub>3</sub>RH<sub>3</sub>) was likely influenced by temperature and humidity of the incubator. During the treatment, the average incubator temperature (34.5°C) was above the optimum temperature (20 – 30°C) at which generally mosquitoes lay their eggs. The optimum temperature for growth and developments of mosquitoes is 25 – 27°C, and the mosquito growth stops when temperature is lower than 10°C or higher than 40°C.<sup>11</sup>

Mosquito eggs are easily broken and some embryonic processes are crashed in condition not optimum or placed in a moistened environment. Those reared in a high moisture environment hatched at a high rate compared with their to a drier environment.<sup>13</sup> In the present study, the average air RH in all three treatment groups ranged 69.8 – 90.3%. This was within the average range of optimum humidity for mosquito development of 70 – 90% and, therefore, air humidity during treatment did not affect much on the growth and development of mosquito eggs. The experimental condition of room humidity was optimum for the embryonic processes and embryo survival resulting in longer mosquito lifespan.<sup>11</sup>

DEN virus affected to the ability of life and growth and development of progeny. It was indicated by the difference of the number of hatching eggs into adult mosquitoes. As shown in Table 2, hatching eggs of infected mosquitoes (68% of F0 and 72.6% of F1) were smaller than the uninfected ones (85% of F0 and 76% of F1). Previous research showed that hatching rate of eggs from DEN-3 virus mosquitoes was only 30 – 68.1%.<sup>14</sup>

In the present study, factors associated with pathogenesis of DEN-2 virus in mosquito system or other factors cannot be explained because the presence of DEN antigen and viral titer of egg, larva, and pupa were not checked. However, it is known that the pathogenesis of the virus also relies on the offspring, where less adaptable offspring causes higher pathogenesis. Meanwhile, mosquitoes infected with DEN-3 virus showed higher mortality compared with control mosquitoes. This differs from other finding that classically arbovirus has no disturbing effect on the vector.<sup>11</sup> As a result, DEN virus breeding in cells of *Ae. albopictus* does not always cause CPE (cytopathogenic effect) so that the infection is persistent. This may be analogous to the presence of natural DEN-2 virus in *Ae. aegypti* where the virus replicates did not cause the mosquito dead due to the absence of CPE.

Effect of DEN-2 virus transmitted transovarially can be regarded as one of biological controlling factors over the reproductive potency of *Ae. Aegypti*. *Ae. aegypti* eggs, which still survive and hatch up to become adult mosquitoes, is probably the best mosquito progeny of genetic superiority and important transovarial dynamic transmission of dengue for the next generation.

## Conclusions

The DEN-2 virus was able to infect *Ae. aegypti* by transovarial transmission, where infection rate in F0 was higher than that in F1 generation. In both generations, DEN-2 virus-infected *Ae. aegypti* mosquitoes produced smaller number of eggs than those of uninfected mosquitoes. Temperature and humidity affected the ability of *Ae. aegypti* eggs to live and grow to adulthood, when longer storage resulted in smaller survival rates.

## Acknowledgment

We thanks to Dr Sustriayu Nalim, National Institute for Health and Research for her excellent reviewed of the manuscript. Special thanks to Mr Purwono, Mr Heru Sudibyo, Mr. Joko Trimuratno and Ms Kiswati, laboratory of Parasitology Faculty of Medicine Gadjah Mada University for their technical help.

## References

1. Zhang Y, Bi P, Hiller JE. Climate change and the transmission of vector-borne diseases: a review. *Asia Pacific Journal of Public Health*. 2008; 20(1): 64-76.
2. Halstead SB. Dengue. In: Warren KS, Mahmoud, eds. *Tropical and geographical medicine*. London: Imperial College Press; 1990. p. 675-85.
3. Angel B, Joshi V. Distribution and seasonality of vertically transmitted dengue viruses in *Aedes* mosquitoes in arid and semi-arid areas of Rajasthan, India. *Journal of Vector Borne Disease*. 2008; 45: 56-9.
4. Joshi V, Mourya DT, Sharma RC. Persistence of Dengue-3 Virus through transovarial transmission passage in successive generations of *Aedes aegypti* mosquitoes. *American Journal of Tropical Medicine and Hygiene*. 2002; 67: 158-61.
5. Mourya DT, Joshi V. Horizontal and vertical transmission of dengue virus in *Aedes aegypti* mosquitoes and its persistence in successive generations. The 6th International Symposium on Vector and Vectors Borne Diseases. 2002 Feb 9-11, Bhubaneswar, India. National Academy of Vector Borne Diseases & Regional Medical Research Centre, Bhubaneswar: 2002.
6. Farnesi LC, Martins AJ, Valle D, Rezende GL. Embryonic development of *Aedes aegypti* (Diptera: Culicidae): Influence of different constant temperatures. *Memorias do Instituto Oswaldo Cruz*. 2009; 104 (1): 124-6.
7. Graves PM. Studies on the use of membrane feeding technique for infecting anopheles gambiae with *Plasmodium falciparum*. *Transaction of the Royal Society of Tropical Medicine and Hygiene*. 1980; 74: 738-42.
8. Medina F, Medina JF, Colón C, Vergne E, Santiago GA, Muñoz-Jordán JL. Dengue virus: isolation, propagation, quantification, and storage. *Curr Protoc Microbiol* [serial on the internet]. 2012; 15: 150 [cited 2012 Nov 10]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23184594>.
9. Umniyati SR, Sutaryo, Wahyono D, Artama WT, Mardihusodo SJ, Soeyoko, et al. Application of monoclonal antibody DSSC7 for detecting dengue infection in *Aedes aegypti* based on immunocytochemical streptavidin Biotin Peroxidase Complex Assay (ISBPC). *Dengue Bulletin*. 2008; 32: 83-98.

10. Foster JJ. Data analysis using SPSS for widows versions. London: SAGE Publication Ltd; 2002.
11. Beaty BJ, Jennifer LW, Higgs S. Natural cycles of vector-borne pathogens. In: Beaty BJ, Marquardt WC, eds. The Biology of Disease Vectors. Niwot Colorado: University Press of Colorado; 1996. p. 51-70.
12. Clements NA. The biology of mosquitoes. New York: CABI Publishing; 1999 .p. 619- 26.
13. Dieng H, Boots M, Tamori N, Higashihara J, Okada T, Kato K, Eshita Y. Some technical and ecological determinants of hatchability in *Aedes albopictus*, a potential candidate for transposonmediated transgenesis. Journal of the American Mosquito Control Association. 2006; 22(3): 382-9.
14. Joshi V, Sharma RC. Impact or vertically transmitted dengue virus on viability of eggs of virus-inoculated *Aedes aegypti*. Dengue Bulletin. 2001; 25: 103-6.