



## Assessment of Probiotic Properties of Lactic Acid Bacteria from Traditional Sourdoughs for Bread-Making in Turkey against Some Gut Conditions

Murat Doğan<sup>1\*</sup>, İsmail Hakkı Tekiner<sup>2</sup>

<sup>1</sup>Gastronomy and Culinary Arts Department, Faculty of Fine Arts, İstanbul Gelişim University, İstanbul, Turkey

<sup>2</sup>Nutrition and Dietetics Department, Faculty of Health Sciences, İstanbul Sabahattin Zaim University, İstanbul, Turkey

\*Corresponding author (muratdogan72@gmail.com)

### Abstract

This study aims to assess the probiotic properties of Lactic Acid Bacteria isolated from the traditional sourdoughs used for bread making in Turkey against some gut conditions. A total number of 29 samples from twelve provinces of Turkey were collected, and screened for the presence of lactic acid bacteria using microbiological methods. The microbiological screening yielded 148 presumptive isolates. Of them, 62.8% were characterized as lactic acid strains by VITEK® MS. Following that, the characterized isolates were subjected to probiotic property testing, including gastric acid resistance, bile resistance and hydrophobic ability. The results showed that 44.1% exceeded gastric pH resistance, 33.3% survived under gastrointestinal system bile salt conditions, and 10.8% exhibited high hydrophobicity ability. In conclusion, our study revealed that only 4.3% (*Enterococcus faecium*, *Lactobacillus brevis*, *Lactobacillus pentosus*, and *Lactobacillus plantarum*) out of 93 lactic acid bacteria isolated from the traditional sourdoughs could meet all probiotic requirements against some gut conditions.

Keywords: lactic acid bacteria, probiotic, probiotic property, sourdough.

Article information:

Received: 25 August 2019

Accepted: 06 October 2019

Available online:

10 October 2019

Keywords:

lactic acid bacteria

probiotic

probiotic property

sourdough

© 2019

Indonesian Food Technologists

This is an open access article under the CC BY-NC-ND license

doi: 10.17728/jaft.5662

### Introduction

Probiotics are living microorganisms of the human intestinal microflora that show beneficial functions by keeping the human intestine in balance. Such bacteria form a colony in the human intestine, and compete with harmful microorganisms. The beneficial effects of probiotics have been known by Metchnikoff's studies since the early 1900s. Metchnikoff has attributed the longer lifespan and healthiness of Caucasian societies to fermented products and probiotics present in these products (Metchnikoff, 2004; Otlés, 2014; Doğan *et al.*, 2019).

Probiotics can ferment non-digestible oligosaccharide dietary fibers in the colon to short-chain fatty acids, and thus prevent colon cancer. They contribute to lactose digestion by fermenting lactose, increases immunity by adjusting IgA production, and inhibit the absorption of intestines by breaking down antigens, resulting in reduced allergy. The introduction of

probiotics as fermented food and food supplement are important for public health, and these studies have also increased public interest (Cui *et al.*, 2011; Kailasapathy, 2013; Butel, 2014; Polewski *et al.*, 2016).

Lactic Acid Bacteria (LAB) can reduce the redox potential in the intestine and generally break down carbohydrates and proteins. Anaerobic fermentation produces microbial metabolites such as lactic acid, succinate, acetate, propionate, butyrate, short chain fatty acids, microbial metabolites such as hydrogen, carbon dioxide, methane. *Lactobacillus casei*, *Lactobacillus acidophilus*, *Lactobacillus rhamnosus*, *Lactobacillus lactis*, *Lactobacillus brevis*, *Lactobacillus plantarum*, *Lactobacillus paraplantarum*, *Lactobacillus pentosus*, *Lactobacillus fermentum*, *Enterococcus faecium*, *Pediococcus pentosaceus*, and *Pediococcus acidilactici* were determined as the probiotic LAB strains (Dhanasekaran *et al.*, 2008; Lee and Salminen, 2009).

In order for a microorganism to exhibit probiotic

properties, it must reach the intestines alive. For this reason, they should primarily resist the gastric acid (pH 1.5-3.0), and maintain their viability. Probiotics that exceed the acidity of the stomach then encounter bile in the small intestine, and therefore must be resistant to the bile to maintain their viability. Probiotics, which pass into colon while maintaining their viability in the small intestine, must be able to attach to the epithel and mucosal surfaces in order to proliferate in the intestinal surfaces to form a colony. Probiotics have gastric acid resistance and bile resistance properties that enable them to form colonies by attaching to the colon, thus show their viability, and function in the gastrointestinal system (GIS) and these properties are among the most important probiotic selection criteria. Detection of microorganisms that provide probiotic properties is one of the objectives of food biotechnology (Dogan and Ozpinar, 2017).

Probiotics are also available in some foods and dietary supplements, in particular the fermented products, and similar to the probiotics that exist naturally in the Human gut. Probiotics has a shell that naturally protects it from acidic environments in the stomach and upper intestines until it reaches the desired location. For instance, sourdough is an ingredient in the cereal fermentation such as bread making. It is an initial preparation by fermenting flour and water mixture with yeast and LAB in order to improve sensory, nutritious, functional and technological properties of cereal based foods (Arendt *et al.*, 2011; Gobbetti *et al.*, 2014).

In terms of the media sources and environmental conditions of sourdough, LAB and yeasts compete with other microorganisms to form the dominant flora. LAB is a very important bacteria that plays significant role in the production of many fermented foods, including yoghurt, cheese, boza and kefir, that are mostly traditional Turkish fermented foods (Terpou *et al.*, 2017).

In this study, we aimed to assess the probiotic properties of Lactic Acid Bacteria isolated from the traditional sourdoughs used for bread making in the different provinces of Turkey against some gut conditions.

## Materials and Methods

### Sampling

Twenty-nine traditional sourdough samples were taken from the bakeries located in the twelve provinces of Turkey (2 İstanbul, 2 Kocaeli, 2 Adapazarı, 2 Tekirdağ, 2 Ankara, 3 Konya, 3 Karaman, 2 Aksaray, 2 Isparta, 3 Erzurum, 3 Malatya, and 3 Elazığ) (Table 1). No industrial yeast or starter culture was used. All the sourdough samples were selected among the traditionally used ones. The collected sourdough samples were initially mixed with wheat flour, water, and salt according to the Instructions by ISO 11133 (2014). Subsequently, the wet doughs were fermented for 6 hours at least. After that, the fermented dough was taken to the laboratory at 4°C, and cultured in the same day for further analysis.

### Chemicals and Reagents

The chemicals and reagents used in the study

were DeMan, Rogosa and Sharpe (MRS) agar (Merck 1.10660, Germany), MRS Broth (Merck 1.10661), M17 agar (1.15108 Merck) and M17 broth (1.15029 Merck) for cultural examination, pre-identification and storage of LAB strains; crystal violet, safranin, and lugol dyes for biochemical and morphological tests; and physiological saline (8.5 g NaCl dissolved in water, autoclaved 15 minutes at 121°C, and cooled to room temperature) for dilution and 20% glycerol (Merck 10494) for storage of culture, respectively. MRS and M17 broths were prepared with 1 N sterile Hydrochloric Acid (HCL) (Merck H9892), and 0.3% (w/v) Oxgall Bile bovine (B3883 Sigma Aldrich, USA) were used to determine gastric acid and bile resistance of the cultures. As much as 0.1 M Potassium Nitrate (KNO<sub>3</sub>) (P8394 Sigma Aldrich) and 0.3 ml Xylene (Sigma Aldrich) chemicals were used to calculate % hydrophobicity of the cultures. All the chemicals and reagents were selected and prepared according to the Instructions by ISO 11133 (2014).

Table 1. Distribution of Traditional Sourdough Samples by Province

Province	Number of Sample (n)
Istanbul	2
Kocaeli	2
Adapazarı	2
Tekirdağ	2
Ankara	2
Konya	3
Karaman	3
Aksaray	2
Isparta	2
Erzurum	3
Malatya	3
Elazığ	3

### Sample Preparation and Microbiological Analysis

Ten grams of sourdough sample were homogenized in 90 ml of sterile physiological saline solution. After that, each solution was diluted using physiological saline, and 1 mL was transferred onto *Lactobacillus* Agar acc. to MRS agar and M17 agar in parallel, followed by incubation for 24-48 hours at 37°C. At the end of the incubation, suspected LAB colonies were examined morphologically under microscope. To ensure the purity of the suspected colonies, MRS and/or M17 were inoculated into broth tubes, and activated at 37°C for 24 hours under aerobic/anaerobic conditions. Then, the matte-cream colored colonies were evaluated as LAB strains. Dyeing was performed for pure cultures; Gram (+), cocci and rods were determined under the light microscope, and followed by the catalase test. Those negative for catalase test were selected. Finally, these single colonies were re-enriched in MRS and/or M17 broths containing 20% glycerol, and stored at -80°C for the further analysis. The cultural examination of the suspected LAB strains were made according to the Instructions by ISO 6887-6 (2013) and ISO 11133 (2014).

Table 2. Source of Sourdough Samples by Provinces, LAB Microbial Load ( log CFU/g), pH and TTA Values (%), Culture Distribution and Co-Growing Cultures

Sample no	Sources	LAB	pH	TTA Lactic Acid	<i>E. faecium</i>	<i>L. brevis</i>	<i>L. casei</i>	<i>L. fermentum</i>	<i>L. lactis</i>	<i>L. pentosus</i>	<i>L. plantarum</i>	<i>L. rhamnosus</i>
1	Istanbul	8.57±0.03	3.9	0.51	+	+				+		
2	Istanbul	7.75±0.28	3.6	0.57	+			+			+	
3	Kocaeli	6.8±0.05	3.8	0.49					+		+	
4	Kocaeli	6.7±0.09	4.4	0.38			+			+		+
5	Adapazari	8.48±0.02	4.2	0.52		+				+		+
6	Adapazari	7.9±0.54	3.6	0.61		+		+				
7	Tekirdağ	6.83±0.4	4.9	0.33		+	+					+
8	Tekirdağ	6.95±0.18	4.2	0.37			+				+	+
9	Ankara	6.32±0.32	5.3	0.23		+		+				
10	Ankara	7.79±0.03	3.9	0.49			+		+	+		
11	Konya	6.04±0.18	4.8	0.29	+	+						
12	Konya	6.08±0.4	4.3	0.27		+						
13	Konya	6.59±0.14	5.6	0.18							+	
14	Karaman	6.74±0.22	5	0.21		+						+
15	Karaman	6.86±0.01	4.7	0.31				+	+			
16	Karaman	6.69±0.45	4.9	0.33			+					
17	Aksaray	7.74±0.05	3.7	0.58			+				+	+
18	Aksaray	6.92±0.03	5.4	0.22			+					
19	Isparta	8.75±0.02	3.4	0.69		+		+	+		+	
20	Isparta	6±0.5	4.7	0.38		+						+
21	Erzurum	6.92±0.19	3.9	0.5					+			+
22	Erzurum	6.96±0.06	3.5	0.65		+			+			
23	Erzurum	6.96±0.09	5.8	0.19			+					
24	Malatya	6.83±0.4	5.1	0.13			+				+	
25	Malatya	6.98±0.5	4.6	0.31				+				+
26	Malatya	6.23±0.02	4.1	0.37							+	
27	Elazığ	7.8±0.48	4.4	0.31	+	+			+			
28	Elazığ	6.87±0.2	3.7	0.52		+		+				
29	Elazığ	8.95±0.05	3.9	0.46		+	+	+				

Note: All experiments were repeated three times and the results were given as mean ± standard deviation

Table 3. High Performance Isolates According to Probiotic Properties

No	Type	Viability	Viability	Hydrophobicity (%)
		(CFU/g, pH 2.5)	(CFU/g, 0.3% w/v Oxgall Bile)	
1	<i>E. faecium</i>	5.92±0.05	5.36±0.06	63.82±7.4
2	<i>L. brevis</i>	4.84±0.07	4.98±0.12	98.46±2.5
3	<i>L. pentosus</i>	5.91±0.04	5.38±0.08	62.59±13.6
4	<i>L. plantarum</i>	5.04±0.06	4.63±0.10	74.48±5.0

All experiments were repeated three times and the results were given as mean ± standard deviation

#### Characterization by Mass Spectrometry

The suspected LAB strains stored in MRS and/or M17 broths at  $-80^{\circ}\text{C}$  were initially allowed to stand at room temperature for 2 hours. Subsequently, they were activated in aerobic and anaerobic conditions  $37^{\circ}\text{C}$  for 24-48 hours according to the Instructions by ISO 6887-6 (2013). The characterization of the strains was done by using VITEK® MS (bioMerieux, Marcy l'Etoile, France). A reference strain of *Escherichia coli* was used for the positive test control (Dubois *et al.*, 2012; Rifaat *et al.*, 2014).

#### Evaluation of Probiotic Properties of LAB Strains

The probiotic properties of the characterized isolates, including gastric acid resistance, bile resistance, and hydrophobicity for adhesion to the intestines abilities were assessed according to the studies by Yadav *et al.* (2016) and de Melo Pereira *et al.*

(2018). All probiotic property tests were repeated three times, and the results were given as mean±standard deviation.

#### Gastric Acid Resistance

As much as 1 N sterile HCl, basically the acid present inside the stomach-comprising HCl around 5,000 to 10,000 ppm, was added to MRS and/or M17 broths to adjust pH to 2.5 – the normal pH in the human stomach ranges from 1 to 3. The activated culture was centrifuged at 10,000 rpm for 10 minutes, and the supernatant was removed away. The resulting pellet were initially suspended in 7 ml of physiological saline, and subsequently incubated for 3 hours at  $37^{\circ}\text{C}$  with 1% inoculation into 10 ml broths with pH 2.5. Then, serial dilutions were made again, and allowed for incubation at  $37^{\circ}\text{C}$  for 72 hours. Finally, active bacteria counting were performed (Lee *et al.*, 2016).

### Bile Resistance Analysis

Seven milliliters of MRS/M17 broths were prepared with 0.3% (w/v) Oxgall Bile bovine. The activated cultures were centrifuged at 10,000 rpm for 10 minutes. The resulting pellets were suspended in 7 ml of physiological saline, and then incubated for 3 hours at 37°C with 1% inoculation into 10 ml broths containing Ox-Bile. Finally, serial dilutions were made, and incubated again at 37°C for 72 hours, followed by active bacteria counting (Liong and Shah, 2005).

### Hydrophobicity Analysis

The active cultures were centrifuged at 10,000 rpm for 15 minutes. The precipitate was washed twice with phosphate buffer solution, dissolved in 0.1 M KNO<sub>3</sub> (pH 6.2) buffer, and plated. The spectrophotometer was adjusted to an optical density (OD) of 600 (A0). One milliliter of the suspension was placed on 0.3 ml of xylene hydrocarbon, and incubated at room temperature for 4 hours. The OD of the aqueous phase was measured again on a spectrophotometer at 600 nm (A1). The adhesion percentage of the cultures to hydrocarbons was calculated using the formula  $[(A0-A1)/A0] \times 100$  (Mishra and Prasad, 2005).

## Results and Discussion

In this study, the probiotic properties of LAB strains isolated from the traditionally used sourdoughs for bread making in the twelve provinces of Turkey were assessed. Our analysis showed that microbiological screening yielded 148 presumptive isolates of LAB, and 62.8% of them were characterized as LAB by VITEK® MS. Among the characterized ones, 44.1% exceeded gastric pH resistance, 33.3% survived under gastrointestinal system bile salt conditions, and 10.8% exhibited high hydrophobicity ability. Overall, only 4.3% (*Enterococcus faecium*, *L. brevis*, *L. pentosus*, and *L. plantarum*) could meet all required probiotic properties.

The studies previously conducted in Turkey detected two identical strains from the sourdoughs, in particular *E. faecium* and *L. lactis*, except for another LAB species different from that of our study, which was *L. rhamnosus* (Diğrak and Özçelik, 1991; Menteş *et al.*, 2004; Ertekin and Çon, 2014; Ekinci *et al.*, 2016; Bakırcı and Köse, 2017). On the other hand, our work detected 148 presumptive isolates by microbiological methods, and VITEK® MS could identify 93 (62.8%) as LAB strains, which were *P. pentosaceus* (n=18), *L. brevis* (n=14), *L. casei* (n=10), *P. acidilactici* (n=10), *L. rhamnosus* (n=9), *L. pentosus* (n=8), *L. plantarum* (n=8), *L. fermentum* (n=6), *L. lactis* (n=6), and *E. faecium* (n=4) in number, respectively (Table 2).

To survive in the gut, the organisms must be tolerant to low pH and bile toxicity prevalent in the upper digestive tract. For colonization, they should exhibit good surface hydrophobicity and aggregation properties (Singh *et al.*, 2014). Gastric acid resistance, bile resistance and hydrophobicity were the probiotic properties of the identified LAB isolates from the sourdoughs. One of the most important selection criteria of probiotics is the resistance to the acidity of stomach. Active bacteria were counted in the medium that had a pH similar to the real gastric acidity (Ramirez-Chavarin

*et al.*, 2013; Dianawati *et al.*, 2016). Our analysis provided that 41 isolate (44%) (*E. faecium*, *L. brevis*, *L. casei*, *L. fermentum*, *L. lactis*, *L. pentosus*, *L. plantarum*, *L. rhamnosus*, *P. acidilactici*, and *P. pentosaceus*) were found to be resistant to gastric acidity (Figure 1). In order to protect LAB against gastric acid, microencapsulation method is recommended for the isolates that do not exhibit resistance to gastric acidity so that they can pass through the stomach with minimal harm to be used as probiotics. Firstly, the active isolates are powdered by lyophilization and then encapsulated with suitable material. In this way, active isolates are least damaged by the acidity of the stomach (Burgain *et al.*, 2011; Singh *et al.*, 2017).

LABs that exceed the acidity of the stomach encounter bile in the small intestine. In the process of selection of probiotics, resistance of the probiotics to bile resistance is the most important criteria after gastric acid resistance. Therefore, isolates that are considered as probiotic should be resistant to bile, and remain alive in the small intestine (Maldonado and Nader, 2015). A medium was formed with Oxgall Bile bovine to create a broth media similar to the small intestine medium and counts of the active cultures that survived in this medium were performed (Son *et al.*, 2018). In this study, 31 isolates (*E. faecium*, *L. brevis*, *L. casei*, *L. fermentum*, *L. lactis*, *L. pentosus*, *L. plantarum*, *L. rhamnosus*, *P. acidilactici*, and *P. pentosaceus*) was able to pass the bile resistance test among 41 isolates, that could pass the gastric acid resistance test (Figure 2). Within the bile resistant strains, *L. brevis*, *L. plantarum*, and *P. acidilactici* were obtained to have the higher bile resistance. Our results on the resistant ability of *L. brevis* and *L. plantarum* strains against bile were similar to the previous studies carried out by Ramos *et al.* (2013) and Landa-Salgado *et al.* (2019).

Another important criteria in the selection of probiotics is the adherence ability of the probiotic strains to the epithelial surfaces of intestine. A positive correlation was obtained between adherence of bacterial cells and cell surface hydrophobicity (Angmo *et al.*, 2016). For this reason, the hydrophobicity of the cultures that passed the gastric acid resistance and bile resistance tests were examined, and the percentages of adherence to hydrocarbons were calculated as previously described by Mishra and Prasad (2005). Our study showed that 10 isolates (*E. faecium*, *L. brevis*, *L. pentosus*, *L. plantarum*, *L. rhamnosus*, *P. acidilactici*, and *P. pentosaceus*) out of 31 isolates exhibited high hydrophobicity, ranging from 60% to 100% as previously given by Singh *et al.* (2017) (Figure 3). Some other works also reported the same findings on the significant hydrophobic ability of some LAB strains such as *L. brevis*, *L. pentosus*, *L. plantarum*, *L. rhamnosus*, *P. acidilactici*, and *P. pentosaceus* (Ramos *et al.*, 2013; Arasu *et al.*, 2015; Ayyash *et al.*, 2018; Aarti *et al.*, 2018; Palachum *et al.*, 2018; Maldonado *et al.*, 2018). Overall, only 4 (or 4.3%) strains (*E. faecium*, *L. brevis*, *L. pentosus*, and *L. plantarum*) out of the 93 characterized LAB isolates could pass the gastric acid resistance, bile resistance and hydrophobicity tests (Table 3, Figure 4, Figure 5).




Figure 1. Viability of Isolates Resistant to Gastric Acidity




Figure 2. Viability of Isolates Resistant to Bile




Figure 3. Percentage of Hydrophobicity of Isolates




Figure 4. Comparison of Isolates with pH 2.5 and Oxgall Bile




Figure 5. Hydrophobicity of Isolates

**Conclusion**

In conclusion, our study revealed that only 4.3% of the identified LAB strains (*E. faecium*, *L. brevis*, *L. pentosus*, and *L. plantarum*) from the traditional sourdoughs could meet all probiotic requirements against some gut conditions.

**References**

Aarti, C., Khusro, A., Varghese, R., Arasu, M.V.,

Agastian, P., Al-Dhabi, N.A., Choi, K.C. 2018. In vitro investigation on probiotic, anti-Candida, and antibiofilm properties of *Lactobacillus pentosus* strain LAP1. Archives of Oral Biology 89:99-106.  
 Angmo, K., Kumari, A., Bhalla, T.C. 2016. Probiotic characterization of lactic acid bacteria isolated from fermented foods and beverage of Ladakh. LWT-Food Science and Technology 66:428-435.  
 Arasu, M.V., Al-Dhabi, N.A., Rejiniemon, T. S., Lee, K.D., Huxley, V.A.J., Kim, D.H., Choi, K.C. 2015. Identification and characterization of *Lactobacillus brevis* P68 with antifungal, antioxidant and probiotic functional properties. Indian Journal of Microbiology 55(1):19-28.  
 Arendt, E.K., Moroni, A., Zannini, E. 2011. Medical nutrition therapy: use of sourdough lactic acid bacteria as a cell factory for delivering functional biomolecules and food ingredients in gluten free bread. In Microbial Cell Factories 10(1):S15.  
 Ayyash, M., Abushelaibi, A., Al-Mahadin, S., Enan, M.,

- El-Tarabily, K., Shah, N. 2018. In-vitro investigation into probiotic characterization of *Streptococcus* and *Enterococcus* isolated from camel milk. *LWT* 87:478-487.
- Bakırcı, F., Köse, E. 2017. Ekşi Hamurlardan Laktik Asit Bakterileri ve Mayaların İzolasyonu ve Tanımlanması.  *Akademik Gıda* 15(2):149-154.
- Burgain, J., Gaiani, C., Linder, M., Scher, J. 2011. Encapsulation of probiotic living cells: From laboratory scale to industrial applications. *Journal of Food Engineering* 104(4):467-483.
- Butel, M.J. 2014. Probiotics, gut microbiota and health. *Médecine et maladies infectieuses* 44(1):1-8.
- Cui, S.W., Nie, S., Roberts, K.T. 2011. Functional properties of dietary fibre. *Biotechnology* 4:517–525.
- de Melo Pereira, G.V., de Oliveira Coelho, B., Júnior, A.I.M., Thomaz-Soccol, V., Soccol, C.R. 2018. How to select a probiotic? A review and update of methods and criteria. *Biotechnology Advances* 36(8):2060-2076.
- Dhanasekaran, D., Saha, S., Thajuddin, N., Panneerselvam, A. 2008. Probiotics effect of lactobacillus isolates against bacterial pathogens in clarias orientails. *Facta Universitatis Medicine And Biology* 15:97-102.
- Diğrak, M., Özçelik, S. 1991. Elazığ ve Yöresinde Kullanılan Ekşi Mayanın Bileşimi, Morfolojik Fizyolojik ve Biyokimyasal Özellikleri. *Gıda* 16(5):325-331.
- Dianawati, D., Mishra, V., Shah, N.P., 2016. Viability, acid and bile tolerance of spray dried probiotic bacteria and some commercial probiotic supplement products kept at room temperature. *Journal of Food Science* 81(6):1472-1479.
- Dogan, M., Ozpinar, H. 2017. Investigation of probiotic features of bacteria isolated from some food products. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 23(4):554-562.
- Doğan, M., Tekiner, İ.H., Demirkesen Biçak, H. 2019. Probiotics From Food Products and Gastrointestinal Health. In: *Dietary Interventions in Gastrointestinal Diseases* (Ed.), (pp. 169-177). USA, Academic Press Elsevier.
- Dubois, D., Grare, M., Prere, M.F., Segonds, C., Marty, N., Oswald, E. 2012. Performances of the Vitek MS matrix-assisted laser desorption ionization–time of flight mass spectrometry system for rapid identification of bacteria in routine clinical microbiology. *Journal of Clinical Microbiology* 50(8):2568-2576.
- Ekinci, R., Şimşek, Ö., Küçükçuban, A., Nas, S. 2016. Evaluation of mono or mixed cultures of lactic acid bacteria in type II sourdough system. *Preparative Biochemistry and Biotechnology* 46(3):247-253.
- Ertekin, Ö., Çon, A. H. 2014. Farklı Gıdalardan İzole Edilen Laktik Asit Bakterilerinin Endüstriyel ve Probiyotik Özellikleri. *Academic Food Journal*, 12(4):6-16.
- Gobbetti, M., Rizzello, C.G., Di Cagno, R., De Angelis, M., 2014. How the sourdough may affect the functional features of leavened baked goods. *Food Microbiology* 37:30-40.
- ISO 6887-6. 2013. Microbiology of food and animal feed-Preparation of test samples, initial suspension and decimal dilutions for microbiological examination-Part 6: Specific rules for the preparation of samples taken at the primary production stage, Geneva, Switzerland.
- ISO 11133. 2014. Microbiology of food, animal feed and water-Preparation, production, storage and performance testing of culture media, Geneva, Switzerland.
- Kailasapathy K. 2013. Commercial sources of probiotic strains and their validated and potential health benefits-a review. *International Journal of Fermented Foods* 2(1):1-17.
- Landa-Salgado, P., Caballero-Cervantes, Y., Ramirez-Bribiesca, E., Maria Hernandez-Anguiano, A., Mariana Ramirez-Hernandez, L., Espinosa-Victoria, D., Hernandez-Sanchez, D. 2019. Isolation and identification of potentially probiotic lactic acid bacteria for Holstein calves in the Mexican Plateau. *Revista Mexicana De Ciencias Pecuarias* 10(1):68-83.
- Lee, K.W., Shim, J.M., Park, S.K., Heo, H.J., Kim, H.J., Ham, K.S., Kim, J.H. 2016. Isolation of lactic acid bacteria with probiotic potentials from kimchi, traditional Korean fermented vegetable. *LWT-Food Science and Technology* 71:130-137.
- Liong, M.T., Shah, N.P. 2005. Acid and bile tolerance and cholesterol removal ability of lactobacilli strains. *Journal of Dairy Science* 88(1):55-66.
- Lee, Y.K., Salminen, S. 2009. *Handbook of probiotics and prebiotics*. USA, John Wiley and Sons. NJ.
- Maldonado, N.C., Nader, M.E.F. 2015. Functional properties (acid and bile tolerance) and antibiotic susceptibility of lactic acid bacteria isolated from newborn calves for the design of a probiotic product. *International Journal of Veterinary Science and Research* 1(1):011-022.
- Maldonado, N.C., Ficooseco, C.A., Mansilla, F.I., Melián, C., Hébert, E.M., Vignolo, G.M., Nader-Macías, M.E. F., 2018. Identification, characterization and selection of autochthonous lactic acid bacteria as probiotic for feedlot cattle. *Livestock Science* 212:99-110.
- Menteş, Ö, Akçelik, M, Ercan, R. 2004. Türkiyede Üretilen Ekşi Hamurlardan *Lactobacillus* Suşlarının İzolasyonu, İdentifikasyonu ve Bu Suşların Temel Endüstriyel Özellikleri. *Gıda*, 29 (4):307-315.
- Metchnikoff, I.I., 2004. *The prolongation of life: optimistic studies*. USA, Springer Publishing Company, NY.
- Mishra, V., Prasad, D.N., 2005. Application of in vitro methods for selection of *Lactobacillus casei* strains as potential probiotics. *International Journal of Food Microbiology* 103(1):109-115.
- Otles, S., 2014. *Probiotics and Prebiotics*. In: *Food, Nutrition and Health* (Ed.), 2th ed (pp. 30-33). USA, CRC Press, NY.
- Palachum, W., Chisti, Y., Choorit, W., 2018. In-vitro assessment of probiotic potential of *Lactobacillus plantarum* WU-P19 isolated from a traditional

- fermented herb. *Annals of Microbiology* 68(2):79-91.
- Polewski, M.A., Krueger, C.G., Reed, J.D., Leyer, G., 2016. Ability of cranberry proanthocyanidins in combination with a probiotic formulation to inhibit in vitro invasion of gut epithelial cells by extra-intestinal pathogenic *E. coli*. *Journal of Functional Foods* 25:123-134.
- Ramirez-Chavarin, M.L., Wachter, C., Eslava-Campos, C.A., Perez-Chabela, M. L., 2013. Probiotic potential of thermotolerant lactic acid bacteria strains isolated from cooked meat products. *International Food Research Journal* 20(2):991-1000.
- Ramos, C.L., Thorsen, L., Schwan, R.F., Jespersen, L. 2013. Strain-specific probiotics properties of *Lactobacillus fermentum*, *Lactobacillus plantarum* and *Lactobacillus brevis* isolates from Brazilian food products. *Food Microbiology* 36(1):22-29.
- Rifaat, E.A., Tekiner, İ.H., Özpınar, H. 2014. Halk Sağlığı Açısından İçme ve Kullanma Sularında Koliform ve Fekal Koliform Bakterilerin Varlıklarının Klasik ve MASS Spektrometresi Yöntemleriyle İncelenmesi. *Electronic Journal of Food Technologies* 9(2):20-32.
- Singh, T.P., Malik, R.K., Renuka, G.K., 2014. Safety assessment and evaluation of probiotic potential of *Lactobacillus reuteri* strains under in vitro conditions. *International Journal of Current Microbiology and Applied Science* 3(2):335-348.
- Singh, P., Medronho, B., Alves, L., da Silva, G.J., Miguel, M.G., Lindman, B. 2017. Development of carboxymethyl cellulose-chitosan hybrid micro- and macroparticles for encapsulation of probiotic bacteria. *Carbohydrate Polymers* 175:87-95.
- Son, S.H., Jeon, H.L., Yang, S.J., Sim, M.H., Kim, Y.J., Lee, N.K., Paik, H.D. 2018. Probiotic lactic acid bacteria isolated from traditional Korean fermented foods based on  $\beta$ -glucosidase activity. *Food Science and Biotechnology* 27(1):123-129.
- Terpou, A., Gialleli, A.I., Bekatorou, A., Dimitrellou, D., Ganatsios, V., Barouni, E., Kanellaki, M. 2017. Sour milk production by wheat bran supported probiotic biocatalyst as starter culture. *Food and Bioproducts Processing* 101:184-192.
- Yadav, R., Puniya, A.K., Shukla, P. 2016. Probiotic properties of *Lactobacillus plantarum* RYPR1 from an indigenous fermented beverage Raabadi. *Frontiers in Microbiology* 7:1683.