



Original Article

The association between dietary fat and mineral intake with semen parameters: A cross-sectional study in infertile men

Mehran Nouri^{1, 2} Ph.D. Candidate, Sanaz Mehrabani³ Ph.D. Candidate, Hossein Firoozbakht⁴ M.Sc., Elmira Vataniyan⁵ M.Sc., Homayoun Abbasi⁶ M.D., Mahsa Shirani^{3, 7} M.Sc.

Corresponding Author:

Mahsa Shirani; School of Nutrition and Food Science, Isfahan University of Medical Sciences, Hezar Jereeb St., Isfahan, Iran.

Postal Code: 8174673461

Tel: (+98) 9132066179

Email: mahsa.shirani1373@yahoo.com

ORCID:

<https://orcid.org/0000-0002-6630-7473>

Received: 28 March 2021

Revised: 4 August 2021

Accepted: 13 December 2021

Production and Hosting by Knowledge E

© Nouri et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editor-in-Chief:

Aflatoonian Abbas M.D.

¹Students' Research Committee, School of Nutrition and Food Science, Shiraz University of Medical Sciences, Shiraz, Iran.

²Department of Community Nutrition, School of Nutrition and Food Science, Shiraz University of Medical Sciences, Shiraz, Iran.

³Department of Community Nutrition, School of Nutrition and Food Science, Food Security Research Center, Isfahan University of Medical Sciences, Isfahan, Iran.

⁴Department of Nutrition, Science and Research Branch, Islamic Azad University, Tehran, Iran.

⁵Department of Cellular and Molecular Nutrition, School of Nutritional Science and Dietetics, Tehran University of Medical Science, Tehran, Iran.

⁶Isfahan Fertility and Infertility Center, Isfahan, Iran.

⁷Students' Research Committee, Isfahan University of Medical Sciences, Isfahan, Iran.

Abstract

Background: Infertility has been a significant problem for couples in recent decades, and the issue can often lie with the male rather than the female.

Objective: This study aimed to investigate the effects of fats and minerals intake on semen parameters in infertile men.

Materials and Methods: This cross-sectional study was performed on 260 infertile men referred to Isfahan Infertility Clinic, Isfahan, Iran in the summer of 2018. Semen parameters regarding sperm concentration, total motility, normal morphology, and sperm volume were considered. To assess dietary intakes, a validated food frequency questionnaire was used.

Results: In the adjusted model, the prevalence of abnormal concentration was 5.23 times higher in the top quartile of calcium intake, compared with the bottom quartile ($p = 0.03$). Also, the prevalence of abnormal morphology was 68% lower in the third quartile of calcium intake ($p = 0.03$). Additionally, the prevalence of abnormal concentration was 84% lower in the top quartile of folate intake in comparison to the bottom quartile ($p = 0.01$) and the prevalence of abnormal morphology was 70% lower in the top quartile of folate intake ($p = 0.03$). Additionally, the prevalence of abnormal concentration was 72% lower in the top quartile of selenium intake in comparison to the bottom quartile ($p = 0.04$). Furthermore, in the crude model, the prevalence of abnormal volume was 64% lower in the second quartile of linoleic acid intake rather than the first quartile ($p = 0.01$).

Conclusion: In conclusion, diets containing higher amounts of folate and selenium, and lower amounts of cholesterol, saturated fatty acid and calcium were associated with more favorable semen quality parameters.

Key words: Infertility, Semen quality, Minerals, Fats, Cross-sectional study.

OPEN ACCESS

1. Introduction

Over recent decades, infertility has been a significant issue in developing countries, affecting 15% of all couples falling within fertile age brackets (1). A meta-analysis estimated the infertility rate within the Iranian population to be about 10.9%. In approximately 40% of infertile couples, male infertility is a factor (2). Infertility is defined as the inability of a couple to conceive within 12 months or more of regular unprotected intercourse (3).

Infertility can be caused by various disorders including: anatomical or hormonal problems, genetic abnormalities, and infections. In addition, environmental factors such as air pollution, industrial chemicals, stress and depression, alcohol consumption, or smoking are all considered to be risk factors, which reduce sperm quality parameters in developed countries (4). Moreover, studies have shown a relationship between semen quality and lifestyle factors, including physical activity and diet (5).

Dietary fats are one of the dietary components that can contribute to infertility. Due to large amounts of unsaturated fatty acids in the membrane of sperm cells, the chance of lipid peroxidation in sperm cells and production of free radicals such as hydrogen peroxide, superoxide, and hydroxyl radicals is high (6, 7). Lipid peroxidation can drastically alter membrane integrity and permeability, alter cellular enzyme activation and cell apoptosis, and thus affect spermatozoa. As a result, sperm count and motility decrease and sperm shapes become abnormal (8).

Furthermore, over-consumption of saturated and trans-fatty acids can reduce sperm quality by increasing the rate of inflammation (9). Meanwhile, previous studies have shown that there is an important association between the intake of certain dietary minerals, zinc and iron, and semen quality (10-12). Although the association between dietary

fat and the quality of sperm has been evaluated before, the results are controversial.

Therefore, this study aimed to evaluate the association between dietary fats and minerals with factors related to semen quantity and quality.

2. Materials and Methods

2.1. Study design

This cross-sectional study was performed on 260 infertile men referred to Isfahan Infertility Clinic in Isfahan, Iran, in June-August 2018. Participants were aged between 18-55 yr with a history of primary or secondary infertility in the last 5 yr. All participants voluntarily entered the study. Exclusion criteria were: a history of urinary infection, use of supplements, androgens, anticoagulants, cytotoxic drugs or immunosuppressant, and metabolic diseases such as diabetes or renal disease (13). Additionally, participants with incomplete information or caloric intakes outside the range of 800-4200 kcal/day were excluded from the study. Finally, the data for 260 cases were used in the study.

2.2. Sperm parameters

After 3 days of abstinence, semen samples were taken from participants. The samples were collected in sterile containers and placed at room temperature for 30 min to ensure liquidity. Analysis of the semen samples was done according to the 5th edition of the World Health Organization laboratory manual for human semen (14). 4 dependent semen parameters were measured: sperm concentration (SC); total sperm movement (TSM); normal sperm morphology (NSM); and seminal volume (SV). Abnormal semen parameters were defined as oligospermia: SC < 20 M/ml, TSM < 60%, sperm volume < 3 ml and NSM < 65% (13).

As we were not able to analyze data with NSM < 65% in our study, the World Health Organization cut point of NSM < 4% was considered for normal morphology (13).

2.3. Assessment of dietary intakes

To assess dietary intakes of individuals, a validated detailed food frequency questionnaire was used in this study. It included 168 food items comprising the most common Iranian foods and has been previously shown to be valid and reproducible for use in Iranian adults (15).

For each food item, participants reported their average consumption during the last year in terms of frequency of use. This classification includes the following: ≥ 6 times a day, 3-5 times a day, 2-3 times a day, once a day, 5-6 times a week, 2-4 times a week, once a week, 1-3 times a month and < 1 a month. The selected group was then converted to the daily intake for each food item. Nutritionist IV for Windows software was used to determine micronutrient intakes for each participant.

2.4. Assessment of other variables

Demographic data, medical history, and intake of supplements were collected via a structured questionnaire for all participants. In this study, weight and height were measured with high accuracy and then body mass index was calculated according to the formula (kg/m^2).

2.5. Ethical considerations

The study was ethically approved by the Ethical Committee of Isfahan University of Medical Sciences, Isfahan, Iran (Code: IR.MUI.RESEARCH.REC.1397.232). All of the participants completed the informed consent form.

2.6. Statistical analysis

Statistical analyses were carried out using the Statistical Package for the Social Sciences (SPSS) for Windows software (version 20.0, SPSS Inc, Chicago, IL, USA). P-values < 0.05 were considered statistically significant. To describe quantitative variables, mean \pm standard deviation (SD) and for the categorical outcome, frequency and percentage were used. Analysis of covariance (ANCOVA) was used to determine the association between semen parameters as dependent variables, and fats and minerals covariates as independent variables.

Multiple logistic regression (odds ratios [ORs] with 95% confidence intervals [CI]) was used to evaluate the relationship between dietary fat and mineral intakes with sperm quality parameters in the crude and adjusted models. In the adjusted model, potential confounding variables including age, BMI, education, total energy intake, alcohol, smoking, and vitamin-mineral use were justified by ANCOVA, and the first quartile was considered as the reference level.

3. Results

3.1. Baseline characteristics of participants

Finally, 260 participants aged between 18 and 55 yr were included in this study. 36% of the subjects were smokers, and 20% of participants used alcohol. Furthermore, 30% of participants used multivitamins. All of the participants had semen analyses performed and the semen analysis parameters included: the mean SC; SV; TSM; and NSM. Also, the dietary intake of participants is shown in table I.

3.2. Correlation between sperm related parameters and dietary components

There was a significant association between dietary intakes of protein and sperm volume ($p < 0.001$), dietary intakes of carbohydrates and SC ($p = 0.01$), and also dietary intakes of carbohydrates and sperm motility ($p = 0.03$, Table II).

3.3. Association between dietary mineral intakes and sperm parameters

In the adjusted model, the prevalence of abnormal SC was 5.23 times higher in the top quartile of calcium intake, compared with the bottom quartile ($p = 0.03$). Also, the prevalence of abnormal sperm morphology was 68% lower in the third quartile of calcium intake, compared with the first quartile ($p = 0.03$). Additionally, the prevalence of abnormal SC was 84% lower in the top quartile

of folate intake in comparison to the bottom quartile ($p = 0.007$) and the prevalence of abnormal sperm morphology was 70% lower in the top quartile of folate intake, compared with the first quartile ($p = 0.03$). Also, the prevalence of abnormal SC was 72% lower in the top quartile of selenium intake in comparison to the bottom quartile ($p = 0.04$, Table III).

3.4. Association between dietary fat intakes and sperm parameters

As is shown in the adjusted model, the prevalence of abnormal SC was 5.6 and 4.8 times higher in the third quartile of cholesterol and saturated fatty acid (SFA) intake, respectively, compared with the first quartile ($p = 0.01$). Additionally, the prevalence of abnormal sperm volume was 64% lower in the second quartile of linoleic acid intake than in the first quartile ($p = 0.01$) in the crude model; however, this was not significant in the adjusted model (Table IV).

Table I. Baseline characteristics of participants

Characteristics	Mean \pm SD (n = 260)
Age (yr)	31.24 \pm 4.23
BMI (kg/m ²)	26.94 \pm 4.09
MET (MET-h/wk)	29.27 \pm 2.12
WC (cm)	94.51 \pm 10.35
Dietary intake	
Energy (kcal)	2516.51 \pm 686.94
Carbohydrate (gr/day)	361.38 \pm 104.45
Protein (gr/day)	94.07 \pm 27.04
Fat (gr/day)	86.77 \pm 36.48
Caffeine (mg/day)	79.41 \pm 61.05
Semen parameters	
SV (mL)	4.13 \pm 2.06
SC (mill/mL)	13.11 \pm 6.01
TSM (%)	29.76 \pm 18.08
NSM (%)	2.00 \pm 1.68

BMI: Body mass index, MET: Metabolic equivalents, WC: Waist circumference, SV: Semen volume, SC: Sperm concentration, TSM: Total sperm motility, NSM: Normal sperm morphology

Table II. Correlation between sperm-related parameters and dietary components (n = 260)

Mean range of value	Calorie intake (kcal)	Protein (gr/day)	Carbohydrate (gr/day)	Fat (gr/day)	Caffeine
SV (mL)	0.10 (0.08)	0.18 (0.00)	0.08 (0.19)	0.10 (0.10)	0.06 (0.29)
SC (10 ⁶ /ml)	0.07 (0.24)	0.02 (0.64)	0.14 (0.01)	-0.05 (0.37)	0.03 (0.56)
TSM (%)	0.05 (0.34)	-0.02 (0.72)	0.13 (0.03)	-0.05 (0.34)	0.05 (0.37)
NSM (%)	-0.03 (0.61)	-0.09 (0.12)	0.00 (0.98)	-0.05 (0.36)	0.05 (0.37)

Values are reported by correlation coefficient (p-value) as assessed by regression. SV: Semen volume, SC: Sperm concentration, TSM: Total sperm motility, NSM: Normal sperm morphology

Table III. Abnormal semen quality across quartiles of mineral intake (n = 260)

Variables	SC < 20 M/ml	P-value	TSM < 60%	P-value	NSM < 4%	P-value	SV < 3 ml	P-value
Calcium								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	1.00 (0.38-2.59)	1.00	0.65 (0.10-4.06)	0.65	0.66 (0.27-1.62)	0.36	1.07 (0.50-2.27)	0.84
Q3	1.13 (0.42-2.99)	0.80	0.31 (0.06-1.60)	0.16	0.40 (0.17-0.96)	0.04	1.23 (0.59-2.60)	0.57
Q4	2.18 (0.70-6.78)	0.17	0.48 (0.08-2.74)	0.41	1.13 (0.42-2.99)	0.80	0.66 (0.29-1.47)	0.31
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	1.41 (0.50-3.99)	0.51	1.17 (0.17-8.04)	0.87	0.58 (0.22-1.50)	0.26	1.19 (0.53-2.66)	0.66
Q3	1.95 (0.61-6.27)	0.25	0.74 (0.11-4.76)	0.75	0.32 (0.11-0.89)	0.03	1.45 (0.60-3.49)	0.40
Q4	5.23 (1.13-24.25)	0.03	2.18 (0.22-21.32)	0.50	0.83 (0.24-2.90)	0.78	0.84 (0.30-2.40)	0.75
Zinc								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	0.40 (0.14-1.14)	0.08	0.18 (0.02-1.65)	0.13	0.60 (0.26-1.36)	0.22	1.32 (0.63-2.77)	0.45
Q3	0.63 (0.21-1.89)	0.41	0.18 (0.02-1.65)	0.13	1.00 (0.42-2.36)	1.00	0.92 (0.43-1.98)	0.84
Q4	1.00 (0.30-3.28)	1.00	0.23 (0.02-2.19)	0.20	1.78 (0.68-4.64)	0.23	0.72 (0.33-1.59)	0.42
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	0.49 (0.16-1.49)	0.21	0.35 (0.03-3.34)	0.36	0.72 (0.30-1.73)	0.46	1.49 (0.67-3.32)	0.32
Q3	0.88 (0.24-3.13)	0.84	0.55 (0.05-5.72)	0.61	1.35 (0.48-3.76)	0.56	1.12 (0.45-2.76)	0.79
Q4	1.78 (0.34-9.23)	0.48	1.85 (0.10-33.09)	0.67	3.04 (0.78-11.87)	0.10	1.01 (0.33-3.04)	0.97
Folate								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	0.84 (0.26-2.65)	0.77	0.00 (0.00)	0.99	1.00 (0.40-2.50)	1.00	1.00 (0.47-2.13)	1.00
Q3	1.55 (0.41-5.77)	0.51	0.00 (0.00)	0.99	0.81 (0.33-1.98)	0.65	1.00 (0.47-2.13)	1.00
Q4	0.28 (0.10-0.78)	0.01	0.00 (0.00)	0.99	0.53 (0.22-1.23)	0.14	0.92 (0.43-1.98)	0.84
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	0.67 (0.20-2.24)	0.52	0.00 (0.00)	0.99	0.78 (0.30-2.06)	0.62	1.15 (0.52-2.56)	0.72
Q3	1.06 (0.25-4.44)	0.93	0.00 (0.00)	0.99	0.54 (0.19-1.49)	0.23	1.31 (0.55-3.12)	0.53
Q4	0.16 (0.04-0.60)	0.007	0.00 (0.00)	0.99	0.30 (0.09-0.91)	0.03	1.48 (0.55-3.12)	0.43
Selenium								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	2.70 (1.03-7.11)	0.04	1.00 (0.19-5.14)	1.00	0.90 (0.36-2.21)	0.81	0.85 (0.39-1.860)	0.69
Q3	2.32 (0.91-5.90)	0.07	3.09 (0.31-30.57)	0.33	0.90 (0.36-2.21)	0.81	1.16 (0.54-2.47)	0.70
Q4	6.74 (1.86-24.48)	0.056	0.34 (0.08-1.36)	0.12	0.53 (0.22-1.23)	0.14	1.24 (0.58-2.64)	0.56
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	0.79 (0.25-2.42)	0.68	1.42 (0.26-7.64)	0.68	0.76 (0.29-1.94)	0.56	1.02 (0.45-2.32)	0.95
Q3	2.17 (0.50-9.46)	0.29	6.28 (0.54-72.80)	0.14	0.69 (0.25-1.85)	0.46	1.58 (0.68-3.68)	0.28
Q4	0.28 (0.08-0.94)	0.04	0.88 (0.17-4.64)	0.88	0.36 (0.13-1.04)	0.06	2.33 (0.91-5.99)	0.07

Data presented as OR (95%CI). Adjusted model: Adjusted for age, BMI, Education, Energy intake, Alcohol, Smoking, Vitamin-mineral use. Q: Quartile, Multivariable-adjusted odds ratio was used. SV: Semen volume, SC: Sperm concentration, TSM: Total sperm motility, NSM: Normal sperm morphology

Table IV. Abnormal semen quality across quartiles of dietary fat intake

Variables	SC < 20 M/ml	P-value	TSM < 60%	P-value	NSM < 4%	P-value	SV < 3 ml	P-value
Cholesterol								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	0.81 (0.33-1.98)	0.65	1.00 (0.23-4.18)	1.00	0.63 (0.27-1.47)	0.29	1.54 (0.72-3.27)	0.25
Q3	4.21 (1.11-15.88)	0.03	2.06 (0.36-11.69)	0.41	1.00 (0.41-2.42)	1.00	0.92 (0.41-2.03)	0.84
Q4	1.68 (0.61-4.66)	0.31	0.78 (0.20-3.07)	0.73	1.00 (0.41-2.42)	1.00	1.16 (0.54-2.51)	0.69
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	0.99 (0.39-2.53)	0.98	1.61 (0.36-7.18)	0.53	0.60 (0.25-1.45)	0.26	1.80 (0.81-3.98)	0.14
Q3	5.63 (1.39-22.82)	0.01	4.13 (0.65-26.28)	0.13	0.93 (0.36-2.41)	0.88	1.14 (0.48-2.69)	0.75
Q4	2.57 (0.78-8.42)	0.11	2.22 (0.44-11.18)	0.33	0.92 (0.33-2.53)	0.87	1.62 (0.66-3.97)	0.28
SFA								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	1.00 (0.4-2.50)	1.00	2.06 (0.36-11.69)	0.41	0.63 (0.27-1.47)	0.29	0.69 (0.33-1.47)	0.34
Q3	3.10 (0.93-10.33)	0.06	1.00 (0.23-4.18)	1.00	1.00 (0.41-2.42)	1.00	0.64 (0.30-1.37)	0.25
Q4	1.45 (0.54-3.88)	0.45	0.78 (0.20-3.07)	0.73	1.00 (0.41-2.42)	1.00	0.64 (0.30-1.37)	0.25
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	1.27 (0.48-3.34)	0.31	3.54 (0.58-21.29)	0.16	0.63 (0.26-1.52)	0.31	0.76 (0.34-1.65)	0.49
Q3	4.82 (1.30-17.87)	0.01	2.34 (0.48-11.42)	0.29	0.96 (0.35-2.59)	0.94	0.71 (0.30-1.68)	0.43
Q4	3.50 (0.84-14.60)	0.08	5.44 (0.74-39.93)	0.09	0.95 (0.27-3.27)	0.94	0.80 (0.27-2.34)	0.69
Mufa								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	1.00 (0.38-2.59)	1.00	0.73 (0.15-3.43)	0.69	1.09 (0.47-2.56)	0.82	0.81 (0.39-1.68)	0.57
Q3	1.50 (0.53-4.23)	0.43	1.52 (0.24-9.43)	0.65	1.09 (0.47-2.56)	0.82	0.64 (0.30-1.37)	0.25
Q4	1.50 (0.53-4.23)	0.43	0.47 (0.11-1.99)	0.30	1.09 (0.47-2.56)	0.82	0.54 (0.25-1.18)	0.12
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	1.33 (0.48-3.65)	0.57	1.21 (0.24-5.91)	0.81	1.07 (0.43-2.63)	0.87	0.82 (0.37-1.80)	0.63
Q3	2.48 (0.75-8.17)	0.13	3.79 (0.52-27.47)	0.18	1.02 (0.38-2.73)	0.96	0.63 (0.26-1.52)	0.30
Q4	3.93 (0.83-18.53)	0.08	2.93 (0.34-24.90)	0.32	0.98 (0.28-3.35)	0.97	0.54 (0.17-1.66)	0.28
Pufa								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	1.50 (0.53-4.23)	0.43	0.73 (0.15-3.43)	0.69	1.29 (0.57-2.90)	0.53	0.46 (0.21-1.02)	0.05
Q3	1.00 (0.38-2.59)	1.00	3.09 (0.31-30.57)	0.33	1.73 (0.74-4.07)	0.20	0.55 (0.26-1.18)	0.13
Q4	1.50 (0.53-4.23)	0.43	0.40 (0.09-1.62)	0.20	1.73 (0.74-4.07)	0.20	0.81 (0.39-1.68)	0.58
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	1.78 (0.60-5.24)	0.29	1.04 (0.21-5.15)	0.95	1.41 (0.60-3.30)	0.42	0.52 (0.23-1.18)	0.12
Q3	1.34 (0.45-3.95)	0.58	5.98 (0.54-65.28)	0.14	2.07 (0.79-5.40)	0.13	0.70 (0.30-1.66)	0.42
Q4	2.63 (0.63-10.83)	0.18	1.39 (0.20-9.49)	0.73	2.41 (0.75-7.73)	0.13	1.25 (0.46-3.40)	0.65
Linoleic acid								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	1.82 (0.62-5.34)	0.27	0.75 (0.16-3.49)	0.71	1.59 (0.69-3.68)	0.27	0.36 (0.15-0.82)	0.01
Q3	0.92 (0.36-2.35)	0.87	3.19 (0.3-31.56)	0.32	1.47 (0.64-3.35)	0.35	0.72 (0.34-1.50)	0.38
Q4	1.53 (0.54-4.31)	0.41	0.40 (0.10-1.65)	0.20	1.77 (0.75-4.16)	0.18	0.91 (0.44-1.88)	0.80

Table IV. (Continued)

Variables	SC < 20 M/ml	P-value	TSM < 60%	P-value	NSM < 4%	P-value	SV < 3 ml	P-value
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	2.04 (0.67-6.18)	0.20	1.01 (0.21-4.87)	0.98	1.70 (0.71-4.02)	0.22	0.42 (0.18-1.00)	0.05
Q3	1.88 (0.41-3.40)	0.74	6.35 (0.58-69.37)	0.12	1.69 (0.67-4.25)	0.26	1.04 (0.45-2.39)	0.92
Q4	2.39 (0.60-9.46)	0.21	1.40 (0.21-9.28)	0.72	2.24 (0.72-6.95)	0.16	1.70 (0.64-4.52)	0.28
Linolenic acid								
Crude								
Q1	Reference		Reference		Reference		Reference	
Q2	1.03 (0.38-2.81)	0.94	0.76 (0.16-3.55)	0.73	1.50 (0.64-3.50)	0.34	0.66 (0.30-1.40)	0.28
Q3	0.93 (0.35-2.47)	0.89	0.77 (0.16-3.60)	0.74	1.53 (0.65-3.56)	0.32	0.69 (0.33-1.47)	0.34
Q4	1.66 (0.55-4.98)	0.36	0.77 (0.16-3.60)	0.74	1.38 (0.60-3.18)	0.43	0.69 (0.33-1.47)	0.34
Adjusted								
Q1	Reference		Reference		Reference		Reference	
Q2	1.10 (0.40-3.03)	0.84	0.90 (0.18-4.29)	0.89	1.49 (0.63-3.50)	0.35	0.68 (0.31-1.47)	0.33
Q3	1.14 (0.40-3.24)	0.79	1.60 (0.30-8.42)	0.57	1.55 (0.63-3.78)	0.33	0.80 (0.36-1.77)	0.58
Q4	2.45 (0.67-8.95)	0.17	3.07 (0.45-20.81)	0.25	1.40 (0.52-3.75)	0.50	0.88 (0.36-2.15)	0.78

Data presented as OR (95% CI). Adjusted model: Adjusted for age, BMI, Education, Energy intake, Alcohol and smoking, Vitamin-mineral use. Q: Quartile, SFA: Saturated fatty acid, Mufa: Mono-unsaturated fatty acid, Pufa: Poly-unsaturated fatty acid. Multivariable-adjusted odds ratio was used for analysis. SV: Semen volume, SC: Sperm concentration, TSM: Total sperm motility, NSM: Normal sperm morphology

4. Discussion

In this cross-sectional study on 260 infertile men, we found that calcium intake was reversely correlated with SC and positively related to NSM. Furthermore, folate intake showed a significant relationship with SC and morphology and there was a significant relationship between SC and selenium intake. In addition, there was a reverse association between cholesterol and SFA intake with SC.

The results of recent studies have both been in favor of, and contradict, the results of the present study. The possible effect of dairy foods and calcium on male fertility is highly controversial. Unlike the present study, the results of other studies have found no association between blood calcium concentration, the time of first serving, and the time of withdrawal from pregnancy, respectively (16, 17). In addition, one study found that serum calcium concentrations were in contrast to Ca^{2+} concentrations, which could be affected by blood pH and protein concentrations

(16). Also, in plasma, calcium is available in 3 forms: ionized, complex, and bound protein. Among these, calcium ionization (Ca^{2+}) is the physiologically active form, so measuring plasma Ca^{2+} concentration, as opposed to total plasma concentration, is an indicator of physiologically present calcium (18). In another study, no significant relationship was observed between calcium and sperm parameters (19). However, high-fat dairy products and cheese have been inversely related to sperm quality parameters (20), while low fat milk and skimmed milk were associated with better classical semen indicators. Consumption of low-fat and skimmed milk can be associated with higher levels of insulin and insulin-like growth factor-1 and thus spermatogenesis, while the reason for the adverse effect of high-fat dairy on infertility can be ascribed to the high content of SFA (21).

As seen in our study, selenium intake indicated a preservative effect against SC. For example, some minerals such as selenium and zinc may develop semen quality via their anti-inflammatory and

protective roles against free radicals (22). There is a direct relation between the antioxidant status and the production of reactive oxygen species (ROS) in sperm (23). In addition, high concentrations of ROS negatively affect sperm DNA and, in turn, impact sperm motility, vitality and concentration, as well as potentially contributing to miscarriages and developmental abnormalities in offspring (24). Antioxidants are considered as 'scavengers' of ROS and their use has been studied as a treatment to neutralize the negative effect of high concentrations of ROS on semen parameters. A recent study also indicated the importance of selenium in completing sperm maturation and quality (25). Selenium, as a selenoprotein, is more involved in male reproductive function. Adequate selenium intake is important for normal spermatogenesis and sperm maturation (26).

Additionally, folate intake displayed significant correlations with SC and morphology in our study. In men with high folate intake, lower levels of aneuploid sperm have been reported, suggesting that this vitamin (found mainly in green leafy vegetables) may play a role in spermatogenesis due to its role in DNA retention, RNA transport, and protein synthesis (27). In addition, the beneficial effects of dietary antioxidants such as vitamin C, vitamin E, and folate on sperm motility have already been shown. The results of a randomized controlled trial showed that after treatment with sulfate and folic acid, the number of normal sperm increased in infertile and fertile men (28). In a clinical trial, analysis of DNA methylation in sperm with exposure to folic acid suggested the susceptibility of mice and humans in sequences of potential importance to germ cells and embryonic growth. This study also suggested hypomethylation of sperm DNA and involvement in the regulation of Methyl tetrahydrofolate reductase as a possible

mechanism for folic acid supplementation for different species (29).

Furthermore, supporting the findings of our study, a previous study found an inverse relationship between dietary fat, especially cholesterol and SFA intake and SC. Trans-fatty acids, SFAs, and preservatives or hormonal residues such as xenobiotic or anabolic steroids may alter sperm quality (30). High SFA concentrations and low omega-3, poly-unsaturated fatty acids (PUFA) levels are associated with decreased fertility parameters (19). In an animal study, some dietary SFAs did not affect sperm quality parameters. However, one human study showed that higher levels of palmitic acid or stearic acid were present in the sperm of infertile men (31). Also, in our study, linoleic acid intake showed some preservative effects on sperm volume in the second quartile of intake. Several mechanisms have been proposed for the beneficial effects of PUFA and MUFA on sperm quality. Adipose tissue plays an important role in causing oxidative stress after an inflammatory disorder that may alter normal reproductive pathways and sperm activity (32, 33). In fact, under stress conditions, mitochondria are likely to cause a cascade of oxidative damage in the testicular environment. MUFA and PUFA fatty acids have protective effects against increased ROS levels due to their anti-inflammatory and antioxidant properties (34, 35).

4.1. Limitations

The present study, like other cross-sectional studies, is not able to definitively determine the cause-and-effect relationship between the variables and it is suggested that clinical trial or case-control studies be performed to confirm the results. Another limitation is that plasma or semen levels of minerals and fatty acids were not

considered in this study. Finally, the 168-item food frequency questionnaire has some limitations in recording food intakes and relies on the individual's memory, and is not always as accurate when employed by the elderly or illiterate people; therefore, the use of 3-day food recall could be useful in future studies.

5. Conclusion

The results of the present study showed that high calcium intake was inversely related to SC and directly related to normal morphology. Also, folate and selenium intake showed protective effects on SC. On the other hand, the high levels of cholesterol and SFA intake showed an inverse relationship with SC.

Acknowledgments

We appreciate Isfahan infertility clinic, Isfahan, Iran. This study received no financial support.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- [1] Benedetti S, Tagliamonte MC, Catalani S, Primiterra M, Canestrari F, De Stefani S, et al. Differences in blood and semen oxidative status in fertile and infertile men, and their relationship with sperm quality. *Reprod Biomed Online* 2012; 25: 300–306.
- [2] Aflakseir A, Mahdiyar M. The role of religious coping strategies in predicting depression among a sample of women with fertility problems in Shiraz. *J Reprod Infertil* 2016; 17: 117–122.
- [3] Zegers-Hochschild F, Adamson GD, de Mouzon J, Ishihara O, Mansour R, Nygren K, et al. The international committee for monitoring assisted reproductive technology (ICMART) and the world health organization (WHO) revised glossary on ART terminology, 2009. *Hum Reprod* 2009; 24: 2683–2687.
- [4] Karayiannis D, Kontogianni MD, Mendorou C, Douka L, Mastrominas M, Yiannakouris N. Association between adherence to the Mediterranean diet and semen quality parameters in male partners of couples attempting fertility. *Hum Reprod* 2017; 32: 215–222.
- [5] Gabrielsen JS, Tanrikut C. Chronic exposures and male fertility: The impacts of environment, diet, and drug use on spermatogenesis. *Andrology* 2016; 4: 648–661.
- [6] Lancellotti TES, Boarelli PV, Monclus MA, Cabrillana ME, Clementi MA, Espínola LS, et al. Hypercholesterolemia impaired sperm functionality in rabbits. *PLoS One* 2010; 5: e13457.
- [7] Dadkhah H, Kazemi A, Nasr-Isfahani MH, Ehsanpour S. The relationship between the amount of saturated fat intake and semen quality in men. *Iran J Nurs Midwifery Res* 2017; 22: 46–50.
- [8] Eslamian Gh, Amirjannati N, Rashidkhani B, Sadeghi MR, Baghestani AR, Hekmatdoost A. Dietary fatty acid intakes and asthenozoospermia: A case-control study. *Fertil Steril* 2015; 103: 190–198.
- [9] Attaman JA, Toth TL, Furtado J, Campos H, Hauser R, Chavarro JE. Dietary fat and semen quality among men attending a fertility clinic. *Hum Reprod* 2012; 27: 1466–1474.
- [10] Moslemi MK, Tavanbakhsh S. Selenium-vitamin E supplementation in infertile men: Effects on semen parameters and pregnancy rate. *Int J Gen Med* 2011; 4: 99–104.
- [11] Ammar O, Houas Z, Mehdi M. The association between iron, calcium, and oxidative stress in seminal plasma and sperm quality. *Environ Sci Pollut Res Int* 2019; 26: 14097–14105.
- [12] Fallah A, Mohammad-Hasani A, Colagar AH. Zinc is an essential element for male fertility: A review of Zn roles in men's health, germination, sperm quality, and fertilization. *J Reprod Infertil* 2018; 19: 69–81.
- [13] Biswas TK, Pandit S, Mondal S, Biswas SK, Jana U, Ghosh T, et al. Clinical evaluation of spermatogenic activity of processed Shilajit in oligospermia. *Andrologia* 2010; 42: 48–56.
- [14] Morselli MG, Colombo M, Faustini M, Luvoni GC. Morphological indices for canine spermatozoa based on the World Health Organization laboratory manual for human semen. *Reprod Domest Anim* 2019; 54: 949–955.
- [15] Mirmiran P, Hosseini Esfahani F, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public Health Nutr* 2010; 13: 654–662.

- [16] Wilhelm AL, Maquivar MG, Bas S, Brick TA, Weiss WP, Bothe H, et al. Effect of serum calcium status at calving on survival, health, and performance of postpartum Holstein cows and calves under certified organic management. *J Dairy Sci* 2017; 100: 3059–3067.
- [17] Chamberlin WG, Middleton JR, Spain JN, Johnson GC, Eilersieck MR, Pithua P. Subclinical hypocalcemia, plasma biochemical parameters, lipid metabolism, postpartum disease, and fertility in postparturient dairy cows. *J Dairy Sci* 2013; 96: 7001–7013.
- [18] Rosol TJ, Capen C. Calcium-regulating hormones and diseases of abnormal mineral (calcium, phosphorus, magnesium) metabolism. In: Kaneko JJ, Harvey JW, Bruss ML. Clinical biochemistry of domestic animals. 5th Ed. San Diego: Academic Press; 1997.
- [19] Eslamian G, Amirjannati N, Rashidkhani B, Sadeghi MR, Hekmatdoost A. Nutrient patterns and asthenozoospermia: A case-control study. *Andrologia* 2017; 49: e12624.
- [20] Afeiche M, Williams PL, Mendiola J, Gaskins AJ, Jørgensen N, Swan SH, et al. Dairy food intake in relation to semen quality and reproductive hormone levels among physically active young men. *Hum Reprod* 2013; 28: 2265–2275.
- [21] Afeiche MC, Bridges ND, Williams PL, Gaskins AJ, Tanrikut C, Petrozza JC, et al. Dairy intake and semen quality among men attending a fertility clinic. *Fertil Steril* 2014; 101: 1280–1287.
- [22] Ross C, Morriss A, Khairy M, Khalaf Y, Braude P, Coomarasamy A, et al. A systematic review of the effect of oral antioxidants on male infertility. *Reprod Biomed Online* 2010; 20: 711–723.
- [23] Showell MG, Mackenzie-Proctor R, Brown J, Yazdani A, Stankiewicz MT, Hart RJ. Antioxidants for male subfertility. *Cochrane Database Syst Rev* 2014; 12: CD007411.
- [24] Aitken RJ, Gibb Z, Baker MA, Drevet J, Gharagozloo P. Causes and consequences of oxidative stress in spermatozoa. *Reprod Fertil Dev* 2016; 28: 1–10.
- [25] Gharagozloo P, Aitken RJ. The role of sperm oxidative stress in male infertility and the significance of oral antioxidant therapy. *Hum Reprod* 2011; 26: 1628–1640.
- [26] Ahsan U, Kamran Z, Raza I, Ahmad S, Babar W, Riaz M, et al. Role of selenium in male reproduction: A review. *Anim Reprod Sci* 2014; 146: 55–62.
- [27] Molloy AM. Genetic aspects of folate metabolism. *Subcell Biochem* 2012; 56: 105–130.
- [28] Wong WY, Merkus HMWM, Thomas ChMG, Menkveld R, Zielhuis GA, Steegers-Theunissen RPM. Effects of folic acid and zinc sulfate on male factor subfertility: A double-blind, randomized, placebo-controlled trial. *Fertil Steril* 2002; 77: 491–498.
- [29] Aarabi M, Christensen KE, Chan D, Leclerc D, Landry M, Ly L, et al. Testicular MTHFR deficiency may explain sperm DNA hypomethylation associated with high dose folic acid supplementation. *Hum Mol Genet* 2018; 27: 1123–1135.
- [30] Chavarro JE, Mínguez-Alarcón L, Mendiola J, Cutillas-Tolín A, López-Espín JJ, Torres-Cantero AM. Trans fatty acid intake is inversely related to total sperm count in young healthy men. *Hum Reprod* 2014; 29: 429–440.
- [31] Fair S, Doyle DN, Diskin MG, Hennessy AA, Kenny DA. The effect of dietary n-3 polyunsaturated fatty acids supplementation of rams on semen quality and subsequent quality of liquid stored semen. *Theriogenology* 2014; 81: 210–219.
- [32] Bachir BG, Jarvi K. Infectious, inflammatory, and immunologic conditions resulting in male infertility. *Urol Clin North Am* 2014; 41: 67–81.
- [33] Morielli T, O'Flaherty C. Oxidative stress impairs function and increases redox protein modifications in human spermatozoa. *Reproduction* 2015; 149: 113–123.
- [34] Ferramosca A, Moscatelli N, Di Giacomo M, Zara V. Dietary fatty acids influence sperm quality and function. *Andrology* 2017; 5: 423–430.
- [35] Gürlür H, Calisici O, Calisici D, Bollwein H. Effects of feeding omega-3-fatty acids on fatty acid composition and quality of bovine sperm and on antioxidative capacity of bovine seminal plasma. *Anim Reprod Sci* 2015; 160: 97–104.