

How to Cite:

Sodhi, J. K. ., Shrivastava, B., Lamba, H. S., Kaur, A., & Sharma, P. K. (2022). *Caesalpinia crista* seeds-in vivo screening of hepatoprotective efficacy against CCL4 induced hepatotoxicity. *International Journal of Health Sciences*, 6(S1), 10330–10340. <https://doi.org/10.53730/ijhs.v6nS1.7446>

***Caesalpinia crista* seeds-in vivo screening of hepatoprotective efficacy against CCL₄ induced hepatotoxicity**

Jaspreet Kaur Sodhi

H R Institute of Pharmacy, Ghaziabad - 201003, Uttar Pradesh, India and School of Pharmaceutical Sciences, Jaipur National University, Jaipur - 302017, Rajasthan, India
Corresponding author email: jassodhi.kaur@gmail.com

Birendra Shrivastava

School of Pharmaceutical Sciences, Jaipur National University, Jaipur - 302017, Rajasthan, India

Hardarshan Singh Lamba

Spectrum Institute of Pharmaceutical Sciences & Research, Greater Noida - 201309, Uttar Pradesh, India

Avneet Kaur

Gurugram University, Gurugram– 122003, Haryana, India

Pankaj Kumar Sharma

Apex Institute of Pharmaceutical Sciences, Apex University, Jaipur, 302020, Rajasthan, India

Abstract---Ethanolic extract of *Caesalpinia crista* (commonly known as Latakaranja) seeds, belonging to family Caesalpiniaceae, was evaluated for hepatoprotective efficacy in Wistar rats against hepatic injury carbon induced tetrachloride. The toxicant CCl₄ (1:1 mixture in olive oil) was administered to rats (3 mL/kg) *i.p.*, twice a week for 4 weeks to induce liver toxicity. The ethanolic extract of seeds of *Caesalpinia crista* was administered at doses of 150 and 300 mg/kg *p.o.*, daily for 4 weeks and Silymarin (100 mg/kg *p.o.*) was used as a standard drug. Biochemical parameters such as serum alanine transaminase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP) and total bilirubin were assessed and histopathological studies of the liver were performed to evaluate the hepatoprotective efficacy of the extract. Treatment of rats administered with *C. crista* seed extract showed a significant decrease in CCl₄-induced elevated serum enzyme levels as well as a significant

increase in total protein levels that was lowered by the hepatotoxic compound used and the results were comparable to Silymarin.

Keywords---*Caesalpinia crista*, Wistar rats, carbon tetrachloride, hepatoprotective efficacy.

Introduction

The liver is the body's largest internal organ as well as the largest gland. At the same time it serves as a metabolic as well as a biochemical transformation factory. It receives blood carrying substances via the portal vein and oxygen-rich blood via the hepatic artery (Muriel). As a result, the liver is referred to as a "bidirectional biofilter" (Arias) that is capable of performing a wide range of tasks (mohan and Muriel).

As the liver assists nearly every organ in the body; is imperative for endurance and due to its essential positioning and multifaceted functionalities, it is additionally inclined to numerous illnesses (Eswar). Also as the liver's role is to eliminate substances through the portal circulation, it is vulnerable to constant attack by exogenous agents, resulting in hepatic diseases (Bodakhe). Hepatic problems thus continue to be one of the most serious threats to overall health, posing serious health concerns around the world (Asha).

Extended drug treatment, unrestrained utilization of a number of regularly utilized medicines like paracetamol or acetaminophen, diclofenac and so forth; liquor addiction, vulnerability to particular xenobiotics, toxins and diseases influence liver working to a certain extent and sooner or later result in several liver disorders like hepatitis, cirrhosis, cancer and alcoholic hepatic disorder (Adewusi and Afolayan, 2010).

Due to the absence of any compelling medication that can restore hepatic functions, provide safety from harm or help in recovering hepatic cells (Chattopadhyay, 2003); inaccessibility of well-advised treatment in present day medication and no or extremely more negative impact of synthetic medications in liver-damage; have encouraged specialists in this field to anticipate natural medications with better hepatoprotective activity (Ravishankar and Bhavsar, 1993). Powerful and intense herbal medications require assessment by standard scientific techniques in order to be approved for the treatment of an illness (Jannu et al., 2012).

Utilizing natural medications for treating hepatic disorders has a long background and medicinal plants alongwith their derivatives are as yet utilized everywhere in the world in one form or the other for this reason. Hepatoprotective plants contain an assortment of chemical components like phenols, coumarins, monoterpenes, glycosides, alkaloids and xanthenes (Bhawna and Kumar, 2009). The retardation of free radical generation also can provide an effortless model for assessing the action of hepatoprotective agents (Mohamed Saleem et al., 2008).

Caesalpinia crista, commonly known as Fever nut or Latakaranja (Khare) has been traditionally used in Ayurveda to treat a vast range of diseases like fever, malaria,

skin diseases, and inflammation. It is found to possess anthelmintic [Jabbar], antidiabetic [Gupta], analgesic and antipyretic [Ishan; Gill], wound healing [Patil], antiulcer [Chauhan], antibacterial and antiviral [Afrin; Patil], and cardioprotective [Kumar] activities. Different parts of the plant like leaves, flowers, fruit, root, bark, seeds and seed oil were used medicinally [Suryawanshi]. Taking into account the physiologically active nature of *Caesalpinia crista*, the current study seeks to assess the hepatoprotective potential of *C. cristaseeds* against carbon tetrachloride-induced hepatotoxicity.

Materials and Methods

Chemicals

All chemicals and solvents utilized in the study were of analytical grade; DPPH (Sigma Chemicals Co.); and Silymarin (Micro Labs Ltd) were procured from local supplier of Ghaziabad. The assay kits employed for the estimation of various biochemical parameters were acquired from ERBA (Himachal Pradesh, India).

Plant collection and authentication

The seeds of *Caesalpinia crista* were procured from the local market of Old Delhi, India. The seeds of the plant were authenticated by an emeritus scientist, Dr. Sunita Garg of Raw Material Herbarium and Museum (RHMD), CSIR- National Institute of Science Communication and Information Resources, Delhi (taxonomic reference number: NISCAIR/RHMD-3632-33).

Preparation of the extract

The procured seeds of *Caesalpinia crista* were washed properly and dried. They were then ground to powder, first with the help of pestle-mortar and then an electric grinder. The powdered seeds were subjected to continuous extraction with hot ethanol using Soxhlet apparatus [Harborne, Evans]. The extract obtained, denoted as EECC, was filtered and the filtrate was made solvent free using a vacuum evaporator, yield was calculated and then preliminary phytochemical analysis was performed [Bulugahapitiya].

Total phenolic and flavonoid content

Total phenolic content of ethanolic extract of *Caesalpinia crista* was determined using Folin-Ciocalteu reagent method with some modifications, [Sen et al (2013)] taking gallic acid as standard. All tests were done in triplicate and the result of total phenolic content for all plant extracts was expressed as milligrams of gallic acid equivalent (GAE) to per gram of extract (mgGAE/ g). Similarly total flavonoid content of EECC was determined spectrophotometrically by the method using aluminium chloride [Sandip et al (2014)] with light modifications. The samples were prepared in triplicate and average absorbance was noted. Concentration of total flavonoid contents in the extracts was expressed as milligram Quercetin Equivalents (QE) per gram of the plant extract (mg of QE/ g of plant extract).

In vitro antioxidant activity

To study the antioxidant efficacy, the DPPH scavenging potential of *Caesalpinia crista* seed extract was determined (sodhi). Ascorbic acid was used as a positive control, due to its well-known antioxidant properties. The bleaching of DPPH indicates the free radical scavenging capacity of ethanolic seed extract of *Caesalpinia crista*, the activity being highest at $71.6 \pm 0.42\%$ as compared to the standard ascorbic acid, $84.6 \pm 0.45\%$ by DPPH free radical scavenging method. As a result, the findings demonstrated that EECC could work as a natural antioxidant as well as a free radical inhibitor or scavenger.

Animal studies

Wistar rats of either sex weighing between 300-350 g, were acquired from the animal house of MIET, Meerut, Uttar Pradesh, India. They were housed in polypropylene cages with dust free rice husk as a bedding material and maintained under standard laboratory conditions (controlled temperature of $23 \pm 2^\circ\text{C}$, relative humidity of $40 \pm 10\%$ and natural, 12 hour each, light-dark cycle) for acclimatization for seven days prior to and during experimentation as per guidelines of Institutional Animal Ethics Committee. They were fed with standard rodent pellet diet and water *ad libitum*. The research protocol of this study was approved by IAEC of Meerut Institute of Engineering And Technology (registration no. IAEC/MIET/2021/27).

Acute toxicological studies

Wistar rats (n=3) were selected by random sampling technique and OECD-423 guidelines (OECD) were followed to study the acute oral toxicity. The ethanolic extract was administered orally starting with 5 mg/kg body weight up to 2000 mg/kg body weight. Initially for the first an hour after administering the dose, individual animals were monitored, at least once and then regularly throughout the first 24 hours, with specific care being given during the first 4 hours, every-day after that for a total of 14 days.

Hepatoprotective activity against carbon tetrachloride induced hepatotoxicity

Carbon tetrachloride, a hepatotoxin, was used to induce liver injury and Silymarin was used as reference standard. The animals were randomly divided into 5 different groups (n=6) as follows:

Group I: Served as normal control, normal saline (10 ml/kg of body wt.) p.o. was administered to rats daily for 4 weeks.

Group II: Served as disease control, CCl_4 dissolved in olive oil (1:1) was administered to rats (3 mL/kg of body wt.) intraperitoneally, twice a week for 4 weeks.

Group III: Carbon tetrachloride (CCl_4) was pre-administered as in group 2 to rats followed by Silymarin (100 mg/kg p.o.) daily for 4 weeks.

Group IV: CCl_4 was pre-administered as in group 2 to rats followed by ethanolic extract of stem bark of *Caesalpinia crista* (150 mg/kg p.o.) daily for 4 weeks.

Group V: CCl_4 was pre-administered as in group 2 to rats followed by ethanolic extract of seeds of *Caesalpinia crista* (300 mg/kg p.o.) daily for 4 weeks.

After 24 hours of the last dose, the animals were euthanized and liver and blood samples were collected for the biochemical assessment and histopathological analysis.

Assessment of biochemical parameters

Various biochemical parameters were evaluated for assessing liver functions where estimation of serum enzymes like SGOT, SGPT (Reitman) and serum alkaline phosphatase (Burtis); total bilirubin, total proteins (Dumas, Tietz), serum albumin (Dumas) and TBARS (Wills, Ohkawa) was performed to study the effect of EECC on liver toxicity induced by CCl₄.

Histopathological analysis

The liver tissues fixed in 10% formalin, were dried in graduated amounts of ethanol before being submerged in xylene and then embedded in paraffin. 4 µm thick sections were prepared and stained with haematoxylin and eosin dye (Clayden) to observe gross histopathological changes.

Statistical analysis

The results were given as mean ± standard error mean (SEM). The significance of difference among the control group and various treated groups was analysed using analysis of variance (ANOVA) and $P < 0.05$ was considered statistically significant. The data was statistically analysed using GraphPad Prism 5 software (GraphPad Inc., La Jolla CA).

Results and Discussion

In the Indian system of medicine, a variety of plants are believed to provide relief from liver problems but the claimed medicinal reputation of such herbs' must be scientifically validated. Thus in the current research work, the ethanolic extract of *Caesalpinia cristaseeds* was evaluated for its hepatoprotective potential. The percentage yield obtained from the ethanolic extract was found to be 8.1% and the preliminary phytochemical evaluation revealed the presence of plant secondary metabolites such as alkaloids, flavonoids, tannins, carbohydrates, terpenoids and amino acids in the ethanolic seed extract of *Caesalpinia crista* (table 1). The total phenolic and flavonoid content found to be present in ethanolic extract of *C. crista* seeds was 60.45 ± 8.76 mgGAE/g and 11.78 ± 1.02 mgQE/g, respectively.

Table 1
Preliminary phytochemical screening of ethanolic extract of *Caesalpinia cristaseeds*

Phytochemical Group	EECC
Flavonoids	+
Phenols	+
Alkaloids	+

Tannins	+
Steroids and sterols	-
Terpenoids	+
Glycosides	+
Saponins	-
Carbohydrates	+
Amino acids and proteins	+

+ = Presence of active constituents

- = Absence of active constituents

Considering the presence of greatest amount of phytoconstituents, the ethanolic seed extract of *C. crista* was tested for its hepatoprotective efficacy against carbon tetrachloride. It was found to be nontoxic up to a dose of 2000mg/kg with no signs of mortality in test animals. CCl₄, a best known example of a hepatotoxin, on metabolising forms a hazardous trichloromethyl radical, which causes necrosis and liver failure.

It was perceived that the rats treated with carbon tetrachloride developed a significant hepatic damage as seen from augmented serum levels of hepato-specific enzymes like ALT, AST, ALP, and total bilirubin; and lowered serum total protein levels when compared to normal control. Pre-treatment with Silymarin and EECC showed good protection against carbon tetrachloride induced toxicity to liver. Tests indicate a significant reduction in augmented serum enzyme levels and a considerable increase in total protein levels with extract treated animals as evident from table 2.

Lipid peroxidation study showed that CCl₄ treated group exhibited significant increase in malondialdehyde level when compared to normal control group whereas EECC was significantly able to lower this rise in MDA level as evident from table 2.

Table 2
Effect of ethanolic extracts of stem bark of *Caesalpinia crista* (EECC) on ALT, AST, ALP, total bilirubin, total proteins, serum albumin and LPO levels in carbon tetrachloride induced hepatotoxicity in rats

Group	Serum ALT Levels (IU/L)	Serum AST Levels (IU/L)	Serum ALP Levels (IU/L)	Total bilirubin Levels (mg/dl)	Total protein levels (g/dl)	Serum albumin (g/dl)	LPO (nMole of MDA/mg of protein)
1	39.88±5.52	41.77±5.42	101.66±1.88	0.72±0.09	7.59±0.86	4.61±0.53	0.78±0.08
2	82.67±10.67	91.35±10.25	226.91±4.34	1.81±0.19	3.06±0.41	1.51±0.18	2.63±0.27
3	51.88±4.96***	61.62±6.40***	127.54±3.45***	1.01±0.14**	5.58±0.3***	3.68±0.40***	1.55±0.18**
4	62.86±	74.65±	171.77±	1.16±	3.99±	2.91±	1.98±

	6.29*	9.32*	17.58*	0.11*	0.4**	0.21**	0.2*
5	39.7±	55.87±	125.23±	0.96±	5.97±	3.76±	1.09±
	4.34***	6.1**	12.06***	0.083***	0.45*	0.31***	0.09***

Values are mean \pm SEM (n=6) one way ANOVA. Where, * represents significant at $p < 0.05$, ** represents highly significant at $p < 0.01$, *** represents very significant at $p < 0.001$. All values are compared with toxicant.

Histopathological studies of the liver in carbon tetrachloride induced hepatotoxicity

The histopathological evaluation of carbon tetrachloride toxicity in all the groups was examined and shown in figure 1. The rat liver section of normal control group showed liver parenchyma with intact architecture (the usual morphology) whereas the architecture of the liver in the disease control group was partially effaced, with certain hepatocytes showing apoptotic changes, perivenular mononuclear inflammatory infiltration, and scattered inflammatory infiltration within the parenchyma as signs of toxicity. On the other hand, liver section of Silymarin treated group showed liver parenchyma with intact architecture, though some of the central veins showed congestion with diffuse congestion of sinusoids. Also, the liver section in EECC treated groups (150 and 300 mg/kg) showed intact architecture, few regenerative hepatocytes, sinusoidal congestion and scattered mononuclear inflammatory cells that is similar to Silymarin treated group.

Based on these findings, tannins and flavonoids found to be present in EECC may be responsible for the hepatoprotective efficacy. The outcomes of this study, supported by histological findings, showed that *C. cristata* promises to be a pharmacologically effective treatment for a variety of liver illnesses, demonstrating an improvement in liver function, similar to Silymarin.




Figure 1: Histopathology of the rat liver in carbon tetrachloride induced hepatotoxicity

Conclusion

Pursuant to the results, treatment with an ethanolic extract of *Caesalpinia acrista* seeds afforded considerable protection against CCl₄-induced hepatotoxicity. In contrast to 150 mg/kg, the hepatoprotective potential of 300 mg/kg was somewhat higher. The extract's therapeutic effects were endorsed by histological findings.

Conflicts of interest: the authors declare no conflicts of interest.

Acknowledgements

The authors would like to thank Dr. Vipin Garg, Professor and Head at Meerut Institute of Engineering and Technology (MIET) for providing facilities to carry out the pharmacological activity.

Reference

1. Muriel P. Liver Pathophysiology- Therapies and Antioxidants. Academic Press, Elsevier, London: 2017.
2. Arias IM, Alter HJ, Boyer JL, Cohen DE, Shafritz DA, Thorgeirsson SS, Wolkoff AW. The Liver-biology and Pathobiology, 6th Ed. Hoboken NJ: Wiley-Blackwell, 2020.
3. Mohan H. Textbook of Pathology, 6th Edition. Jaypee Brothers Medical Publishers; India: 2010. Pp: 592-648.
4. EswarKA, Susmitha K, Swathy B, Ramu E, Venkatesh B. A review on liver disorders and screening models of hepatoprotective agents. International Journal of Allied Medical Sciences and Clinical Research. 2014; 2(2):136-150.
5. Bodakhe SH, Ram A. Hepatoprotective Properties of *Bauhinia variegata* Bark Extract. Yakugaku Zasshi. 2007; 127:1503-1507.
6. Asha VV, Pushpangadan P. Preliminary evaluation of the anti-hepatotoxic activity of *Phyllanthus kozhikodanus*, *Phyllanthus maderaspatensis* and *Solanum indicum*, Fitoterapia. 1998; 59: 255-259.
7. Adewusi EA and Afolayan AJ. A review of natural products with hepatoprotective activity. Journal of Medicinal Plants Research. 2010; 4(13):1318-1334.
8. Chattopadhyay RR. Possible mechanism of hepatoprotective activity of *Azadirachta indica* leaf extract: Part II. Journal of Ethnopharmacology. 2003; 89: 217-219.
9. Ravishankar SB, Bhavsar GC. Plants with hepatoprotective activity. Indian Drugs. 1993; 30: 363.
10. Jannu V, Baddam PG, Boorgula AK, Jambula SR. A Review on Hepatoprotective Plants. International Journal of Drug Development and Research. 2012; 4(3):1-8.
11. Bhawna S, Kumar SU. Hepatoprotective activity of some indigenous plants. International Journal of PharmTech Research. 2009; 4:1330-1334.
12. Mohamed Saleem TS, Christina AJM, Chidambaranathan N, Ravi V, Gauthaman K. Hepatoprotective activity of *Annona squamosa* Linn. on experimental animal model. International journal of applied research in natural products. 2008; 1(3):1-7.
13. Khare CP. Indian Medicinal Plants- An Illustrated Dictionary. Springer Science and Business Media, LLC; 2007. p. 116-17.
14. Jabbar A, Zaman MA, Iqbal Z, Yaseen M, Shamim A. Anthelmintic Activity of *Chenopodium album* (L.) and *Caesalpinia crista* (L.) against trichostrongylid nematodes of sheep. Journal of Ethnopharmacology. 2007; 114(1): 86-91.
15. Gupta N, Sharma I, Agarwal M, Mohammed SM, Chauhan P, Anwer T, Khan G. Antidiabetic activity of seed extracts of *Caesalpinia crista* Linn. in experimental animals. Afr J Pharm Pharmacol, 2013; 7: 1808- 1813.

16. Ishan S, Nakul G, Safhi MM, Agrawal M, Prerna C. Antipyretic activity of *Caesalpinia crista* Linn. seeds extract in experimental animals. International Journal of Current Research, 5(5), 2013, 1202-1205.
17. Gill NS, Kaur R, Arora R, Bali M. Phytochemical investigation of *Caesalpinia crista* seed extract for their therapeutic potential. Res J Med Plants, 2012; 6: 100-107.
18. Patil KS. Wound healing activity of the seed kernels of *Caesalpinia crista* Linn. J Nat Remed, 2005; 5: 26- 30.
19. Chauhan P, Gupta N, Safhi MM, Nomier Y, Agarwal M, Nayem M. Evaluation of anti-ulcer activity of *Caesalpinia crista* Linn. seeds on pylorus ligation and indomethacine induced gastric lesions in albino rats. Int J Res Pharm Sci, 2015; 5: 9-13.
20. Afrin S, Pervin R, Sabrin F, Sohrab MH, Rony SR, Islam ME *et al.* Assessment of antioxidant, antibacterial and preliminary cytotoxic activity of chloroform and methanol extracts of *Caesalpinia crista* L. leaf. Bangladesh Journal of Botany 2016;45(5):1061-8.
21. Patil U, Sharma MC. Antiviral activity of lathakaranja (*Caesalpinia crista* L.) crude extracts on selected animal viruses. Global J Res Med Plants Indigen Med 2012;1(9):440-7.
22. Kumar SR, Kumar SA. Cardio protective effect of *Caesalpinia crista* Linn. on isoproterenol induced myocardial necrosis in rats. Int J Res Pharm Sci, 2013;3:119-30.
23. Suryawanshi HP, Patel MR. Traditional uses, Medicinal and Phytopharmacological Properties of *Caesalpinia crista* Linn –An overview. International Journal of Research in Pharmacy and Chemistry 2011;1(4):1179-83.
24. Harborne JB: Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis. Chapman and Hall Ltd, 1973.
25. Evans WC: Trease and Evans Pharmacognosy. Saunders, Elsevier, Edition 16, 2009: 136.
26. Bulugahapitiya VP: Plants Based Natural products: Extraction, Isolation and Phytochemical screening methods. First Edition 2013.
27. Sen S, De B, Devanna N, Chakraborty R. Total Phenolic, Total Flavonoid Content, and Antioxidant Capacity of the Leaves of *MeynaspineseRoxb.*, an Indian Medicinal Plant. Chin. J. Nat. Medicines. 2013; 11(2): 0149-0157.
28. Sandip P, Shreewardhan R, Sandeephan M, Abhay C, Debjani D. Phytochemical evaluation and free radical scavenging potential of *HugoniaMystax* (L) leafextract. Bionano Frontier. 2014; 7(2): 20.
29. Sodhi JK, Kaur A, Shrivastava B and Lamba HS, Sharma J: *In vitro* antioxidant activity of ethanolic extract of bark of *Alstonia scholaris*. International Journal of Psychosocial Rehabilitation 2020; 24(04):10502-10509.
30. OECD. Test No. 423: Acute Oral toxicity - Acute Toxic Class Method, OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, 2002. <https://doi.org/10.1787/9789264071001-en>.
31. Reitman S and Frankel S: A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. American Journal of Clinical Pathology 1957; 28(1):56-63.
32. Burtis CA, Ashwood ER andBruns DE: Textbook of Clinical Chemistry and Molecular Diagnostics. WB Saunders Comp., Fifth Edition 2012.

33. Doumas BT: Standards for Total Serum Protein Assays- A Collaborative Study. *Clinical Chemistry* 1975; 21(8):1159-1166.
34. Tietz NW: Textbook of Clinical Chemistry. WB Saunders, Philadelphia, 1986: 579.
35. Doumas BT, Arends RL, Pinto PC: Standard Methods of Clinical Chemistry. Academic Chicago Press, Vol. VI, 1972: 175-189.
36. Wills ED: Lipid peroxide formation in microsomes. General considerations. *Biochemical Journal* 1969; 113(2):315-324.
37. Ohkawa H, Ohishi N and Yagi K: Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry* 1979; 95:351-358.
38. Clayden EC: Practical Section Cutting and Staining. Churchill Livingstone: Edinburgh, Fifth Edition 1971.