

**How to Cite:**

Kumar, V. K., & Singh, A. (2022). A stability indicating RP-HPLC method for related impurities of dexamethasone in tobramycin and dexamethasone otic suspension. *International Journal of Health Sciences*, 6(S6), 119–132. <https://doi.org/10.53730/ijhs.v6nS6.9177>

# **A stability indicating RP-HPLC method for related impurities of dexamethasone in tobramycin and dexamethasone otic suspension**

**Vala Kiran Kumar\***

\*Research Scholar, Bhagwant University, Ajmer, Rajasthan, India-305004

Email: [kitturx2014@gmail.com](mailto:kitturx2014@gmail.com)

**Dr. Anoop Singh**

Research Supervisor, Department of Pharmaceutics, Bhagwant University, Ajmer, Rajasthan, INDIA-305004

**Abstract---**A Stability Indicating RP-HPLC Method for Related Impurities of Dexamethasone in Tobramycin and Dexamethasone Otic Suspension USP 0.3% and 0.1%. This RP-HPLC method is also validated for various parameters as per ICH guidelines. The system suitability parameters proved that the method is suitable for the quantification of Dexamethasone. System suitability parameters were within the limits as indicated by good resolution. The precision was within the acceptance criteria of %RSD i.e., not more than 2%. Linearity was observed in concentration range of LOQ to 200% and the Correlation coefficient was found to be within the limit. Accuracy was performed with the concentration ranges 50%, 100%, and 200% and was found to be within the limit i.e., 85 to 115%. Stability was evaluated by subjecting the ophthalmic suspension to thermal, acidic, basic, oxidative, and UV stress condition. Hence it can be concluded that the proposed method was a good approach for obtaining reliable results and can be used as a quality – control tool for routine analysis of Dexamethasone in Tobramycin and Dexamethasone Otic Suspension USP 0.3% and 0.1%.


**Keywords---**Otic, Stability, quantification, thermal, acidic, stress condition, Suspension, USP

## **1. Introduction**

Tobramycin (Russ H et al.,) (Fig-1) is an amino-glycoside, broad-spectrum antibiotic produced by *Streptomyces tenebrarius*. Tobramycin can be used in topical or systemic treatment. It is effective against gram-negative bacteria, especially the *pseudomonas* species. It is a 10% component of the antibiotic


complex, produced by the same species. It is chemically (2S,3R,4S,5S,6R)-4-amino-2-[[[(1S,2S,3R,4S,6R)-4,6-diamino-3-[[[(2R,3R,5S,6R)-3-amino-6-(aminomethyl)-5-hydroxyoxan-2-yl]oxy]-2-hydroxycyclohexyl]oxy]-6-(hydroxymethyl) oxane-3,5-diol.

**Fig No 1 Structure of Tobramycin**



Dexamethasone (Fig. 2) is a kind of corticosteroids and prevents the release of substances in the body that cause inflammation. It is used in the treatment of many conditions including a number of skin diseases, severe allergies and asthma. Dexamethasone chemically is 9-fluoro-11 $\beta$ , 17, 21-trihydroxy-16 $\alpha$ -methylpregna-1, 4-diene, 3, 20-dione. It is a synthetic adrenocortical steroid used to treat many different conditions such as allergic disorders, skin conditions, ulcerative colitis, arthritis, lupus, psoriasis, or breathing disorders. The effects of dexamethasone are frequently seen within a day and last for about three days (Sreelakshmi et al).

**Fig No 1 Structure of Dexamethasone**



From literature survey, it has been observed that few methods found to quantitative analysis for estimation of dexamethasone. Official method for assay of dexamethasone available but with challenging chromatographic conditions and some authors also reported study on dexamethasone (S. Shaikh et al.,). As per ICH guideline a specific and stability-indicating procedure should be included to determine the content of the new drug substance (Thamaraikani et al.,) The main objective of this study is to develop a simple, suitable, cost effective and

environment friendly HPLC method required for analysis of dexamethasone from dexamethasone finished product.

Dexamethasone working standard were a kind gift of Remidium Laboratories, Hyderabad, India. Test samples purchased from market store. HPLC grade Acetonitrile and HPLC Water were purchased from Ultra Fine Chemicals Ltd., India. Analytical grade orthophosphoric acid, HCl, NaOH pellets and H<sub>2</sub>O<sub>2</sub> purchased from Merck, India. High performance liquid chromatographic system (Agilent (1100) Gradient System) equipped with UV-visible detector was used for the analysis.

## 2. Method Development

Various trials performed with respect to mobile phase and stationary phase to optimize the suitable chromatographic conditions. Method was done using Agilent 1260 UV/PDA Detector. A non-polar analytical chromatographic column Waters, Symmetry, C18, 3.9 mm x 150 mm, 5 $\mu$  (or) any equivalent Column was chosen as the stationary phase. With Flow Rate: 1.5 mL/min Run time: For standard: 15mins; For Sample: 60 mins, Column Temperature: 35°C, Detection WL: 254nm and Injection Volume: 50  $\mu$ L

### 2.1 Preparation of Standard Solutions:

**Dexamethasone Standard Stock Preparation:** Accurately weigh about 25mg of Dexamethasone Working Standard transfer to a 25mL volumetric flask. Add 15mL of Acetonitrile and sonicate for 3 mins; make up to volume Acetonitrile. (Concentration: 1 mg/mL).

**Preparation of Standard Solution A:** Transfer 1mL of the Standard stock solution (1mg/mL) into 25mL volumetric flask, make up the volume with diluent (Concentration: 0.04mg/mL).

**Preparation of Standard Solution B:** Transfer 1mL of the Standard solution A (0.04mg/mL) into 10mL volumetric flask, make up the volume with diluent (Concentration: 0.004mg/mL).

#### 2.1.1 Dexamethasone Acetate Imp Stock Solution:

2.0 mg of Dexamethasone Acetate Imp in 1mL of Acetonitrile. (Concentration: 2mg/mL)

**2.1.2 System Suitability Solution preparation:** Transfer 2mL of the Standard stock solution (1mg/mL), 1mL of the Dexamethasone Acetate impurity Stock (2mg/mL) into 10mL volumetric flask, make up the volume with diluent (Concentrations: Dexamethasone 0.2mg/mL and Dexamethasone Acetate impurity 0.2mg/mL).

**2.1.3 Preparation of Sample Solution:** Accurately weigh about 2.0gm of sample in 10mL volumetric flask, add about 4mL of diluent, and sonicate for 5mins, make up volume with diluent and mix well. (Concentrations: 0.2mg/mL).


**2.1.3 Preparation of Placebo:** Accurately weigh about 2.0gm of Placebo in 10mL volumetric flask, add about 4mL of diluent, and sonicate for 5mins, make up volume with diluent and mix well.

## 2.2 Procedure:


Set the chromatographic conditions as described above and equilibrate the column with mobile phase till a stable baseline is obtained. Inject diluent (as

blank) solution in duplicate into the chromatograph and record the chromatogram. Inject standard solution for six times into the chromatograph record the chromatograms and measure the peak areas.

**Fig-3 Chromatogram for Standard:**



**Fig-4 Chromatogram for System suitability:**



**TABLE 1 System Suitability**

S. No	Name of the compound	Retention time(mins)	Area	USP Plate Count	USP Resolution	USP Tailing
1	Dexamethasone	8.76	1574556877	1723	0.00	1.03
2	Dexamethasone Acetate	38.65	641723744	2805	15.89	1.24

### 3. Validations of RP-HPLC method

#### 3.1 System Precision:

##### Preparation of Standard Solutions:

##### Dexamethasone Standard Stock Preparation:

Accurately weighed about 25.25 mg of Dexamethasone Working Standard and transferred to a 25mL volumetric flask. 15mL of Acetonitrile was added and sonicated for 3mins; made up the volume with Acetonitrile. (Concentration 1.0mg/ml).

**Preparation of Standard Solution – A:** Transferred 1mL of the Standard stock solution, into 25mL volumetric flask, made up the volume with Acetonitrile (Concentrations: 0.04mg/mL).

**Preparation of Standard Solution – B:** Transferred 1mL of the Standard solution A into 10mL volumetric flask, made up the volume with diluent (Concentrations: 0.004mg/mL).

**Table 2- System suitability Results**

Dexamethasone	Area
Standard 1	4078673
Standard 2	4095825
Standard 3	4097963
Standard 4	4119881
Standard 5	4123177
Standard 6	4131259
<b>Mean</b>	4107796
SD	20106.1
<b>%RSD</b>	0.5

#### 3.2 Method Precision:

**Table 3- Method Precision Results**


S. No	Weight of Sample (gm)	Diluent added (mL)	Sonication time	9,11 Epoxide Impurity Stock Solution (Conc: 0.1mg/mL)	17-Carboxy-17-deoxy Impurity Stock Solution Conc: 0.052mg/mL)	Dexamethasone-21-Acetate Impurity Stock Solution Conc: 0.1mg/mL)	21-dehydro, 17 deoxy Impurity Stock Solution Conc: 0.1mg/mL)	Made up to volume with Mobile phase
1.	2.02424	2	5min	0.2mL	1mL	0.2mL	0.2mL	10mL
2.	2.00950	2		0.2mL	1mL	0.2mL	0.2mL	
3.	2.02259	2		0.2mL	1mL	0.2mL	0.2mL	
4.	2.02107	2		0.2mL	1mL	0.2mL	0.2mL	

5.	2.01937	2		0.2mL	1mL	0.2mL	0.2mL	
6.	2.02001	2		0.2mL	1mL	0.2mL	0.2mL	

**Table 4 Impurities – Retention Times, Relative Retention Times and Relative Response Factors:**

S.No	Name of compound	Retention time (approx.)	RRT	RRF
1	9,11 Alpha Epoxide	5.93	0.647	1.23
2	17-Carboxy-17 Deoxy	27.41	2.96	1.05
3	Dexamethasone-21-Acetate	40.50	4.37	1.37
4	21-Dehydro-17-deoxy	45.40	4.90	4.67
5	Dexamethasone	9.25	1	1

**Fig 5 Chromatogram for Method Precision**



### 3.3 LOQ Precision

**Table 5 LOQ Dexa-21 Acetate Epoxide**

Dexa-21 Acetate	
Sample_1	128976
Sample_2	143891
Sample_3	143735
Sample_4	136300
Sample_5	146936
Sample_6	137166
AVG	139501
STD	6617.44
%RSD	4.7

**Table 6 LOQ 9,11Alpha**

9,11Alpha Epoxide	
Sample_1	255736
Sample_2	266415
Sample_3	264462
Sample_4	260527
Sample_5	265011
Sample_6	257043
AVG	261532
STD	4456.16
%RSD	1.7

**Table 7 LOQ Dexa-21 Acetate Acetate**

<b>21-Dehydro 17-Deoxy</b>	
Sample_1	96640
Sample_2	81609
Sample_3	86820
Sample_4	90467
Sample_5	86174
Sample_6	82228
AVG	87323
STD	5601.31
%RSD	6.4

**Table 8 LOQ Dexa-21**


<b>17-Carboxy 17-Deoxy</b>	
Sample_1	269867
Sample_2	264402
Sample_3	270675
Sample_4	259174
Sample_5	263390
Sample_6	252663
AVG	263362
STD	6765.88
%RSD	2.6

**3.4 Linearity:**

The linearity calibration curve (Peak area vs. Concentration) of Dexamethasone impurity was checked over the concentration ranges LOQ to 200% the linearity was evaluated by linear regression analysis.


**Table 9 Linearity Dexamethasone**

<b>S. No</b>	<b>Linearity Level (%)</b>	<b>Linearity Preparations</b>	<b>Final concentration (ppm)</b>	<b>Area</b>
		<b>Dexamethasone Standard Solution A</b> (Stock conc. 0.05mg/mL)		
<b>1</b>	<b>LOQ</b>	0.05mg/mL × 0.02ml/20ml	0.05	78006
2	10	0.05mg/mL × 0.02ml/10ml	0.10	136680
3	25	0.05mg/mL × 0.05ml/10ml	0.26	344989
4	50	0.05mg/mL × 0.1ml/10ml	0.51	656934
5	100	0.05mg/mL × 0.2ml/10ml	1.02	1399427
6	200	0.05mg/mL × 0.4ml/10ml	2.04	2481866
<b>Correlation Coefficient I</b>				0.998
<b>Slope</b>				1223009.48
<b>Intercept</b>				38477



**Table 10 Linearity 9,11 Alpha Epoxide**


S. No	Linearity Level (%)	Linearity Preparations	Final concentration (ppm)	Area
		9,11Alpha Epoxide Impurity		
1	LOQ	0.005mg/mL × 0.4ml/ 20ml	0.10	109348
2	10	0.102mg/mL × 0.02ml/ 10ml	0.20	196940
3	25	0.102mg/mL × 0.05ml/ 10ml	0.51	473516
4	50	0.102mg/mL × 0.1ml/ 10ml	1.02	937445
5	100	0.102mg/mL × 0.2ml/ 10ml	2.05	1983247
6	200	0.102mg/mL × 0.4ml/ 10ml	4.10	4054964
Correlation Coefficient I				1.000
Slope				991525.16
Intercept				27084






**Table 11 Linearity 17-Carboxy-17-deoxy Impurity**

S.No	Linearity Level (%)	Linearity Preparations	Final concentration (ppm)	Area
		17-Carboxy-17-deoxy Impurity		
1	LOQ	0.0052mg/mL ×0.4ml/20ml	0.11	242874
2	10	0.052mg/mL ×0.1ml/10ml	0.53	552304
3	25	0.052mg/mL ×0.25ml/10ml	1.32	1525580
4	50	0.052mg/mL ×0.5ml/10ml	2.64	2783789
5	100	0.052mg/mL ×1.0ml/10ml	5.29	5724467
6	200	0.052mg/mL ×2.0ml/10ml	10.57	12456437
<b>Correlation Coefficient I</b>				0.999
<b>Slope</b>				1167719.68
<b>Intercept</b>				-10115

**Table 12 Linearity 17-Carboxy-17-deoxy Impurity**


S. No	Linearity Level (%)	Linearity Preparations	Final concentration (ppm)	Area
		Dexamethasone-21-Acetate Impurity		
1	LOQ	0.005mg/mL ×0.4ml/20ml	0.10	91684
2	10	0.100mg/mL ×0.02ml/10ml	0.20	166163
3	25	0.100mg/mL ×0.05ml/10ml	0.50	423392
4	50	0.100mg/mL ×0.1ml/10ml	1.01	1184468
5	100	0.100mg/mL ×0.2ml/10ml	2.01	1822787
6	200	0.100mg/mL ×0.4ml/10ml	4.02	3609379
<b>Correlation Coefficient I</b>				0.996

<b>Slope</b>	893778.91
<b>Intercept</b>	47428



**Table 13 Linearity Dexamethasone-21-Acetate Impurity**

S.No	Linearity Level (%)	Linearity Preparations	Final concentration (ppm)	Area
		21-dehydro-17-Deoxy Impurity		
1	LOQ	0.005mg/mL × 0.4ml/20ml	0.09	98508
2	10	0.094mg/mL × 0.02ml/10ml	0.19	41088
3	25	0.094mg/mL × 0.05ml/10ml	0.47	117034
4	50	0.094mg/mL × 0.1ml/10ml	0.94	234046
5	100	0.094mg/mL × 0.2ml/10ml	1.88	464864
6	200	0.094mg/mL × 0.4ml/10ml	3.76	1018454
<b>Correlation Coefficient I</b>				0.995
<b>Slope</b>				261785.09
<b>Intercept</b>				8893



From the above observation, the correlation coefficient for the Dexamethasone and their Impurities was found to be within the limit.

### 3.5 Accuracy

Accuracy was performed with the concentration range 50%, 100% and 200% was found to be within the limit i.e., 85 to 115%. The % Recovery of 50%, 100% and 200% was found to be within the limits.

**Table 14 Accuracy Results**

S. No	Sample Name	Concentration (mg/ml)			% Recovery		
		50%	100%	200%	50%	100%	200%
1.	9,11 Epoxide Impurity	0.001	0.002	0.004	88.0	85.3	101.2
2.	17-Carboxy-17-deoxy Impurity	0.026	0.052	0.104	102.9	103.6	101.9
3.	Dexamethasone-21-Acetate Impurity	0.001	0.002	0.004	107.4	97.7	99.1

### 4.0 Forced Degradation Studies:

Degradation studies were carried out as per ICH guidelines. The objective of the study was to find out the degradation products, which in turn help in the establishment of degradation pathways and the intrinsic stability of drug molecule. In order to check the selectivity of the proposed method, degradation studies were carried out by using acidic, basic, neutral, oxidative conditions.

#### 4.1 Acidic degradation:

**Test sample Preparation:** Weighed about 4.03049g of sample and transferred to a 10ml volumetric flask, 6mL of 0.1N HCl was added, sonicated for 10 mins and made up the volume with 0.1N HCl. Mixed the contents well and leave undisturbed for 30mins at 80°C. After cooling 5mL of the above solution was transferred to a 10mL volumetric flask, made up the volume with diluent. This solution was injected into the HPLC system and chromatogram was recorded.

**4.2 Alkali degradation:****Test sample Preparation:**

Weighed about 4.03138g of sample and transferred to a 10ml volumetric flask, 6mL of 0.01N NaOH was added, sonicated for 10 mins and made up the volume with 0.01N NaOH. Mixed the contents well and leave undisturbed for 30mins at 80°C. After cooling 5mL of the above solution was transferred to a 10mL volumetric flask, made up the volume with diluent. This solution was injected into the HPLC system and chromatogram was recorded.

**4.3 Neutral degradation:****Test sample Preparation:**

Weighed about 4.03794g of sample and transferred to a 10ml volumetric flask, 6mL of water was added, sonicated for 10 mins and made up the volume with water. Mixed the contents well and leave undisturbed for 30mins at 80°C. After cooling 5mL of the above solution was transferred to a 10mL volumetric flask, made up the volume with diluent. This solution was injected into the HPLC system and chromatogram was recorded.

**4.4 Thermal degradation:****Test sample Preparation:**

Weighed about 4.03817g of sample (B.No. 40192-003A) and transferred to a 10ml volumetric flask, and leave undisturbed for 30mins at 80°C. After cooling 5mL of the above solution was transferred to a 10mL volumetric flask, made up the volume with diluent. This solution was injected into the HPLC system and chromatogram was recorded.

**4.5 Peroxide degradation:****Test sample Preparation:**

Weighed about 4.04564g of sample (B.No.: 40192-003A) and transferred to a 10ml volumetric flask, 6mL of 0.1%H<sub>2</sub>O<sub>2</sub> was added, sonicated for 10 mins and made up the volume with 0.1%H<sub>2</sub>O<sub>2</sub>. Mixed the contents well and leave undisturbed for 30mins at 80°C. After cooling 5mL of the above solution was transferred to a 10mL volumetric flask, made up the volume with diluent. This solution was injected into the HPLC system and chromatogram was recorded.

**4.6 Photo degradation:**

**Test sample Preparation:** About 4gms of sample was kept under UV Cabinet for 2hrs.

Weighed about 2.02276gms of above sample and transferred to a 10 ml volumetric flask, 7mL of diluent was added, sonicated for 5 mins and made up the volume with diluent. This solution was injected into the HPLC system and chromatogram was recorded.

**Table 15 Peak purity data of Dexamethasone Peak in Forced degradation:**

Condition	Sample Name	Peak Purity	% Assay	Total % of Impurities	Mass Balance
Acid Degradation	Dexamethasone	1.000	88.72	3.01	91.73
Alkali Degradation	Dexamethasone	1.000	87.57	2.03	89.60
Neutral Degradation	Dexamethasone	1.000	86.40	3.14	89.31
Peroxide Degradation	Dexamethasone	1.000	86.17	2.26	96.21
Thermal Degradation	Dexamethasone	1.000	93.95	2.03	88.43
Photo Degradation	Dexamethasone	1.000	92.09	2.05	94.14

## 5. Conclusion

Method was done using Agilent 1260 UV/PDA Detector. A non-polar analytical chromatographic column Waters, Symmetry, C18, 3.9 mm x 150 mm, 5 $\mu$  (or) any equivalent Column was chosen as the stationary phase. With Flow Rate: 1.5 mL/min Run time: For standard: 15mins; For Sample: 60 mins, Column Temperature: 35°C, Detection WL: 254nm and Injection Volume: 50  $\mu$ L

This RP-HPLC method is also validated for various parameters as per ICH guidelines. The system suitability parameters proved that the method is suitable for the quantification of Dexamethasone. System suitability parameters were within the limits as indicated by good resolution. The precision was within the acceptance criteria of %RSD i.e., not more than 2%. Linearity was observed in concentration range of LOQ to 200% and the Correlation coefficient was found to be within the limit. Accuracy was performed with the concentration ranges 50%, 100%, and 200% and was found to be within the limit i.e., 85 to 115%. Hence it can be concluded that the proposed method was a good approach for obtaining reliable results and can be used as a quality – control tool for routine analysis of Dexamethasone in Tobramycin and Dexamethasone Otic Suspension USP 0.3% and 0.1%.

## References

1. <https://www.drugbank.ca/drugs/DB00873>
2. <https://www.drugbank.ca/drugs/DB00684>
3. <https://www.drugs.com/sfx/Tobramycin-side-effects.html>

4. C.A.C. Jessurun, A.F.C. Hulsbergen, L.D. Cho, L.S. Aglio, R.D.S.N. Tewarie and M.L.D. Broekman, J. Neurooncol., 144. 249 (2019); <https://doi.org/10.1007/s11060-019-03238-4>
5. S. Shaikh, H. Verma, N. Yadav, M. Jauhari and J. Bullangowda, ISRN Anesthesiol., 2012, Article ID 985495 (2012); <https://doi.org/10.5402/2012/985495>
6. Dexamethasone USP monograph, USP29-NF24
7. T. Thamaraikani, S. Mounika and S. Vijith, Int. Res. J. Pharm. App. Sci., 2, 5 (2012).
8. International Conference on Harmonization ICH guidelines, Q6A, Test Procedures and Acceptance Criteria for New Drug Substances and New Drug Products: Chemical Substances (2000);
9. Valentini F, Buldini PL, Landi E, Tampieri A and Tonelli D. (2008). HPLC determination of tobramycin in a simulated body fluid. Microchemical Journal, 90(2), 113-7.
10. Comstock TL and Holland EJ. (2010). Loteprednol and tobramycin in combination: a review of their impact on current treatment regimens. Expert opinion on pharmacotherapy, 11(5), 843-52.
11. Mashat M, Chrystyn H, Clark BJ and Assi KH. (2008). Development and validation of HPLC method for the determination of tobramycin in urine samples post-inhalation using pre-column derivatisation with fluorescein isothiocyanate. Journal of Chromatography B, 869(1-2), 59-66.
12. Patel AB and Patel DB. (2013). Development and validation of RP-HPLC method for simultaneous estimation of gatifloxacin and loteprednol etabonate in pharmaceutical dosage form. Asian Journal of Research in Chemistry,
13. Dagneaux PK and Elhorst JK. (1981). HPLC-estimation of tobramycin in serum. Pharmaceutisch weekblad, 3(1),
14. Russ H, McCleary D, Katimy R, Montana JL, Miller RB, Krishnamoorthy R and Davis CW. (1998). Development and validation of a stability-indicating HPLC method for the determination of tobramycin and its related substances in an ophthalmic suspension. Journal of liquid chromatography and related technologies, 21(14), 2165-81.
15. Pfeifer C, Fassauer G, Gerecke H, Jira T, Remane Y, Frontini R, Byrne J and Reinhardt R. (2015). Purity determination of amphotericin B, colistin sulfate and tobramycin sulfate in a hydrophilic suspension by HPLC. Journal of Chromatography B, 990, 7-14.
16. Sreelaksmi et al. Method Development and Validation of Tobramycin and Dexamethasone by RP-HPLC. International Journal for Advances and Research in Pharmaceutical Sciences. 2020; 1(1):25-31