How to Cite:

Sintia Dewi Septiani, M. K., Andriati, A., & Narasinta, I. (2022). A randomized controlled trial of modified traditional Javanese dance on cardiorespiratory fitness in elderly. *International Journal of Health Sciences*, 6(S4), 3132–3140. https://doi.org/10.53730/ijhs.v6nS4.10002

A randomized controlled trial of modified traditional Javanese dance on cardiorespiratory fitness in elderly

Sintia Dewi Septiani M K

Department of Physical Medicine and Rehabilitation Faculty of Medicine Universitas Airlangga, Dr Soetomo General Academic Hospital, Surabaya, Indonesia

Corresponding author email: luv_se3n@yahoo.com

Andriati

Department of Physical Medicine and Rehabilitation Faculty of Medicine Universitas Airlangga, Dr Soetomo General Academic Hospital, Surabaya, Indonesia

Inggar Narasinta

Department of Physical Medicine and Rehabilitation Faculty of Medicine Universitas Airlangga, Dr Soetomo General Academic Hospital, Surabaya, Indonesia

> **Abstract**---The aim of this study is to determine changes of cardiorespiratory fitness in elderly after giving modified Javanese traditional dance for 12 weeks. This study is a experimental study with pre and post-test randomized control group design. The subjects were the elderly woman over 60 years old in the Hargodedali nursing home, Surabaya, Indonesia. Subjects in the treatment group participated in the modified Javanese traditional dance for 12 weeks, 3 times per week with a duration of 35 minutes per session. The control group subjects did not attend the dance session. Six minutes walking test (6MWT) and VO2 max were measured before before treatment and 12 weeks after treatment was finished. There was a statistically significant increase in VO_2 max value of the treatment group (p = 0.01). While in the control group there was a decrease in vo2 max, but not significant (p = 0.221). The effect size of the treatment group is 0.93. There is a significant difference in delta vo2max between the treatment and control groups (p=0.004). The effect size after treatment between groups is 1.4. The adherence of subject in this study is 95.3%. Thus, modified traditional Javanese dance increase VO_2 max for elderly.

International Journal of Health Sciences ISSN 2550-6978 E-ISSN 2550-696X © 2022. Manuscript submitted: 18 Feb 2022, Manuscript revised: 27 April 2022, Accepted for publication: 9 June 2022 3132

Keywords---dance movement therapy, cardiorespiratory fitness, elderly, Javanese dance, VO2max, 6MWT.

Introduction

Globally the population aged over 65 years is 9%, which is increasing over time, due to the increasing life expectancy (United Nation, 2019). Population in Indonesia in 2018, there were around 24.49 million elderly (Kemenkes, 2014). Data shows the elderly morbidity rate in Indonesia in 2018 was 25.99% (BPS, 2018). Elderly brings physiological changes in all organ systems including the cardiorespiratory system. Changes in the cardiovascular and respiratory systems due to aging will cause a decrease in cardiorespiratory fitness. Maximum oxygen consumption (VO₂ max) as an indicator of cardiorespiratory fitness will decrease with age (Cifu, 2018; Bonder & Bello, 2009). Most of the elderly show physical inactivity, and have a sedentary lifestyle. Physical inactivity contributes to many health conditions, including decreased cardiorespiratory fitness as assessed by VO_2 max. Low levels of cardiorespiratory fitness are associated with high rates of associated morbidity and mortality with various etiologies, especially cardiovascular disorders (ACSM, 2018). Exercise or physical activity in the elderly can help overcome this problem, but the compliance and interest of the elderly in physical exercise is still very low. Dancing has the potential to be a good and interesting physical activity for the elderly, which can be adapted to the target population of age, physical limitations, and culture (Hwang & Braun, 2017).

Dance movement therapy (DMT) is an alternative exercise for elderly to improve cardiorespiratory function. DMT can also improve the psychological aspects of the elderly which can lead to attachment and adherence to exercises and therapists (Payne, 2003). Indonesia is a country that has various tribes, customs, races, cultures, and religions. Indonesian traditional dance has the potential to be used as an exercise modality to improve physical fitness (Maghfiroh, 2017; Pradewi, 2012; Pertiwi, 2017). Javanese dance is one of the traditional Indonesian dances originating from the Java area which has its own characteristics and rules, which have a variety of movements that involve all parts of the body, so that it is considered to improve physical fitness, include cardiorespiratory fitness (Poerwanto et al., 2019; Kuswardhani et al., 2020). Modified traditional Javanese dance is a dance modification that uses basic traditional Javanese dance movements, which are adapted to the physical capacity of the elderly. The dance is given with a moderate intensity that has been measured previously in the preliminary study, accompanied by Javanese music that is familiar to the research subject. This study aims to determine changes in the cardiorespiratory function of elderly women at the Hargodedali nursing home Surabaya after receiving modified Javanese traditional dance therapy.

Method

This study is a true experimental randomized control group design with a pre and post intervention design. This study uses simple random sampling. This research received ethical approval from the Health Research Ethics Committee, Faculty of Medicine, Airlangga University with No.26/EC/KEPK/FKUA/2022. All research subjects have signed informed consent.

The study was conducted at the Hargodedali nursing home Surabaya for 12 weeks from February 2022 to May 2022. The subjects of this study were 24 elderlies aged over 60 years. The inclusion criteria for this study are: elderly living in the Hargodedali nursing home, female and 60 years or older, MOCA-Ina score more than 20, able to ambulate independently without assistive devices, stable hemodynamics, good visual and hearing function. The exclusion criteria for this study included: the elderly regularly participated in other forms of physical exercise in the last 3 months, Barthel Index Score less than 80, have dementia, have fractures of the upper and lower limbs that interfere with movement and ambulation, have a chronic disease or disease that causes other long-term disability. Subjects can withdraw from the study because: not willing to continue research for any reason, cannot complete the exercise according to the research protocol, experienced shortness of breath, chest pain and complaints related to heart failure that did not improve with rest or taking medication, and died.

The subjects were randomized into 2 groups, treatment group and the control group. Subjects in the treatment group received modified Javanese traditional dance 3 times per week for 12 weeks with moderate intensity (40-60% resting heart rate). The dance session begins with a warm-up exercise in the form of finger gymnastics, joint movements and stretching for 10 minutes accompanied by a slow-tempo national anthem, a core dance with a gradual increase in complexity of motor sequences for 15 minutes accompanied by folk songs with gradually increasing tempo and cooling down in the form of breathing exercises and stretching for 10 minutes accompanied by a slow tempo national anthem. Prior to the 12-week intervention, the patient was given a one-week preparatory class to introduce dance moves. During the intervention, participants were guided by 2 dance instructors, and supervised by 2 doctors and 2 nursing home staff. Subjects in the control group were given education to continue their daily physical activities as usual.

The parameter evaluated in this study was the level of VO₂ max in each group. Measurements were carried out twice, before and after intervention. The initial assessment to determine VO₂ max was carried out with a six minutes walking test (6MWT). The initial evaluation was carried out one day before preparatory class. The final evaluation was carried out one day after the last session. Statistical analysis of this study using SPSS 23.0. The data normality test was performed using Shapiro-Wilk test. To compare VO2max before and after treatment in each group, paired T test was used. To compare the differences in pre and post VO₂max between the treatment and control groups, the independent sample t test was used. The p value is considered significant if p < 0.05.

Result and Discussion

Characteristic of Subject

The total subjects of this study were 24 people and all subjects were female. There were 12 subjects in both treatment group and control group. In the final

evaluation, 1 person from the treatment group was drop out because they did not participate in the exercise 3 times in a row and 1 person from the control group was dropout due to death. The final number of subjects analyzed were 11 subjects from the treatment group and 11 subjects from the control group. The initial characteristics of the subject, homogeneity and normality test can be seen in the table 1.

Variable	Treatment	p-value	Control	p-value	p-value
	Group (n=12)	(normality)	Group (n=12)	(normality)	(homogenity)
	Mean±SD		Mean±SD		
Age (years)	74,72±7,9	0,458	77,09±6,13	0,138	0,646
Body weight (kg)	50,45±12,23	0,467	45±6,41	0,704	0,115
Body height	153,7±7,47	0,066	148,36±3,04	0,014	0,020
(cm)					
BMI (kg/m ²)	21,19±3,85	0,990	19,97±2,21	0,257	0,243
Sistolic Blood	129,54±14,2	0,048	118,18±16,0	0,005	0,055
Pressure	2		1		
(mmHg)					
Diastolic Blood	73,63±10,26	0,054	75,45±6,87	0,002	0,823
Pressure					
(mmHg)					
Resting heart	67,81±4,11	0,095	67,54±3,14	0,144	0,616
rate					
(times/minutes)					
6MWT baseline	226,36±88,3	0,044	165,45±62,3	0,034	0,112
(m/s)	4		4		
VO2 max	15,51±4,97	0,136	13,85±3,70	0,767	0,671
baseline					
(ml/kg/minutes					

	Table 1		
Demographic and Clinical	Characteristic	Subject a	at Baseline

6MWT = six minute walking test; VO2 max : maximum oxygen consumption; Shapiro-wilk normality test = data distribution is normal if p>0,05; homogenity test = data distribution is homogen if p>0,05

Outcome

The average VO2 max value of the treatment group before being given a modified Javanese traditional dance was $15.51\pm4.97 \text{ ml/kg/minute}$ and the average after being given exercise for 12 weeks was $17.18\pm4.34 \text{ ml/kg/minute}$. There was a significant increase in the VO2 max value in the treatment group (p = 0.01) (table 2). In the control group, the mean VO2 max value at the initial evaluation was $13.87\pm3.70 \text{ ml/kg/min}$, and the mean VO2 max value at the final evaluation was $13.63\pm3.65 \text{ ml/kg/min}$. Based on the paired t test, there was an increase in the VO2 max value in the control group (p = 0.221). The effect size of Modified Javanese Traditional Dance training program for 12 weeks was large (Cohen's D = 0.93).

The mean delta VO₂ max in the treatment group was $1.66\pm1.78 \text{ m/kg/min}$, while the control group was $-0.42\pm1.13 \text{ ml/kg/min}$. There was a significant difference in delta VO2max between the treatment group and the control group (p=0.004) (Table 3). The effect size based on the delta between groups is large (Cohen's D = 1.4). The value of changes in VO2 max before and after giving the modified Javanese traditional dance training program can be seen in table 3. The average attendance rate of the subjects in the study was 95.3%. From the 12 subjects in the treatment group, only one subject was unable to complete the exercise (dropped out) resulting in a high retention rate of 91.7%.

Table 2							
VO ₂ max pre and	post intervention i	n both group					

	Treatment Group (n=11)			Control Group (n=11)		
	Pre	Post	p-value	Pre	Post	p-value
	intervention	intervention		intervention	intervention	
VO2 max	15,51±4,97	17,18±4,34	0,01*	13,87±3,70	13,63±3,65	0,221
(ml/kg/						
minutes)						

Vomax = maximum oxygen consumption; *paired T-test is meaningful if p<0.05

Table 3					
Δ VO ₂ max in both g	group				

	Treatment Group (n=11)		Control Group (n=11)		p-value		
Δ VO2 max	1,66±1,78		-0,42±1,13		0,004*		
(ml/kg/minute)							
Δ VO2 max =	delta	maximum	oxygen	consumption;	*Indepen	dent	T-test
significant if p<0.	05						

Discussion

In this study, the age and BMI characteristics of the treatment group and control group were considered homogeneous. VO₂ max decreases with age, 4-5% in active individuals, and 8-10% per 10 years for inactive individuals (Rodrigues *et al.*, 2006). Study by Kalyanshetti & Veluru (2017), an increase in body mass index can reduce cardiorespiratory fitness by assessing VO₂ max. Obesity increases the risk of fatigue during work, the higher the body mass index, the easier it is to experience fatigue. This is because more heat is produced, the ratio of surface area to body volume is getting smaller and heat insulation by fat (Rodrigues *et al.*, 2006). In this study, there were 3 obese patients in the treatment group with low VO2max value, 2 subject had increasing of VO2max about 2.3%-37%. This shows that most of the elderly with obesity have a tendency of low VO₂ max, but can increase if given aerobic exercise regularly. Study conducted by Ramos *et al* in 2019 which stated that moderate-intensity aerobic exercise can increase VO₂ max in the elderly with obesity (Ramos *et al.*, 2019).

We used 6 minute walk test (6MWT) to calculate VO2max. 6MWT is a valid clinical assessment for assessing cardiorespiratory fitness in the elderly (Kervio et al., 2003). There is some formula to get VO2 max from 6MWT. In this study, we

3136

used the formula from Manttari *et al* in 2018, because it was considered the closest to the characteristics of the subject (Mantari *et al.*, 2018). The results of this study showed that giving a modified Javanese traditional dance training program for 12 weeks was able to significantly increase the VO₂ max value. The average increase in VO₂ max was 1.67 ml/kg/min in the treatment group. The effect size value based on the treatment group's delta is 0.93 which indicates that the modified Javanese traditional dance training program has a large effect on increasing VO₂ max.

The modified Javanese traditional dance training consists of moderate-intensity aerobic exercise (40-60% HRR), 3 times a week for 12 weeks. This prescription consider the initial conditions of the subjects in a very sedentary nursing home. The subjects in this study had not done regular physical exercise since the beginning of the pandemic, which was about 1.5 years. Conventional aerobic exercise that was performed previously was also only given at a low intensity, so we chose a prescription that was more tolerable by the study subjects. The results of this study are in line with research conducted by Hui et al in 2008, showing that the elderly were given dance movement therapy (DMT) exercises 2 times per week, for 12 weeks (50 minutes per session, 10 minutes of warm-up, 30 minutes of core dance, 10 minutes of cooling) can improve the six minute walking test (6MWT) better than the control group that was not given the intervention (Hui et al., 2009). Another study by Eyigor et al in 2009, with the intervention of Turkish folklore dance 3 times per week with a duration of 1 hour per session, for 8 weeks, can improve functional performance including the 6 minute walking test (6MWT) (Eyigor et al., 2009).

Aerobic exercise consists of repetitive low-resistance movements that engage large muscle groups and increase heart rate over a period of time. The increase in VO2 max after aerobic exercise in the elderly ranged from 6.6% - 30% (Kalapatharakos, 2007). In this study, the increase in VO2 max was 10.7%. The author believes that the increase in VO2 max in this study occurs because the modified Javanese traditional dance movements involve large muscle groups, such as the shoulder muscles (lembehan, ukel ngrayung, menthang miwir sampur, ndayung, singget menthang), trunk (ogek lambung, ongkel pundak, oleng, ogel) and lower limbs (*jinjit-ungkit kaki, laku lombo*). These movements are repeated, with the intensity, frequency and duration according to the aerobic exercise prescription for the elderly, so as to increase VO2 max in the elderly after 12 weeks of intervention. Modified Javanese traditional dance also contain elements of music therapy. Music is closely related and forms the basis for the formation of dance movements. This study uses Javanese-themed music according to the characteristics of the research subjects, of which 92% are Javanese. Familiar music will have a more therapeutic effect, especially for the elderly. This is explained through two mechanisms. First, familiar music reduces the demands of cognitive synchronization, so adaptation to movements and beats will be faster. Second, familiar music will make the patient more enjoy the movement so as to achieve the goal of therapy (Leow et al., 2015).

Music therapy can elicit an emotional response because moving with music activates endorphins-related brain's pleasure circuits, and dancing rhythms to music can create a pleasant movement pattern, and distract from fatigue, so that the elderly can follow the intervention according to the instructions and the exercise goals can be achieved (de Dreu *et al.*, 2012). This study resulted in an increase in the average VO2 max in the intervention group of 1.66 ± 1.78 or about 10.7% of the initial VO2 max value. Aerobic exercise is effective if it can increase VO2 max by 6.6% - 30% (Kalapatharakos, 2007). However, increasing VO2 max is said to be more effective if it can increase 1 metabolic equivalent/mets or VO2 max by 3.5 ml/kg/min. An increase of 1 Mets is associated with a 10-25% increase in survival rate, and a reduced risk of cardiovascular disease (Scribbans *et al.*, 2016). So it is necessary to do further research on this intervention with a higher frequency and duration in future studies to be able to increase VO2 max better.

Patient's Adherence

The average attendance rate in the study was quite good, namely 95.3%. Of the 12 subjects in the treatment group, only one could not complete the exercise (dropped out) according to the criteria, resulting in a high retention rate of 91.7%. The drop out subject did not take part in the exercise 3 times in a row because the person concerned did not want to take part in the exercise for no apparent reason. There are no health problems in this subject that cause not following the exercise. Drop out subjects in the treatment group had a Moca-Ina value of 23, which was included in mild cognitive impairment (MCI). In this study no psychological screening was carried out, so that the incidence of drop out due to psychological disorders could not be explained. Factors associated with a high level of adherence to exercise include good socioeconomic status, marital status, better health conditions, better physical abilities, better cognitive abilities, and fewer depressive symptoms (Rivera-Torres *et al.*, 2019).

The modified traditional Javanese dance is an alternative aerobic exercise, which contains elements of a strong Indonesian tradition, so that it can increase the interest of the elderly for participating in exercise regularly. Dance is also reported more fun for the elderly, improves psychological aspects, integrates physical functions, cognitive and social elements, which provides stimulation of balance, coordination, flexibility, and cardiorespiratory function (Meng *et al.*, 2020; Payne, 2003).

Limitation of Study

The limitations of this study include: 1) this study only took elderly female subjects, 2) this study only assessed the parameters before and immediately after exercise, so the long-term effects of this exercise program are not known, 3) not carried out assessment of daily physical activity and screening for psychological disorders in both the intervention group and the control group.

Conclusion

There is an increase in VO_2 max after modified Javanese traditional dance for 12 weeks to elderly women at the Hargodedali nursing home in Surabaya. There is also delta difference in VO_2 max values in the treatment group that received

modified Javanese traditional dance training for 12 weeks compared to the control group.

Acknowledgments

The authors are grateful for the assistance from all parties in the implementation of this research.

References

- American College of Sports Medicine (ACSM). 2018. ACSM's guidelines for exercise testing and prescription. Lippincott Williams & Wilkins.
- Badan Pusat Statistik (BPS). 2018. Statistik Penduduk Lansia 2017. Jakarta: BPS.
- Bonder, B.R., & Bello-Haas, V.D. 2009. Functional performance in older adults, 3rd ed. F.A. Davis Company: Philadelphia.
- Cifu, D. X. 2015. *Braddom's physical medicine and rehabilitation E-book*. Elsevier Health Sciences.
- de Dreu, M. J., Van Der Wilk, A. S. D., Poppe, E., Kwakkel, G., & van Wegen, E. E. (2012). Rehabilitation, exercise therapy and music in patients with Parkinson's disease: a meta-analysis of the effects of music-based movement therapy on walking ability, balance and quality of life. *Parkinsonism & related disorders*, 18, S114-S119. https://doi.org/10.1016/S1353-8020(11)70036-0
- Eyigor, S., Karapolat, H., Durmaz, B., Ibisoglu, U., & Cakir, S. (2009). A randomized controlled trial of Turkish folklore dance on the physical performance, balance, depression and quality of life in older women. *Archives of gerontology and geriatrics*, 48(1), 84-88. https://doi.org/10.1016/j.archger.2007.10.008
- Hui, E., Chui, B. T. K., & Woo, J. 2009. Effects of dance on physical and psychological well-being in older persons. *Archives of gerontology and geriatrics*, 49(1), e45-e50. https://doi.org/10.1016/j.archger.2008.08.006
- Hwang, P. W. N., & Braun, K. L. 2015. The effectiveness of dance interventions to improve older adults' health: a systematic literature review. Alternative therapies in health and medicine, 21(5), 64. Available from: file:///C:/Users/Carla Carolina/Desktop/Artigos para acrescentar na qualificação/The impact of birth weight on cardiovascular disease risk in the.pdf
- Kalapotharakos, V. I. (2007). Aerobic exercise in older adults: effects on VO 2 max and functional performance. *Critical Reviews™ in Physical and Rehabilitation Medicine*, 19(3). https://doi.org/10.1615/CritRevPhysRehabilMed.v19.i3.30
- Kalyanshetti, S. B., & Veluru, S. (2017). A cross-sectional study of association of body mass index and VO[^] sub2[^] max by nonexercise test in medical students. *National Journal of Physiology, Pharmacy and Pharmacology*, 7(2), 228. https://doi.org/10.5455/njppp.2017.7.0825804092016
- Kementrian kesehatan RI. 2019. Infodatin disabilitas. Diakses 20 Januari 2021, dari : https://pusdatin.kemkes.go.id/
- Kervio, G., Carre, F., & Ville, N. 2003. Reliability and intensity of the six minute walk test in healty elderly subjects. *Journal of the American College of Sports Medicine, Medicine & Science in Sport & Exercise* 35(1): 169-74. https://doi.org/10.1249/01.MSS.0000043545.02712.A7

- Kuswardhani, R. T., Lesmana, S. I., Adiatmika, I. P. G., Pangkahila, J. A., & Tianing, N. W. 2020. Perbedaan Pengaruh antara Latihan Gerak Tari Jawa Tengah Modifikasi dengan Latihan Propioseptifterhadap Keseimbangan Dinamis pada Lansia. Sport and Fitness Journal, 8(3), 115-124.
- Leow, L. A., Rinchon, C., & Grahn, J. (2015). Familiarity with music increases walking speed in rhythmic auditory cuing. Annals of the New York Academy of Sciences, 1337(1), 53-61. https://doi.org/10.1111/nyas.12658
- Maghfiroh, I. 2017. Pengaruh Pemanfaatan Gerak Dasar Tari Jawa Terhadap Kualitas Gestur Aktor Teater SS di Universitas Negeri Semarang. Universitas Negeri Semarang
- Mänttäri, A., Suni, J., Sievänen, H., Husu, P., Vähä-Ypyä, H., Valkeinen, H., ... & Vasankari, T. 2018. Six-minute walk test: a tool for predicting maximal aerobic power (VO 2 max) in healthy adults. *Clinical physiology and functional imaging*, 38(6), 1038-1045. https://doi.org/10.1111/cpf.12525
- Meng, X., Li, G., Zhang, G., Yin, H., Jia, Y., Wang, S., ... & Chen, L. (2020). Effects of dance intervention on frailty among older adults. Archives of Gerontology and Geriatrics, 88.

https://doi.org/10.1016/j.archger.2019.104001

- Payne, H. 2003. Dance movement therapy: Theory and practice. NewYork : Brunner Routledge.
- Pertiwi, N. L. T., & Muliarta, I. M. 2017. Persentase Heart Rate Reverse Penari Legong Keraton Lasem pada Mahasiswi Program Studi Seni Tari Institut Seni Indonesia Denpasar Tahun 2014. *E-Jurnal Medika Udayana*, 6(9).
- Poerwanto, A., Haetami, M., & Gustian, U. PENGEMBANGAN GERAK SENAM AEROBIK BERBASIS TARIAN TRADISIONAL. Jurnal Pendidikan dan Pembelajaran Khatulistiwa, 8(10).
- Pradewi, S., & Lestari, W. 2012. Eksistensi Tari Opak Abang Sebagai Tari Daerah Kabupten Kendal. *Jurnal Seni Tari*, 1(1).
- Ramos, R. M., Coelho-Júnior, H. J., Asano, R. Y., Prado, R. C. R., Silveira, R., Urtado, C. B., ... & de Oliveira Assumpção, C. (2019). Impact of moderate aerobic training on physical capacities of hypertensive obese elderly. *Gerontology* and *Geriatric Medicine*, 5. https://doi.org/10.1177%2F2333721419859691
- Rivera-Torres, S., Fahey, T. D., & Rivera, M. A. (2019). Adherence to exercise programs in older adults: informative report. *Gerontology and geriatric medicine*, 5. https://doi.org/10.1177%2F2333721418823604
- Rodrigues, A. N., Perez, A. J., Carletti, L., Bissoli, N. S., & Abreu, G. R. 2006. Maximum oxygen uptake in adolescents as measured by cardiopulmonary exercise testing: a classification proposal. *Jornal de Pediatria*, 82(6), 426-430. https://doi.org/10.1590/S0021-75572006000800006
- Scribbans, T. D., Vecsey, S., Hankinson, P. B., Foster, W. S., & Gurd, B. J. (2016). The effect of training intensity on VO2max in young healthy adults: a meta-regression and meta-analysis. *International journal of exercise science*, 9(2), 230.
- United Nations, World Population Ageing 2019. (2019). World Population Ageing 2019.