

Some fixed point results of F-Contraction mapping in \mathbb{D} -metric spaces by Samet's method

HOJJAT AFSHARI^{a,*}, SEYED MOHAMMAD ALI ALEOMRANINEJAD^b

^a Department of Mathematics, University of Bonab, Bonab 5551761167, Iran.

^b Department of Mathematics, Faculty of Science, Qom University of Technology, Qom 3718146645, Iran.

• Received: 28 June 2021 • Accepted: 17 September 2021 • Published Online: 21 October 2021

Abstract

The aim of this paper is to study the F-contraction mapping introduced by Wardowski to obtain fixed point results by method of Samet in generalized complete metric spaces. Our findings extend the results announced by Samet methods and some other works in generalized metric spaces.

Keywords: F-contraction, \mathbb{D} -metric spaces, Fixed point.

2010 MSC: 74H10, 54H25.

1. introduction

Dhage [11], presented the extended metric or \mathbb{D} -metric spaces and obtained some results about it. Many researchers have taken these results for granted and applied them in studying fixed point results in \mathbb{D} -metric spaces. Rhoades [21], extended Dhages contractive condition by increasing a number of factors and studied the existence and uniqueness of fixed point of some mappings in \mathbb{D} -metric space. Wardowski introduced the concepts of F-contraction and F-weak contraction to generalize the Banach's contraction in many ways (see [28],[29]). Sedghi et al. [26] have introduced the concept of S-metric space and investigated that this is a generalization of a G-metric space and a \mathbb{D}^* -metric space. Also, they have studied properties of S-metric spaces and some fixed point results for a self-map on an S-metric space. In the following, some authors extended this work (see [2, 20, 24, 25]). Samet et al. [23, 22] proved that $\alpha - \phi$ contractions unify large classes of contractive type operators, whose fixed points can be obtained by means of the Picard iteration. Afterward, these results expanded by many mathematicians (see, for example [1, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 17]).

Here, we investigate the result of Wardowski and Samet in generalized applying the result obtained by E. Karapinar in [16] we prove new fixed point theorems in generalized metric spaces which have many applications in solving integral equations ([14, 15, 27]).

*Corresponding author: hojat.afshari@yahoo.com

2. Preliminaries

Let \mathcal{M} be a nonempty set. A generalized \mathbb{D} -metric on \mathcal{M} is a function, $\mathbb{D} : \mathcal{M}^3 \rightarrow \mathbb{R}^+$ that satisfies the following

(D1) $\mathbb{D}(\zeta, \eta, \zeta) \geq 0$,

(D2) $\mathbb{D}(\zeta, \eta, \zeta) = 0$ if and only if $\zeta = \eta = \zeta$,

(D3) $\mathbb{D}(\zeta, \eta, \zeta) = \mathbb{D}(p\{\zeta, \eta, \zeta\})$, (symmetric) where p is a permutation function,

(D4) $\mathbb{D}(\zeta, \eta, \zeta) \leq \mathbb{D}(\zeta, \alpha, \alpha) + \mathbb{D}(\alpha, \eta, \zeta)$,

where $\zeta, \eta, \zeta, \alpha \in \mathcal{M}$. \mathbb{D} is called a generalized \mathbb{D} -metric and the pair $(\mathcal{M}, \mathbb{D})$ is called a generalized \mathbb{D} -metric space.

Let

$$d_{\mathbb{D}}(\zeta, \eta) = \mathbb{D}(\zeta, \eta, \eta) + \mathbb{D}(\eta, \zeta, \zeta), \quad \forall \zeta, \eta \in \mathcal{M}. \quad (2.1)$$

Obviously that $d_{\mathbb{D}}$ is a metric.

Remark 2.1. In a \mathbb{D} -metric space, we have:

(i) $\mathbb{D}(\zeta, \zeta, \eta) \leq \mathbb{D}(\zeta, \zeta, \zeta) + \mathbb{D}(\zeta, \eta, \eta) = \mathbb{D}(\zeta, \eta, \eta)$,

(ii) $\mathbb{D}(\eta, \eta, \zeta) \leq \mathbb{D}(\eta, \eta, \eta) + \mathbb{D}(\eta, \zeta, \zeta) = \mathbb{D}(\eta, \zeta, \zeta)$,

(iii) $\mathbb{D}(\zeta, \zeta, \eta) = \mathbb{D}(\zeta, \eta, \eta)$.

Definition 2.2. [19] Let $\{\zeta_n\}$ be a sequence of $(\mathcal{M}, \mathbb{D})$. $\{\zeta_n\}$ is \mathbb{D} -convergent to $\zeta \in \mathcal{M}$ if

$$\lim_{n, m \rightarrow +\infty} \mathbb{D}(\zeta, \zeta_n, \zeta_m) = 0.$$

that is, for $\varepsilon > 0$, there exists $N \in \mathbb{N}$ with $\mathbb{D}(\zeta, \zeta_n, \zeta_m) < \varepsilon$, for $n, m \geq N$.

Proposition 2.3. [19] Let $(\mathcal{M}, \mathbb{D})$ be a \mathbb{D} -metric space. The following are equivalent

(i) $\{\zeta_n\}$ is \mathbb{D} -convergent to \mathcal{M} ,

(ii) $\mathbb{D}(\zeta_n, \zeta_n, \zeta) \rightarrow 0$ as $n \rightarrow \infty$,

(iii) $\mathbb{D}(\zeta_n, \zeta, \zeta) \rightarrow 0$ as $n \rightarrow \infty$,

(vi) $\mathbb{D}(\zeta_n, \zeta_m, \zeta) \rightarrow 0$ as $n, m \rightarrow \infty$.

Definition 2.4. [19] Let $(\mathcal{M}, \mathbb{D})$ be a \mathbb{D} -metric space. $\{\zeta_n\}$ is called a \mathbb{D} -Cauchy if for any $\varepsilon > 0$, there exists $N \in \mathbb{N}$ with $\mathbb{D}(\zeta_n, \zeta_m, \zeta_l) < \varepsilon$ for all $m, n, l \geq N$, that is, $\mathbb{D}(\zeta_n, \zeta_m, \zeta_l) \rightarrow 0$ as $n, m, l \rightarrow \infty$.

Definition 2.5. [28] Let $F : \mathbb{R}^+ \rightarrow \mathbb{R}$ satisfying,

(F₁) F is strictly increasing, i.e. for $\alpha, \beta \in \mathbb{R}_+$ with $\alpha < \beta$, $F(\alpha) < F(\beta)$;

(F₂) for $\{\alpha_n\}_{n \in \mathbb{N}}$ of positive real numbers $\lim_{n \rightarrow \infty} \alpha_n = 0$ if and only if $\lim_{n \rightarrow \infty} F(\alpha_n) = -\infty$;

(F₃) $\exists k \in (0, 1)$ with $\lim_{\alpha \rightarrow 0^+} \alpha^k F(\alpha) = 0$.

We say that $T : \mathcal{M} \rightarrow \mathcal{M}$ is F -contraction if there exists $\tau > 0$ such that

$$\forall \zeta, \eta \in \mathcal{M}, \quad d(T\zeta, T\eta) > 0 \Rightarrow \tau + F(d(T\zeta, T\eta)) \leq F(d(\zeta, \eta)).$$

Example 2.6. Let $f_i : \mathcal{R}^+ \rightarrow \mathcal{R}$ ($i = 1, 2, 3, 4, 5, 6$), defined by

$$e_1) f_1(t) = -\frac{1}{\sqrt{\alpha}}, \alpha > 0,$$

$$e_2) f_2(t) = \ln(\alpha), \alpha > 0,$$

$$e_3) f_3(t) = \alpha + \ln(\alpha), \alpha > 0,$$

$$e_4) f_4(t) = \ln(\alpha^2 + \alpha), \alpha > 0,$$

$$e_5) f_5(t) = F(\alpha) = \tan(\alpha + \frac{\pi}{2}),$$

$e_6) f_6(t) = F(\alpha) = -\frac{1}{\alpha^2}, \alpha > 0$. Then f_1, f_2, f_3, f_4, f_5 and f_6 satisfy (F_1) – (F_3) and so are F -contractions.

Definition 2.7. Suppose $T : \mathcal{M} \rightarrow \mathcal{M}$ and $\alpha : \mathcal{M} \times \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{R}^+$. T is α -admissible if

$$\zeta, \eta, \zeta \in \mathcal{M}, \quad \alpha(\zeta, \eta, \zeta) \geq 1 \implies \alpha(T\zeta, T\eta, T\zeta) \geq 1.$$

Denote with Φ the family of nondecreasing functions $\phi : [0, +\infty) \rightarrow [0, +\infty)$ continuous in $t = 0$ such that

(i) $\phi(t) = 0$ if and only if $t = 0$,

(ii) $\phi(t+s) \leq \phi(t) + \phi(s)$,

and $\sum_{n=1}^{+\infty} \phi^n(t) < +\infty$ for $t > 0$, where ϕ^n is the n -th iterate of ϕ .

Lemma 2.8. For $\phi : \mathcal{R}^+ \rightarrow \mathcal{R}^+$ the following holds:

if ϕ is nondecreasing then for $t > 0$, $\lim_{n \rightarrow +\infty} \phi^n(t) = 0$ implies $\phi(t) < t$.

Theorem 2.9. Let $\{\mathfrak{A}_j\}_{j=1}^m$ be a family of nonempty \mathbb{D} -closed subsets of complete space $(\mathcal{M}, \mathbb{D})$. Let $\eta = \bigcup_{j=1}^m \mathfrak{A}_j$ and $T : Y \rightarrow Y$ be a α -admissible satisfying

$$T(\mathfrak{A}_j) \subseteq \mathfrak{A}_{j+1}, \quad j = 1, \dots, m, \quad \text{where} \quad \mathfrak{A}_{m+1} = \mathfrak{A}_1.$$

If there exist $\alpha : Y \times Y \times Y \rightarrow \mathcal{R}^+$ and $\phi \in \Phi$ such that

$$\alpha(\zeta, \eta, T\zeta)\mathbb{D}(T\zeta, T\eta, T\zeta) \leq \phi(\mathbb{D}(\zeta, \eta, \zeta)), \quad (2.2)$$

holds for all $\zeta \in \mathfrak{A}_j$ and $\eta, \zeta \in \mathfrak{A}_{j+1}$, $j = 1, \dots, m$, and there exist $\zeta_0 \in Y$ such that $\alpha(\zeta_0, T\zeta_0, T^2\zeta_0) \geq 1$, then T has a unique fixed point in $\bigcap_{j=1}^m \mathfrak{A}_j$.

Proof. Let $\zeta_0 \in Y$ with $\alpha(\zeta_0, T\zeta_0, T^2\zeta_0) \geq 1$. Suppose $\zeta_0 \in \mathfrak{A}_1$. Define the sequence $\{\zeta_n\}$ in Y as follows

$$\zeta_n = T\zeta_{n-1} \quad \text{for all } n \in \mathbf{N}.$$

Since T is cyclic, $\zeta_0 \in \mathfrak{A}_1$, $\zeta_1 = T(\zeta_0) \in \mathfrak{A}_2, \dots$ and so, if $\zeta_{n_0+1} = \zeta_{n_0}$ for some $n_0 \in \mathbf{N}$, clearly, the fixed point of T is ζ_{n_0} . Let $\zeta_n \neq \zeta_{n+1}$ for $n \in \mathbf{N}$. Since T is α -admissible, we have

$$\alpha(\zeta_0, \zeta_1, \zeta_2) = \alpha(\zeta_0, T\zeta_0, T^2\zeta_0) \geq 1 \implies \alpha(T\zeta_0, T\zeta_1, T\zeta_2) = \alpha(\zeta_1, \zeta_2, \zeta_3) \geq 1.$$

By induction, we get

$$\alpha(\zeta_{n-1}, \zeta_n, \zeta_{n+1}) \geq 1, \quad \text{for all } n \in \mathbf{N}. \quad (2.3)$$

Applying (2.9) with $\zeta = \zeta_{n-1}$ and $\eta = \zeta = \zeta_n$, and utilizing (2.10), we deduce

$$\begin{aligned} 0 \leq \mathbb{D}(\zeta_n, \zeta_{n+1}, \zeta_{n+1}) &= \mathbb{D}(T\zeta_{n-1}, T\zeta_n, T\zeta_n) \\ &\leq \alpha(\zeta_{n-1}, \zeta_n, T\zeta_n)\mathbb{D}(T\zeta_{n-1}, T\zeta_n, T\zeta_n) \leq \phi(\mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n)). \end{aligned}$$

Therefore,

$$\mathbb{D}(\zeta_n, \zeta_{n+1}, \zeta_{n+1}) \leq \phi^n(\mathbb{D}(\zeta_0, \zeta_1, \zeta_1)), \quad \text{for all } n \in \mathbf{N}. \quad (2.4)$$

Fix $\varepsilon > 0$ and let $n(\varepsilon) \in \mathbf{N}$ such that $\sum_{n \geq n(\varepsilon)} \phi^n(\mathbb{D}(\zeta_0, \zeta_1, \zeta_1)) < \varepsilon$.

By (D3) and (D4), we have

$$\mathbb{D}(\zeta, \eta, \eta) = \mathbb{D}(\eta, \eta, \zeta) \leq \mathbb{D}(\eta, \zeta, \zeta) + \mathbb{D}(\zeta, \eta, \zeta) = 2\mathbb{D}(\eta, \zeta, \zeta). \quad (2.5)$$

The inequality (2.5) with $\zeta = \zeta_n$ and $\eta = \zeta_{n-1}$ becomes

$$\mathbb{D}(\zeta_n, \zeta_{n-1}, \zeta_{n-1}) \leq 2\mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n). \quad (2.6)$$

Letting $n \rightarrow \infty$ in (2.6), we get

$$\lim_{n \rightarrow \infty} \mathbb{D}(\zeta_n, \zeta_{n-1}, \zeta_{n-1}) = 0,$$

we show $\{\zeta_n\}$ is Cauchy $(\mathcal{M}, d_{\mathbb{D}})$. Let $n, l \in \mathbf{N}$ with $n > l > n(\varepsilon)$ we obtain

$$\begin{aligned} d_{\mathbb{D}}(\zeta_n, \zeta_l) &\leq d_{\mathbb{D}}(\zeta_n, \zeta_{n-1}) + d_{\mathbb{D}}(\zeta_{n-1}, \zeta_{n-2}) + \dots + d_{\mathbb{D}}(\zeta_{l+1}, \zeta_l) \\ &= \mathbb{D}(\zeta_n, \zeta_{n-1}, \zeta_{n-1}) + \mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n) \\ &\quad + \mathbb{D}(\zeta_{n-1}, \zeta_{n-2}, \zeta_{n-2}) + \mathbb{D}(\zeta_{n-2}, \zeta_{n-1}, \zeta_{n-1}) + \dots \\ &\quad + \mathbb{D}(\zeta_{l+1}, \zeta_l, \zeta_l) + \mathbb{D}(\zeta_l, \zeta_{l+1}, \zeta_{l+1}) \\ &= \sum_{i=l+1}^n [\mathbb{D}(\zeta_i, \zeta_{i-1}, \zeta_{i-1}) + \mathbb{D}(\zeta_{i-1}, \zeta_i, \zeta_i)], \end{aligned} \quad (2.7)$$

By using of (2.12) and (2.6) we obtain

$$\begin{aligned} 0 \leq d_{\mathbb{D}}(\zeta_n, \zeta_l) &\leq \sum_{i=l+1}^n [2\mathbb{D}(\zeta_{i-1}, \zeta_i, \zeta_i) + \mathbb{D}(\zeta_{i-1}, \zeta_i, \zeta_i)] \\ &\leq \sum_{i=l+1}^n 3\phi^{i-1}(\mathbb{D}(\zeta_0, \zeta_1, \zeta_1)) \\ &\leq \sum_{l > n(\varepsilon)} 3\phi^l(\mathbb{D}(\zeta_0, \zeta_1, \zeta_1)) < \varepsilon. \end{aligned}$$

Thus $\{\zeta_n\}$ is Cauchy in $(\mathcal{M}, d_{\mathbb{D}})$. $(\mathcal{M}, \mathbb{D})$ is \mathbb{D} -complete, hence $(\mathcal{M}, d_{\mathbb{D}})$ is complete and then, $\{\zeta_n\}$ converges to $u \in \mathcal{M}$. Furthermore, $\{\zeta_n\}$ is \mathbb{D} -Cauchy in $(\mathcal{M}, \mathbb{D})$. Now we show that $u \in \cap_{j=1}^m \mathfrak{A}_j$. If $\zeta_0 \in \mathfrak{A}_1$, then, $\{\zeta_{m(n-1)}\}_{n=1}^{\infty} \in \mathfrak{A}_1$, $\{\zeta_{m(n-1)+1}\}_{n=1}^{\infty} \in \mathfrak{A}_2$, by continuing, $\{\zeta_{mn-1}\}_{n=1}^{\infty} \in \mathfrak{A}_m$. All the m subsequences are \mathbb{D} -convergent so converge to the same limit u . Moreover, the sets \mathfrak{A}_j are \mathbb{D} -closed, thus the limit $u \in \cap_{j=1}^m \mathfrak{A}_j$. In fact $u \in \mathcal{M}$ is a fixed point of T , considering (2.9) and setting $\zeta = \zeta_n$, $\eta = \zeta = Tu$ with assuming that $u \neq Tu$ or $d_{\mathbb{D}}(u, Tu) > 0$, we get,

$$\begin{aligned} 0 \leq d_{\mathbb{D}}(\zeta_n, Tu) &= \mathbb{D}(\zeta_n, Tu, Tu) + \mathbb{D}(Tu, \zeta_n, \zeta_n) \\ &= \mathbb{D}(T\zeta_{n-1}, Tu, Tu) + \mathbb{D}(Tu, T\zeta_{n-1}, T\zeta_{n-1}) \\ &\leq \mathbb{D}(T\zeta_{n-1}, Tu, Tu) + 2\mathbb{D}(T\zeta_{n-1}, Tu, Tu) \\ &\leq 3\alpha(\zeta_{n-1}, u, u)\mathbb{D}(T\zeta_{n-1}, Tu, Tu) \\ &\leq 3\phi(\mathbb{D}(\zeta_{n-1}, u, u)). \end{aligned} \quad (2.8)$$

Tending $n \rightarrow \infty$, we end up with $0 \leq d_{\mathbb{D}}(u, Tu) \leq 0$ which contradicts the assumption $d_{\mathbb{D}}(u, Tu) > 0$, hence $u = Tu$ and then $u \in \mathcal{M}$ is a fixed point of T . To prove the uniqueness, suppose $v \in \mathcal{M}$ is another fixed point of T such that $v \neq u$. Both u and v lie in $\cap_{j=1}^m \mathcal{A}_j$, thus we can substitute $\zeta = u$ and $\eta = \zeta = v$ in (2.9). So

$$\mathbb{D}(Tu, Tv, Tv) \leq \alpha(u, v, v)\mathbb{D}(Tu, Tv, Tv) \leq \phi(\mathbb{D}(u, v, v)).$$

From lemma 2.8 and $v = Tv$ we have

$$\mathbb{D}(u, v, v) \leq \alpha(u, v, v)\mathbb{D}(Tu, Tv, Tv) < \mathbb{D}(u, v, v),$$

which is contradiction, Thus $u = v$, and the fixed point of T is unique. \square

Theorem 2.10. Let $(\mathcal{M}, \mathbb{D})$ be a \mathbb{D} -complete \mathbb{D} -metric space and $\{\mathcal{A}_j\}_{j=1}^m$ be a family of nonempty \mathbb{D} -closed subsets of \mathcal{M} . Let $\eta = \cup_{j=1}^m \mathcal{A}_j$ and $T : Y \rightarrow Y$ be a α -admissible satisfying

$$T(\mathcal{A}_j) \subseteq \mathcal{A}_{j+1}, \quad j = 1, \dots, m, \quad \text{where} \quad \mathcal{A}_{m+1} = \mathcal{A}_1.$$

If there exist two functions $\alpha : Y \times Y \times Y \rightarrow \mathfrak{R}^+$ and $\phi \in \Phi$ such that

$$\forall \zeta, \eta \in \mathcal{M}, \quad (\mathbb{D}(T\zeta, T\eta, T\zeta) > 0 \Rightarrow \tau + \alpha(\zeta, \eta, T\zeta)F(\mathbb{D}(T\zeta, T\eta, T\zeta)) \leq F(\phi(\mathbb{D}(\zeta, \eta, \zeta)))) \quad (2.9)$$

holds for all $\zeta \in \mathcal{A}_j$ and $\eta, \zeta \in \mathcal{A}_{j+1}$, $j = 1, \dots, m$, and there exists $\zeta_0 \in Y$ such that $\alpha(\zeta_0, T\zeta_0, T^2\zeta_0) \geq 1$, then T has a unique fixed point in $\cap_{j=1}^m \mathcal{A}_j$.

Proof. Suppose $\zeta_0 \in Y$ with $\alpha(\zeta_0, T\zeta_0, T^2\zeta_0) \geq 1$, assume that $\zeta_0 \in \mathcal{A}_1$. Define

$$\zeta_n = T\zeta_{n-1} \quad \text{for all } n \in \mathbf{N}.$$

Since T is cyclic, $\zeta_0 \in \mathcal{A}_1$, $\zeta_1 = T(\zeta_0) \in \mathcal{A}_2, \dots$ and so on. If $\zeta_{n_0+1} = \zeta_{n_0}$ for some $n_0 \in \mathbf{N}$, then, the fixed point of T is ζ_{n_0} . If $\zeta_n \neq \zeta_{n+1}$ for all $n \in \mathbf{N}$, since T is α -admissible, we have

$$\alpha(\zeta_0, \zeta_1, \zeta_2) = \alpha(\zeta_0, T\zeta_0, T^2\zeta_0) \geq 1 \Rightarrow \alpha(T\zeta_0, T\zeta_1, T\zeta_2) = \alpha(\zeta_1, \zeta_2, \zeta_3) \geq 1.$$

By induction, we get

$$\alpha(\zeta_{n-1}, \zeta_n, \zeta_{n+1}) \geq 1, \quad \text{for all } n \in \mathbf{N}. \quad (2.10)$$

Applying the inequality (2.9) with $\zeta = \zeta_{n-1}$ and $\eta = \zeta = \zeta_n$, and using (2.10), we obtain

$$\begin{aligned} 0 \leq F(\mathbb{D}(\zeta_n, \zeta_{n+1}, \zeta_{n+1})) &= F(\mathbb{D}(T\zeta_{n-1}, T\zeta_n, T\zeta_n)) \\ &\leq \alpha(\zeta_{n-1}, \zeta_n, T\zeta_n)F(\mathbb{D}(T\zeta_{n-1}, T\zeta_n, T\zeta_n)) \\ &\leq F(\phi(\mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n))) - \tau \\ &< F(\mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n)) - \tau. \end{aligned} \quad (2.11)$$

So,

$$F(\mathbb{D}(\zeta_n, \zeta_{n+1}, \zeta_{n+1})) \leq F(\mathbb{D}(\zeta_0, \zeta_1, \zeta_1)) - n\tau, \quad \text{for all } n \in \mathbf{N}. \quad (2.12)$$

tending $n \rightarrow \infty$ we have

$$\lim_{n \rightarrow \infty} F(\mathbb{D}(\zeta_n, \zeta_{n+1}, \zeta_{n+1})) = -\infty.$$

Thus,

$$\lim_{n \rightarrow \infty} \mathbb{D}(\zeta_n, \zeta_{n+1}, \zeta_{n+1}) = 0.$$

Hence

$$\lim_{n \rightarrow \infty} \mathbb{D}(\zeta_n, \zeta_{n-1}, \zeta_{n-1}) = 0.$$

Also if put $\gamma_n = \mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n)$, then by using (2.11) we obtain,

$$(\gamma_n)^k F(\gamma_n) \leq (\gamma_n)^k F(\gamma_0) - (\gamma_n)^k n \tau. \quad (2.13)$$

Thus

$$(\gamma_n)^k F(\gamma_n) - (\gamma_n)^k F(\gamma_0) \leq (\gamma_n)^k (F(\gamma_0) - n \tau) - (\gamma_n)^k F(\gamma_0) = -(\gamma_n)^k n \tau \leq 0$$

By attention to, $\lim_{n \rightarrow \infty} \gamma_n^k F(\gamma_n) = 0$ and by $\lim_{n \rightarrow \infty} \gamma_n = 0$ and Letting $n \rightarrow \infty$ in (2.13), we get

$$\lim_{n \rightarrow \infty} (\gamma_n)^k n = 0. \quad (2.14)$$

From (2.14) there exists $n_1 \in \mathbb{N}$ with $(\gamma_n)^k n \leq 1$ for all $n \geq n_1$, hen we have

$$\gamma_n \leq \frac{1}{n^{\frac{1}{k}}}, \quad \forall n \geq n_1.$$

We show $\{\zeta_n\}$ is cauchy $(\mathcal{M}, d_{\mathbb{D}})$ where $d_{\mathbb{D}}$ is given in (2.1). Let $n, l \in \mathbb{N}$ with $n > l > n_1$ we obtain

$$\begin{aligned} d_{\mathbb{D}}(\zeta_n, \zeta_l) &\leq d_{\mathbb{D}}(\zeta_n, \zeta_{n-1}) + d_{\mathbb{D}}(\zeta_{n-1}, \zeta_{n-2}) + \dots + d_{\mathbb{D}}(\zeta_{l+1}, \zeta_l) \\ &= \mathbb{D}(\zeta_n, \zeta_{n-1}, \zeta_{n-1}) + \mathbb{D}(\zeta_{n-1}, \zeta_n, \zeta_n) \\ &\quad + \mathbb{D}(\zeta_{n-1}, \zeta_{n-2}, \zeta_{n-2}) + \mathbb{D}(\zeta_{n-2}, \zeta_{n-1}, \zeta_{n-1}) + \dots \\ &\quad + \mathbb{D}(\zeta_{l+1}, \zeta_l, \zeta_l) + \mathbb{D}(\zeta_l, \zeta_{l+1}, \zeta_{l+1}) \\ &= \sum_{i=l+1}^n [\mathbb{D}(\zeta_i, \zeta_{i-1}, \zeta_{i-1}) + \mathbb{D}(\zeta_{i-1}, \zeta_i, \zeta_i)]. \end{aligned} \quad (2.15)$$

Tending $n, l \rightarrow \infty$ we obtain that $\{\zeta_n\}$ is Cauchy in the $(\mathcal{M}, d_{\mathbb{D}})$. $(\mathcal{M}, \mathbb{D})$ is \mathbb{D} -complete, hence, $\{\zeta_n\}$ converges $u \in \mathcal{M}$. Furthermore, $\{\zeta_n\}$ is \mathbb{D} -Cauchy in $(\mathcal{M}, \mathbb{D})$. Now $u \in \bigcap_{j=1}^m \mathfrak{A}_j$. if $\zeta_0 \in \mathfrak{A}_1$, then $\{\zeta_{m(n-1)}\}_{n=1}^{\infty} \in \mathfrak{A}_1$, $\{\zeta_{m(n-1)+1}\}_{n=1}^{\infty} \in \mathfrak{A}_2$, by continuing, $\{\zeta_{mn-1}\}_{n=1}^{\infty} \in \mathfrak{A}_m$. All the m subsequences are \mathbb{D} -convergent and hence, they all converge to the same limit u . In addition, the sets \mathfrak{A}_j are \mathbb{D} -closed, thus the limit $u \in \bigcap_{j=1}^m \mathfrak{A}_j$. $u \in \mathcal{M}$ is a

fixed point of T , because by (2.1) and (2.9) with $\zeta = \zeta_n$, $\eta = \zeta = Tu$ if $u \neq Tu$ or $d_{\mathbb{D}}(u, Tu) > 0$, then,

$$\begin{aligned} 0 \leq d_{\mathbb{D}}(\zeta_n, Tu) &= \mathbb{D}(\zeta_n, Tu, Tu) + \mathbb{D}(Tu, \zeta_n, \zeta_n) \\ &= \mathbb{D}(T\zeta_{n-1}, Tu, Tu) + \mathbb{D}(Tu, T\zeta_{n-1}, T\zeta_{n-1}) \\ &\leq \mathbb{D}(T\zeta_{n-1}, Tu, Tu) + 2\mathbb{D}(T\zeta_{n-1}, Tu, Tu) \\ &= 3\mathbb{D}(T\zeta_{n-1}, Tu, Tu) \\ &= 3(\mathbb{D}(\zeta_n, u, u)). \end{aligned} \quad (2.16)$$

Tending $n \rightarrow \infty$, $0 \leq d_{\mathbb{D}}(u, Tu) \leq 0$ which contradicts the assumption $d_{\mathbb{D}}(u, Tu) > 0$, hence $u = Tu$, so $u \in \mathcal{M}$ is a fixed point of T . Suppose $v \in \mathcal{M}$ is another fixed point of T such that $v \neq u$. Both u and v lie in $\cap_{j=1}^m \mathbb{D}j$, thus we can substitute $\zeta = u$ and $\eta = \zeta = v$ in (2.9). This yields

$$F(\mathbb{D}(Tu, Tv, Tv)) + \tau \leq \alpha(u, v, v)F(\mathbb{D}(Tu, Tv, Tv)) + \tau \leq F(\phi(\mathbb{D}(u, v, v))),$$

therefore

$$F(\mathbb{D}(Tu, Tv, Tv)) \leq F(\phi(\mathbb{D}(u, v, v))).$$

F is strictly increasing, by 2.8 we obtain

$$\mathbb{D}(Tu, Tv, Tv) \leq \phi(\mathbb{D}(u, v, v)) < \mathbb{D}(u, v, v),$$

this is a contradiction, thus $u = v$, and the fixed point of T is unique. \square

3. Conclusion

In the current study, we used the F -contraction mapping introduced by Wardowski to obtain fixed point results by method of Samet in generalized complete metric spaces. Over the last decade authors proved some fixed point results for F -contraction mappings in metric spaces. We showed that this results hold in \mathbb{D} -metric spaces under some conditions.

References

- [1] Abbas M, Berzig M, Nazir T, Karapinar T (2016). *Iterative Approximation of Fixed Points for Presic Type F-Contraction Operators*. U.P.B. Sci. Bull., Series A. **78**(2): 147-160.
- [2] Abdullah NI, Shaakir LK (2020). *Generalize partial Metric spaces*. Ibn AL-Haitham J. Pure Appl. Sci. **34**(1): 47-59. <https://doi.org/10.30526/34.1.2566>.
- [3] Afshari H, Atapour M, Karapinar E (2020). *A discussion on a generalized Geraghty multi-valued mappings and applications*. Advances in Difference Equations. **2020**(1): 1-14. <https://doi.org/10.1186/s13662-020-02819-2>.
- [4] Afshari H, Aydi H, Karapinar, E (2020). *On generalized α - ϕ -Geraghty contractions on b -metric spaces*. Georgian Mathematical Journal. **27**(1): 9-21. <https://doi.org/10.1515/gmj-2017-0063>.
- [5] Afshari H, Aydi H, Karapinar E (2016). *Existence of fixed points of set-valued mappings in b -metric spaces*. East Asian mathematical journal. **32**(3): 319-32. <https://doi.org/10.7858/eamj.2016.024>.
- [6] Afshari H, Kalantari S, Aydi H (2018). *Fixed point results for generalized α - ψ -Suzuki-contractions in quasi- b -metric-like spaces*. Asian-European journal of mathematics. **11**(01): 1850012.
- [7] Afshari H (2019). *Solution of fractional differential equations in quasi- b -metric and b -metric-like spaces*. Advances in Difference Equations. **2019**(1): 1-14. <https://doi.org/10.1186/s13662-019-2227-9>.

- [8] Agarwal RP, Aksoy Ü, Karapınar E, Erhan IM (2020). *F-contraction mappings on metric-like spaces in connection with integral equations on time scales*. Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas. **114** :1-12. <https://doi.org/10.1007/s13398-020-00877-5>.
- [9] Alsulami HH, Gülyaz S, Karapınar E, Erhan IM (2016). *An Ulam stability result on quasi-b-metric-like spaces*. Open Mathematics. **14**(1): 1087-103. <https://doi.org/10.1515/math-2016-0097>.
- [10] Aydi H, Karapınar E, Yazidi H (2017). *Modified F-Contractions via α -Admissible Mappings and Application to Integral Equations*. Filomat. **31**(5): 1141-1148. <https://doi.org/10.102298/FIL1705141A>.
- [11] Dhage BC (2000). *Generalized metric spaces and topological structure I*. Analele scientific ale universities AI. I. Cuzadin Lasi, Sieve Nona. Matematca. **47**: 3-24.
- [12] Hazarika B, Karapınar E, Arab R, Rabbani M (2018). *Metric-like spaces to prove existence of solution for nonlinear quadratic integral equation and numerical method to solve it*. Journal of Computational and Applied Mathematics. **328**(2018): 302-313. <https://doi.org/10.1016/j.cam.2017.07.012>.
- [13] Hussain N, Karapınar E, Salimi P, Vetro P (2013). *Fixed point results for G m-Meir-Keeler contractive and G - α - ψ -Meir-Keeler contractive mappings*. Fixed Point Theory and Applications. **2013**(1): 1-4. <https://doi.org/10.1186/1687-1812-2013-34>.
- [14] Iqbal M, Shah K, Khan RA (2021). *On using coupled fixed-point theorems for mild solutions to coupled system of multipoint boundary value problems of nonlinear fractional hybrid pantograph differential equations*. Mathematical Methods in the Applied Sciences. **44**(10): 8113-8124. <https://doi.org/10.1002/mma.5799>.
- [15] Khalehghli S, Rahimi H, Gordji ME (2020). *Fixed point theorems in R -metric spaces with applications*. AIMS Mathematics. **5**(4): 3125-3137. <https://doi.org/10.3934/math.2020201>.
- [16] Karapınar E (2012). *Edelstein type fixed point theorems*. Fixed Point Theory and Applications. **2012**(1): 1-12. <https://doi.org/10.1186/1687-1812-2012-107>.
- [17] Karapınar E, Fulga A, Agarwal RP (2020). *A survey: F-contractions with related fixed point results*. Journal of Fixed Point Theory and Applications. **22**(3): 1-58. <https://doi.org/10.1007/s11784-020-00803-7>.
- [18] Karapınar E (2010). *Couple fixed point theorems for nonlinear contractions in cone metric spaces*. Computers and Mathematics with Applications. **59**(12): 3656-3668.
- [19] Mustafa Z, Sims B (2006). *A new approach to generalized metric spaces*. Journal of Nonlinear and convex Analysis. **7**(2): 289.
- [20] Qawaqneh H, Noorani MS, Shatanawi W, Aydi H, Alsamir H (2019). *Fixed point results for multi-valued contractions in b -metric spaces and an application*. Mathematics. **7**(2): 132.
- [21] Rhoades BE (1996). *A fixed point theorem for generalized metric spaces*. International Journal of Mathematics and Mathematical Sciences. **19**(3): 457-460. <https://doi.org/10.1155/S0161171296000658>
- [22] Samet B (2014). *Fixed points for α - ψ contractive mappings with an application to quadratic integral equations*. Electron. J. Differ. Equ. **2014**152: 1-18.
- [23] Samet B, Vetro C, Vetro P (2012). *Fixed point theorems for α - ψ -contractive type mappings*. Nonlinear analysis: theory, methods and applications. **75**(4): 2154-2165. <https://doi.org/10.1016/j.na.2011.10.014>
- [24] Shoaib M, Sarwar M, Shah K, Kumam P. (2016). *Fixed point results and its applications to the systems of non-linear integral and differential equations of arbitrary order*. J. Nonlinear Sci. Appl. **96**: 4949-4962.
- [25] Shahraki M, Sedghi S, Aleomraninejad SM, Mitrovic ZD (2020). *Some fixed point results on S -metric spaces*. Acta Univ. Sapientiae, Mathematica. **12**(2): 347-357. <https://doi.org/10.2478/ausm-2020-0024>.
- [26] Sedghi S, Shobe N, Aliouche A (2012). *A generalization of fixed point theorems in S -metric spaces*. Matematički vesnik. **64**(3): 258-266.
- [27] Silin DB (1997). *On set-valued differentiation and integration*. Set-Valued Analysis. **2**(2): 107-146.
- [28] Wardowski D (2012). *Fixed points of a new type of contractive mappings in complete metric spaces*. Fixed point theory and applications. **2012**(1): 1-6. <https://doi.org/10.1186/1687-1812-2012-94>.
- [29] Wardowski D, Van Dung N (2014). *Fixed points of F -weak contractions on complete metric spaces*. Demonstratio Mathematica. **47**(1): 146-155. <https://doi.org/10.2478/dema-2014-0012>.