

Uniqueness and Stability Results on Non-local Stochastic Random Impulsive Integro-Differential Equations

S. VARSHINI ^{a,*}, K. BANUPRIYA ^a, K. RAMKUMAR^a, K. RAVIKUMAR ^a

^a PSG College of Arts and Science, Coimbatore, India.

• Received: 13 September 2021 • Accepted: 11 October 2021 • Published Online: 25 November 2021

Abstract

The paper is concerned with stochastic random impulsive integro-differential equations with non-local conditions. The sufficient conditions guarantees uniqueness of mild solution derived using Banach fixed point theorem. Stability of the solution is derived by incorporating Banach fixed point theorem with certain inequality techniques.

Keywords: Uniqueness, Stability, Random impulse, Stochastic Integro-differential Equations.

2010 MSC: 34A12, 34K20, 58C30.

1. Introduction

Mathematical modelling in Engineering and Scientific fields results in integral, ordinary or partial differential equations, stochastic differential equations or integro-differential equations. Specifically, in the fluid dynamics, chemical kinetics and biological disciplines, systems in general are of integro-differential type refer [1, 2, 3, 4]. Evolution processes from fields of population dynamics, aeronautics, economics, telecommunications and physics experiences drastic change of state at certain moments of time between the intervals of continuous evolution. Comparatively, the duration of these changes are negligible to the total duration that acts instantaneously in the form of impulse. The theory of impulsive differential equations represents a more natural framework for mathematical modelling see [5, 6, 7] and the references therein.

Impulses exist at fixed time or random time. There are many articles featuring the qualitative properties of fixed impulsive type equations [8, 9]. Wu and Meng [10] initiated the study of random impulsive ordinary differential equations and investigated boundedness of solution by Liapunov's direct function. By regular fluctuations in the deterministic models due to noise which appears to be random, the researchers moved to the stochastic

*Corresponding author: svarshini0808@gmail.com

differential equations. There are several articles regarding the study the Stochastic Differential Equations (SDE) [11, 12, 13, 14, 15, 16, 17, 18]. Stochastic effects with the impulsive effect exist in evolution processes in the real world phenomena refer [19, 20]. Sakthivel and Luo [21] investigated the existence and asymptotical stability for mild solution of Impulsive Stochastic Differential Equations.

Pan and cao [22] solved the exponential stability of impulsive stochastic partial differential equations with delays. Cui and yan [23] investigated the existence results for fractional neutral stochastic integro-differential equations with infinite delay. Mao [24] established stability results of stochastic integro-differential equations. Li et.al [25] investigated the existence and Hyers-Ulam stability of random impulsive stochastic functional differential equations with finite delay.

For instance, let us consider the classical stock price model [26]

$$\begin{aligned} d(N(t)) &= uN(t)dt + \sigma N(t)dB_t, & t \geq 0, t \neq \delta_k, \\ N(\delta_k) &= \alpha_k N(\delta_k^-), & k = 1, 2, 3, \dots, \\ N(0) &= N_0. \end{aligned} \quad (1.1)$$

The system 1.1 represents impulsive stochastic differential equations. Here B_t is a Brownian motion or Wiener process. $N(t)$ represents the price of stock at time t and δ_k represents the release time of an information relating to the stock $N(\delta_k^-) = \lim_{t \rightarrow \delta_k^-} N(t)$ and $N_0 \in \mathbb{R}$. In reality, $\{\delta_k\}$ represents a sequence of random variable which satisfies $0 < \delta_1 < \delta_2 < \dots$

El. Borai [27] studied the existence and uniqueness of stochastic fractional integro-differential equations. Ahmed and El. Borai [28] established the existence results of mild solutions of Hilfer fractional stochastic integro-differential equations with non-local conditions. Guo et.al [29] analysed the exponential stability of impulsive stochastic functional differential equations. Liu and Xu [30] studied the averaging results for impulsive fractional neutral stochastic differential equations.

Sayooj et.al [31] considered a non-local random impulsive integro-differential system and calculated the existence, uniqueness and stability results. There are several papers that includes the study of impulsive integro-differential equations involving random impulses [32, 33]. Therefore, the study of random impulsive stochastic differential equations has a room for improvement. Thus the main objective of this work is to present non-local random impulsive stochastic differential equations hoping that the results obtained would contribute to the area.

Let us consider a the non-local stochastic random impulsive integro-differential equation of the form

$$\begin{cases} dx(t) = f(t, x_t)dt + \int_0^t g(\eta, x(t+\eta))d\omega(\eta), & t \neq \xi_k, t \geq r, \\ x(\xi_k) = b_k(\delta_k)x(\xi_k^-), & k = 1, 2, 3, \dots, \\ x_{t_0} + h(x) = x_0. \end{cases} \quad (1.2)$$

where δ_k is a random variable defined from Ω to $\mathcal{D}_k = {}^{\text{def}} (0, d_k)$ for $k = 1, 2, \dots$ where $0 < d_k < +\infty$. Moreover, assume that for $k = 1, 2, \dots$ δ_k follows Erlang distribution, let δ_i and δ_j are independent with each other as $i \neq j$ for $i, j = 1, 2, \dots$. Here we suppose, $T \in (t_0, +\infty)$, $f : [t_0, T] \times \mathcal{C} \rightarrow \mathbb{R}^d$, $g : [t_0, T] \times \mathcal{C} \rightarrow \mathbb{R}^{d \times m}$, $h : [t_0, T] \times \mathcal{C} \rightarrow \mathbb{R}^d$ and

$b_k : \mathcal{D}_k \rightarrow \mathbb{R}^{d \times d}$ and x_t is \mathbb{R}^d -valued stochastic process such that $x(t) \in \mathbb{R}^d, x(t) = \{x(t + \theta) : -\delta \leq \theta \leq 0\}$. The impulsive moments ξ_k form a strictly increasing sequence, i.e., $\xi_0 < \xi_1 < \xi_2 < \dots < \xi_k < \dots < \lim_{k \rightarrow \infty}$, and $x(\xi_k^-) = \lim_{t \rightarrow \xi_k^-} x(t)$. We assume that $\xi_0 = t_0$ and $\xi_k = \xi_{k-1} + \delta_k$ for $k = 1, 2, 3, \dots$. Obviously, ξ_k is a process with increment increments. We assume that $\{N(t), t \geq 0\}$ is a simple counting process generated by $\{\xi_k\}$ and $\{\omega(t) : t \geq 0\}$ is a given Weiner process. We denote \mathfrak{F}^1 the σ -algebra generated by $\{N(t), t \geq 0\}$, and denote \mathfrak{F}^2 the σ -algebra generated by $\{\omega(s), s \geq t\}$. We assume that $\mathfrak{F}_\infty^1, \mathfrak{F}_\infty^2$ and ξ_k are mutually independent.

The manuscript is summarized as follows. Section 2 presents certain preliminaries. Section 3 is devoted to the existence and uniqueness solution of random impulsive stochastic integro-differential equations with non-local condition. In Section 4, with the Lipschitz condition, the stability results are derived.

2. Preliminaries

Let $(\Omega, \mathfrak{F}, \mathbb{P})$ is a probability space with Expectation \mathbb{E} and associate with the normal filtration $\{\mathfrak{F}_t\}$, $t \geq 0$ satisfying $\mathfrak{F}_t = \mathfrak{F}_t^{(1)} \vee \mathfrak{F}_t^{(2)}$. Let $\mathcal{L}^2(\Omega, \mathbb{R}^d)$ be the collection of all strongly measurable, \mathfrak{F}_t measurable, \mathbb{R}^d -valued random variables x with norm $\|x\|_{\mathcal{L}^2} = \left(\mathbb{E} \|x\|^2\right)^{1/2}$, where $\mathbb{E} = \int_{\Omega} x d\mathbb{P}$. Let $\delta > 0$ denote the Banach space of all piecewise continuous \mathbb{R}^d -valued stochastic process $\{\xi(t), t \in [-\delta, 0]\}$ by $\mathfrak{C}([-\delta, 0], \mathcal{L}^2(\Omega, \mathbb{R}^d))$ equipped with the norm,

$$\|\beta\|_{\mathfrak{C}} = \sup_{\theta \in [-\delta, 0]} \left(\mathbb{E} \|\beta(\theta)\|^2\right)^{1/2}.$$

The initial data, [25]

$$x_{t_0} = x_0 = \{\xi(\theta) : -\delta \leq \theta \leq 0\}, \quad (2.1)$$

is an \mathfrak{F}_{t_0} measurable, $[-\delta, 0]$ to \mathbb{R}^d -valued random variable such that $\mathbb{E} \|\xi\|^2 < \infty$.

Definition 2.1. A \mathbb{R}^d valued Stochastic process $x(t)$ on $t - \delta \leq t \leq T$ is called a mild solution to 2.1, if

- (i) For every $t_0 \leq t \leq T$, $x(t_0) = x_0, \{x_t\}_{t_0 \leq t \leq T}$ is \mathfrak{F}_t -adapted;
- (ii)

$$\begin{aligned}
 x(t) = & \sum_{k=0}^{+\infty} \left[\prod_{i=1}^k b_i(\delta_i)(x_0 - h(x)) + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} f(s, x(s)) ds \right. \\
 & + \int_{\xi_k}^t f(s, x(s)) ds + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} \int_0^T g(\eta, x(s + \eta)) d\omega(\eta) ds \\
 & \left. + \int_{\xi_k}^t \int_0^T g(\eta, x(s + \eta)) d\omega(\eta) ds \right] I_{[\xi_k, \xi_{k+1})}(t), \quad t \in [\delta, T].
 \end{aligned} \quad (2.2)$$

where $\prod_{j=p}^q (.) = 1$ as $p > q$,

$$\prod_{j=i}^k b_j(\delta_j) = b_k(\delta_k) b_{k-1}(\delta_{k-1}) \dots b_i(\delta_i),$$

and $I_{(A)}(.)$ is the index function, i.e.,

$$I_A(t) = \begin{cases} 1, & \text{if } t \in A, \\ 0, & \text{if } t \notin A. \end{cases}$$

Lemma 2.2. For any $r \geq 1$ and for arbitrary \mathcal{L}_2^0 -valued predictable process $\Phi(.)$,

$$\sup_{s \in [0, t]} \mathbb{E} \left\| \int_0^s \Phi(u) dW(u) \right\|_{\mathbb{X}}^{2r} \leq (r(2r-1))^r \left(\int_0^t (\mathbb{E} \|\Phi(s)\|_{\mathcal{L}_2^0}^{2r}) ds \right)^r.$$

The following hypotheses are considered to prove our results.

(H1) The function $f : [t_0, T] \times \mathfrak{C} \rightarrow \mathbb{R}^d$ satisfies the Lipschitz condition. For $x, y \in \mathbb{X}$ and $\delta \leq t \leq T$ there exist $\mathcal{L}_0, \mathcal{M}_0 \geq 0$ such that

$$\begin{aligned} \mathbb{E} \|f(t, x) - f(t, y)\|^2 &\leq \mathcal{L}_0 \mathbb{E} \|x - y\|^2 \\ \mathbb{E} \|f(t, 0)\|^2 &\leq \mathcal{M}_0. \end{aligned}$$

(H2) The function $g : [t_0, T] \times \mathfrak{C} \rightarrow \mathbb{R}^{d \times m}$ satisfies the Lipschitz condition. For $x, y \in \mathbb{X}$ and $\delta \leq t \leq T$ there exist $\mathcal{L}_1, \mathcal{M}_1 \geq 0$ such that

$$\begin{aligned} \mathbb{E} \left\| \int_0^T [g(\eta, x(t+\eta)) - g(\eta, y(t+\eta))] d\eta \right\|^2 &\leq \mathcal{L}_1 \mathbb{E} \|x(t+\eta) - y(t+\eta)\|^2, \\ \mathbb{E} \|g(\eta, 0)\|^2 &\leq \mathcal{M}_1. \end{aligned}$$

(H3) The condition $\max_{i,k} \left\{ \prod_{j=1}^k \|b_j(\delta_j)\| \right\}$ is uniformly bounded $\exists B > 0$ such that

$$\max_{i,k} \left\{ \prod_{j=1}^k \|b_j(\delta_j)\| \right\} \leq B,$$

for all $\delta_j \in D_j$, $j = 1, 2, 3, \dots$

(H4) The function $h : [t_0, T] \times \mathfrak{C} \rightarrow \mathbb{R}^d$ satisfies the Lipschitz condition. For $x, y \in \mathbb{X}$ and $\delta \leq t \leq T$ there exist $\mathcal{L}_* \geq 0$ such that

$$\mathbb{E} \|h(x) - h(y)\|^2 \leq \mathcal{L}_* \|x - y\|^2.$$

(H5)

$$\Gamma = B^2 \max\{1, B^2\} (T - \delta)^2 \left[\mathcal{L}_0 + \mathcal{L}_1 + \frac{\mathcal{L}_*}{(T - \delta)^2} \right] < 1.$$

3. Uniqueness

This section is devoted to the study of uniqueness of mild solution of the system 1.2

Theorem 3.1. *Assume that the hypotheses (H1)-(H5) holds. Then the system 1.2 has a unique mild solution in \mathbb{B} .*

Proof. Let T be an arbitrary number $T \leq +\infty$. Initially from definition 2.1, we define a non-linear operator $\phi : \mathbb{B} \rightarrow \mathbb{B}$. Note that the problem 2.1 has a solution if and only if the operator ϕ has a fixed point.

$$\begin{aligned} \phi x(t) = & \sum_{k=0}^{+\infty} \left[\prod_{i=1}^k b_i(\delta_i)(x_0 - h(x)) + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} f(s, x(s)) ds \right. \\ & + \int_{\xi_k}^t f(s, x(s)) ds + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) ds \\ & \left. + \int_{\xi_k}^t \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) ds \right] \times I_{[\xi_k, \xi_{k+1})}(t), \end{aligned}$$

where $t \in [\delta, T]$. We need to show that \mathbb{B} maps \mathbb{B} under ϕ .
 $\|\phi x(t)\|^2$

$$\begin{aligned} \leq & \sum_{k=0}^{+\infty} \left[\left\| \prod_{i=1}^k b_i(\delta_i) \right\| \|(x_0 - h(x))\| + \sum_{i=1}^k \left\| \prod_{j=i}^k b_j(\delta_j) \right\| \int_{\xi_{i-1}}^{\xi_i} \|f(s, x(s))\| ds \right. \\ & + \int_{\xi_k}^t \|f(s, x(s))\| ds + \sum_{i=1}^k \left\| \prod_{j=i}^k b_j(\delta_j) \right\| \int_{\xi_{i-1}}^{\xi_i} \left\| \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) \right\| ds \\ & \left. + \int_{\xi_k}^t \left\| \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) \right\| ds \right] I_{[\xi_k, \xi_{k+1})}(t), \quad t \in [\delta, T] \\ \leq & 3 \left[\sum_{k=0}^{+\infty} \left[\left\| \prod_{i=1}^k b_i(\delta_i) \right\|^2 \|(x_0 - h(x))\|^2 I_{[\xi_k, \xi_{k+1})}(t) \right] + \left[\sum_{k=0}^{+\infty} \left[\sum_{i=1}^k \left\| \prod_{j=i}^k b_j(\delta_j) \right\| \right. \right. \right. \\ & \times \left. \left. \int_{\xi_{i-1}}^{\xi_i} \|f(s, x(s))\| ds + \int_{\xi_k}^t \|f(s, x(s))\| ds \right] I_{[\xi_k, \xi_{k+1})}(t) \right]^2 \\ & + \left[\sum_{k=0}^{+\infty} \left[\sum_{i=1}^k \left\| \prod_{j=i}^k b_j(\delta_j) \right\| \right] \int_{\xi_{i-1}}^{\xi_i} \left\| \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) \right\| \right. \\ & \left. + \int_{\xi_k}^t \left\| \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) \right\| ds \right] I_{[\xi_k, \xi_{k+1})}(t) \right]^2 \end{aligned}$$

$$\begin{aligned}
 &\leq 3 \max_k \left\{ \left\| \prod_{i=1}^k b_i(\delta_i) \right\|^2 \right\} \|x_0 - h(x)\|^2 + 3 \left[\max_{i,k} \left\{ 1, \left\| \prod_{j=i}^k b_j(\delta_j) \right\|^2 \right\} \right]^2 \\
 &\times \left\{ \int_{t_0}^t \|f(s, x(s))\| ds I_{[\xi_k, \xi_{k+1})}(t) \right\}^2 + 3 \left[\max_{i,k} \left\{ 1, \left\| \prod_{j=i}^k b_j(\delta_j) \right\|^2 \right\} \right]^2 \\
 &\times \left\{ \int_{t_0}^t \left\| \int_0^T g(\eta, x(s+\eta)) d\omega(\eta) \right\| ds I_{[\xi_k, \xi_{k+1})}(t) \right\}^2 \\
 &\leq 3B^2 \|x_0 - h(x)\|^2 + 3 \max\{1, B^2\} (t - t_0) \int_{t_0}^t \|f(s, x(s))\|^2 ds \\
 &+ 3 \max\{1, B^2\} (t - t_0) \mathcal{C}_2 \int_{t_0}^t \left[\left\| \int_0^T g(\eta, x(s+\eta)) d\eta \right\| ds \right]^2.
 \end{aligned}$$

Thus we would obtain,

$$\begin{aligned}
 \mathbb{E} \|\phi x(t)\|^2 &\leq 3B^2 \|x_0 - h(x)\|^2 + 3 \max\{1, B^2\} (t - t_0) \int_{t_0}^t \mathbb{E} \|f(s, x(s))\|^2 ds + 3 \max\{1, B^2\} (t - t_0) \mathcal{C}_2 \\
 &\times \int_{t_0}^t \mathbb{E} \left[\left\| \int_0^T g(\eta, x(s+\eta)) d\eta \right\| ds \right]^2 \\
 &\leq 3B^2 \|x_0 - h(x)\|^2 + 3 \max\{1, B^2\} (T - \delta) \int_{t_0}^t \mathbb{E} \|f(s, x(s))\|^2 ds \\
 &+ 3 \max\{1, B^2\} (T - \delta) \mathcal{C}_2 \int_{t_0}^t \mathbb{E} \left[\left\| \int_0^T g(\eta, x(s+\eta)) d\eta \right\| ds \right]^2 \\
 &\leq 3B^2 \|x_0 - h(x)\|^2 + 6 \max\{1, B^2\} (T - \delta) \mathcal{L}_0 \int_{t_0}^t \mathbb{E} \|x(s)\|^2 ds + 6 \max\{1, B^2\} (T - \delta)^2 \mathcal{M}_0 \\
 &+ 6 \max\{1, B^2\} (T - \delta) \mathcal{L}_1 \mathcal{C}_2 \int_{t_0}^t \mathbb{E} \|x(s+\eta)\|^2 ds + 6 \max\{1, B^2\} (T - \delta)^2 \mathcal{C}_2 \mathcal{M}_1
 \end{aligned}$$

Thus,

$$\begin{aligned}
 \sup_{t \in [\delta, T]} \mathbb{E} \|\phi x(t)\|^2 &\leq 3B^2 \|x_0 - h(x)\|^2 + 6 \max\{1, B^2\} (T - \delta) \mathcal{L}_0 \int_{t_0}^t \sup_{t \in [\delta, T]} \mathbb{E} \|x(s)\|^2 ds + 6 \max\{1, B^2\} \\
 &\times (T - \delta)^2 \mathcal{M}_0 + 6 \max\{1, B^2\} (T - \delta) \mathcal{L}_1 \mathcal{C}_2 \int_{t_0}^t \sup_{t \in [\delta, T]} \mathbb{E} \|x(s+\eta)\|^2 ds \\
 &+ 6 \max\{1, B^2\} (T - \delta)^2 \mathcal{C}_2 \mathcal{M}_1, \quad \delta \leq t \leq T.
 \end{aligned}$$

Hence ϕ maps \mathbb{B} into itself. Now we have to show that ϕ is a contraction mapping,

$$\begin{aligned}
 & \|\phi x(t) - \phi y(t)\|^2 \\
 & \leq \left[\sum_{k=0}^{+\infty} \left\| \prod_{i=1}^k b_i(\delta_i) \right\| \|\mathfrak{h}(x) - \mathfrak{h}(y)\| I_{[\xi_k, \xi_{k+1})}(t) \right]^2 + \left[\sum_{k=0}^{+\infty} \left[\sum_{i=1}^k \left\| \prod_{j=i}^k b_j(\delta_j) \right\| \right. \right. \\
 & \quad \times \left. \left. \int_{\xi_{i-1}}^{\xi_i} \|f(s, x(s)) - f(s, y(s))\| ds + \int_{\xi_k}^t \|f(s, x(s)) - f(s, y(s))\| ds \right] I_{[\xi_k, \xi_{k+1})}(t) \right]^2 \\
 & \quad + \left[\sum_{k=0}^{+\infty} \left[\sum_{i=1}^k \left\| \prod_{j=i}^k b_j(\delta_j) \right\| ds \int_{\xi_{i-1}}^{\xi_i} \left\| \int_0^T g(\eta, x(s+\eta)) - g(\eta, y(s+\eta)) d\omega(\eta) \right\| \right. \right. \\
 & \quad + \left. \left. \int_{\xi_k}^t \left\| \int_0^T g(\eta, x(s+\eta)) - g(\eta, y(s+\eta)) d\omega(\eta) \right\| ds \right] I_{[\xi_k, \xi_{k+1})}(t) \right]^2 \\
 & \leq \max_k \left\{ \prod_{i=1}^k \|b_i(\delta_i)\|^2 \right\} \|\mathfrak{h}(x) - \mathfrak{h}(y)\|^2 + \left[\max_{i,k} \left\{ 1, \prod_{j=i}^k \|b_j(\delta_j)\| \right\} \right]^2 \\
 & \quad \times \left[\int_{t_0}^t \|f(s, x(s)) - f(s, y(s))\| ds I_{[\xi_k, \xi_{k+1})}(t) \right]^2 + \left[\max_{i,k} \left\{ 1, \prod_{j=i}^k \|b_j(\delta_j)\| \right\} \right]^2 \\
 & \quad \times \left[\int_{t_0}^t \left\| \int_0^T g(\eta, x(s+\eta)) - g(\eta, y(s+\eta)) d\omega(\eta) \right\| ds I_{[\xi_k, \xi_{k+1})}(t) \right]^2 \\
 & \leq B^2 \|\mathfrak{h}(x) - \mathfrak{h}(y)\|^2 + \max \{1, B^2\} (t - t_0) \int_{t_0}^t \|f(s, x(s)) - f(s, y(s))\|^2 ds \\
 & \quad + \max \{1, B^2\} (t - t_0) \mathcal{C}_2 \int_{t_0}^t \left\| \int_0^T g(\eta, x(s+\eta)) - g(\eta, y(s+\eta)) d\eta \right\|^2 ds.
 \end{aligned}$$

However,

$$\begin{aligned}
 & \mathbb{E} \|\phi x(t) - \phi y(t)\|^2 \\
 & \leq B^2 \mathbb{E} \|\mathfrak{h}(x) - \mathfrak{h}(y)\|^2 + \max \{1, B^2\} (t - t_0) \int_{t_0}^t \mathbb{E} \|f(s, x(s)) - f(s, y(s))\|^2 ds \\
 & \quad + \max \{1, B^2\} (t - t_0) \mathcal{C}_2 \int_{t_0}^t \mathbb{E} \left\| \int_0^T g(\eta, x(s+\eta)) - g(\eta, y(s+\eta)) d\eta \right\|^2 ds \\
 & \leq B^2 \mathbb{E} \|\mathfrak{h}(x) - \mathfrak{h}(y)\|^2 + \max \{1, B^2\} (T - \delta) \mathcal{L}_0 \int_{t_0}^t \mathbb{E} \|x(s) - y(s)\|^2 ds \\
 & \quad + \max \{1, B^2\} (T - \delta) \mathcal{C}_2 \mathcal{L}_1 \int_{t_0}^t \mathbb{E} \|x(s+\eta) - y(s+\eta)\|^2 ds.
 \end{aligned}$$

Taking supremum over t we would obtain,

$$\begin{aligned}
 \|\phi x - \phi y\|^2 &\leq B^2 \mathcal{L}_* \|x - y\|^2 + \max\{1, B^2\} (T - \delta)^2 \mathcal{L}_0 \|x - y\|^2 \\
 &\quad + \max\{1, B^2\} (T - \delta)^2 \mathcal{C}_2 \mathcal{L}_1 \|x - y\|^2 \\
 &\leq [B^2 \mathcal{L}_* + \max\{1, B^2\} (T - \delta) \mathcal{L}_0 + \max\{1, B^2\} (T - \delta) \mathcal{C}_2 \mathcal{L}_1] \|x - y\|^2 \\
 &\leq [B^2 \mathcal{L}_* + \max\{1, B^2\} (T - \delta)^2 \{\mathcal{L}_0 + \mathcal{C}_2 \mathcal{L}_1\}] \|x - y\|^2 \\
 &\leq \Gamma \|x - y\|^2,
 \end{aligned}$$

where

$$\Gamma = B^2 \mathcal{L}_* + \max\{1, B^2\} (T - \delta)^2 \{\mathcal{L}_0 + \mathcal{C}_2 \mathcal{L}_1\}$$

By (H5) and $0 < \Gamma < 1$ we would obtain ϕ is a contraction mapping. By Banach fixed theorem ϕ has a unique fixed point on \mathbb{B} . Hence the system has a unique mild solution. \square

Remark 3.2. Let $f : \mathbb{R}_\delta \times \mathbb{X} \rightarrow \mathbb{X}$, $g : \mathbb{R}_\delta \times \mathbb{X} \rightarrow \mathbb{X}$ and $h : \mathbb{X} \rightarrow \mathbb{X}$ satisfy the assumptions (H1)-(H5). Then there exists a unique, global, continuous solution x to the system 1.2 for any initial value (t_0, x_0) with $t_0 \geq 0$ and $x_0 \in \mathbb{B}$.

Remark 3.3. Assume that (H1)-(H5) holds. Then the mild solution without existence of non-local condition and the solution is

$$\begin{aligned}
 x(t) &= \sum_{k=0}^{+\infty} \left[\prod_{i=1}^k b_i(\delta_i) x_0 + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} f(s, x(s)) ds \right. \\
 &\quad + \int_{\xi_k}^t f(s, x(s)) ds + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \\
 &\quad \times \left. \int_{\xi_{i-1}}^{\xi_i} \int_0^T g(\eta, x(s + \eta)) d\omega(\eta) ds + \int_{\xi_k}^t \int_0^T g(\eta, x(s + \eta)) d\omega(\eta) ds \right] I_{[\xi_k, \xi_{k+1})}(t),
 \end{aligned}$$

where $t \in [\delta, T]$.

4. Stability

Theorem 4.1. Let $x(t)$ and $\hat{x}(t)$ be solution of the system 1.2 with initial value $x_0 - h(x)$ and $\hat{x}_0 - h(\hat{x})$ respectively. If the assumptions (H1)-(H4) of Theorem 3.1 are satisfied, then the system 1.2 is stable in the mean square.

Proof. From the assumptions $x(t)$ and $\hat{x}(t)$ are two solutions of the system 1.2 for every

$t \in [\delta, T]$. Then,

$$\begin{aligned}
 x(t) - \hat{x}(t) &= \sum_{k=0}^{+\infty} \left[\prod_{i=1}^k b_i(\delta_i)(x_0 - \hat{x}_0) + (h(x) - h(\hat{x})) \right. \\
 &+ \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} [f(s, x(s)) - f(s, \hat{x}(s))] ds \\
 &+ \int_{\xi_k}^t [f(s, x(s)) - f(s, \hat{x}(s))] ds + \sum_{i=1}^k \prod_{j=i}^k b_j(\delta_j) \int_{\xi_{i-1}}^{\xi_i} \int_0^T [g(\eta, x(s+\eta)) \\
 &- g(\eta, \hat{x}(s+\eta))] d\omega(\eta) ds + \int_{\xi_k}^t \int_0^T [g(\eta, x(s+\eta)) - g(\eta, \hat{x}(s+\eta))] \\
 &\times d\omega(\eta) ds \Big] I_{[\xi_k, \xi_{k+1})}(t).
 \end{aligned}$$

Using the assumed hypotheses (H1)-(H4) we would obtain,

$$\begin{aligned}
 \|x(t) - \hat{x}(t)\|^2 &\leq 4 \sum_{k=0}^{+\infty} \left[\left\| \prod_{i=1}^k b_i(\delta_i) \right\| [\|x_0 - \hat{x}_0\| + \|h(x) - h(\hat{x})\| \right. \\
 &+ \sum_{i=1}^k \prod_{j=i}^k \|b_j(\delta_j)\| \int_{\xi_{i-1}}^{\xi_i} \|f(s, x(s)) - f(s, \hat{x}(s))\| ds \\
 &+ \int_{\xi_k}^t \|f(s, x(s)) - f(s, \hat{x}(s))\| ds + \sum_{i=1}^k \prod_{j=i}^k \|b_j(\delta_j)\| \\
 &\times \int_{\xi_{i-1}}^{\xi_i} \left\| \int_0^T [g(\eta, x(s+\eta)) - g(\eta, \hat{x}(s+\eta))] d\omega(\eta) \right\| ds \\
 &+ \left. \int_{\xi_k}^t \left\| \int_0^T [g(\eta, x(s+\eta)) - g(\eta, \hat{x}(s+\eta))] d\omega(\eta) \right\| ds \right]^2 \\
 &\times I_{[\xi_k, \xi_{k+1})}(t),
 \end{aligned}$$

and

$$\begin{aligned}
 \mathbb{E} \|x(t) - \hat{x}(t)\|^2 &\leq 4 \max_k \left\{ \left\| \prod_{i=1}^k b_i(\delta_i) \right\|^2 \right\} \left[\mathbb{E} \|x_0 - \hat{x}_0\|^2 + \mathbb{E} \|h(x) - h(\hat{x})\|^2 \right] \\
 &+ 4 \max_{i,k} \left\{ 1, \prod_{j=i}^k \|b_j(\delta_j)\| \right\}^2
 \end{aligned}$$

$$\begin{aligned}
 & \times \mathbb{E} \left\{ \int_{t_0}^t \|f(s, x(s)) - f(s, \hat{x}(s))\| ds I_{[\xi_k, \xi_{k+1})}(t) \right\}^2 + 4 \max_{i,k} \left\{ 1, \prod_{j=i}^k \|b_j(\delta_j)\| \right\}^2 \\
 & \times \mathbb{E} \left\{ \int_{t_0}^t \left\| \int_0^T [g(\eta, x(s+\eta)) - g(\eta, \hat{x}(s+\eta))] d\omega(\eta) \right\| ds I_{[\xi_k, \xi_{k+1})}(t) \right\}^2 \\
 & \leq 4B^2 \left[\mathbb{E} \|x_0 - \hat{x}_0\|^2 + \mathbb{E} \|h(x) - h(\hat{x})\|^2 \right] + 4 \max \{1, B^2\} (t - t_0) \\
 & \times \int_{t_0}^t \|f(s, x(s)) - f(s, \hat{x}(s))\|^2 ds + 4 \max \{1, B^2\} (t - t_0) \mathcal{C}_2 \\
 & \times \int_{t_0}^t \left\| \int_0^T [g(\eta, x(s+\eta)) - g(\eta, \hat{x}(s+\eta))] d\eta \right\|^2 ds \\
 & \leq 4B^2 \left[\mathbb{E} \|x_0 - \hat{x}_0\|^2 + \mathbb{E} \|h(x) - h(\hat{x})\|^2 \right] + 4 \max \{1, B^2\} (t - t_0) \mathcal{L}_0 \\
 & \times \int_{t_0}^t \mathbb{E} \|x(t) - \hat{x}(t)\|^2 ds + 4 \max \{1, B^2\} (t - t_0) \mathcal{L}_1 \mathcal{C}_2 \\
 & \times \int_{t_0}^t \mathbb{E} \|x(t+\eta) - \hat{x}(t+\eta)\|^2 ds.
 \end{aligned}$$

Taking supremum over t ,

$$\begin{aligned}
 & \sup_{t \in [\delta, T]} \mathbb{E} \|x(t) - \hat{x}(t)\|^2 \\
 & \leq 4B^2 \left[\mathbb{E} \|x_0 - \hat{x}_0\|^2 + \sup_{t \in [\delta, T]} \mathbb{E} \|h(x) - h(\hat{x})\|^2 \right] + 4 \max \{1, B^2\} \\
 & \times (T - \delta) \mathcal{L}_0 \int_{t_0}^t \sup_{t \in [\delta, T]} \mathbb{E} \|x(t) - \hat{x}(t)\|^2 ds + 4 \max \{1, B^2\} \\
 & \times (T - \delta) \mathcal{L}_1 \mathcal{C}_2 \int_{t_0}^t \sup_{t \in [\delta, T]} \mathbb{E} \|x(t+\eta) - \hat{x}(t+\eta)\|^2 ds.
 \end{aligned}$$

By Grownwall inequality,

$$\begin{aligned}
 \sup_{t \in [\delta, T]} \mathbb{E} \|x(t) - \hat{x}(t)\|^2 & \leq 4B^2 \mathbb{E} \|x_0 - \hat{x}_0\|^2 \exp [4 \max \{1, B^2\} (T - \delta)^2] \mathcal{L} \\
 & \leq \Gamma \mathbb{E} \|x_0 - \hat{x}_0\|^2.
 \end{aligned}$$

where, $\Gamma = 4B^2 \exp [4 \max \{1, B^2\} (T - \delta)^2] \mathcal{L}$ and $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1 \mathcal{C}_2 + \frac{\mathcal{L}_*}{(T - \delta)^2}$.

For given $\epsilon > 0$, we choose $\alpha = \frac{\epsilon}{\Gamma}$ such that $\mathbb{E} \|x_0 - \hat{x}_0\|^2 < \alpha$ then,

$$\sup_{t \in [\delta, T]} \mathbb{E} \|x(t) - \hat{x}(t)\|^2 \leq \epsilon$$

.

□

Remark 4.2. Random impulsive integro-differential equation with local initial condition is a special case of the system 1.2. So the random impulsive integro-differential equation with local initial condition is stable in the mean square.

5. conclusion

In this paper, the uniqueness and stability of random impulsive stochastic integro-differential system has been calculated using the initial data $x_{t_0} = x_0$. Contraction mapping principle is used to prove the existence and uniqueness. With inequality techniques and Contraction principle the stability is derived. In the future we can extend this work to stochastic partial differential equations.

References

- [1] Ahmad. B and Alghamdi. B.S (2008). *Approximation of solutions of the nonlinear Duffing equation involving both integral and nonlinear forcing terms with separated boundary conditions*, Computer Physics Communications, **179**(6): 409-416. <https://doi.org/10.1016/j.cpc.2008.04.008>
- [2] Ahmad. B (2008). *On the existence of T-periodic solutions for duffing type integrodifferential equations with p-Laplacian*, Lobachevskii Journal of Mathematics, **29**(1): 1-4. <https://doi.org/10.1134/S1995080208010010>
- [3] Haloi. R, Dwijendra. N Pandey and Bahuguna. D (2012). *Existence and Uniqueness of a Solution for a Non-Autonomous Semilinear Integrodifferential Equation with Deviated Argument*, Differ. Equ. Dyn. Syst., **20**(1): 1-16. <https://doi.org/10.1007/s12591-011-0099-x>
- [4] Yan. Z (2010). *Nonlocal Problems for Delay integrodifferential equations in Banach spaces*, Differ. Eqn, Appl., **2**(1): 15-24. <https://doi.org/10.7153/dea-02-02>
- [5] Anguraj. A and Karthikeyan. K (2009). *Existence of solutions for impulsive neutral functional differential equations with non-local conditions*, Nonlinear Analysis Theory methods and Applications., **70**(7): 2717-2721. <https://doi.org/10.1016/j.na.2008.03.059>
- [6] Lakshmikantham. V, Bairov. D.D and Simeonov. P.S (1989). *Theory of Impulsive Differential Equations*, World Scientific, Singapore. <https://doi.org/10.1142/0906>
- [7] Samoilenko.A.M and Perestyuk. N.A (1995). *Impulsive Differential Equations*, World Scientific, Singapore.
- [8] Anguraj. A, Mallika Arjunan. M and Hernandez. E (2007). *Existence results for an impulsive partial neutral functional differential equations with state-dependent delay*, Applicable Analysis., **86**(7): 861-872. <https://doi.org/10.1080/00036810701354995>
- [9] Hernandez. E, Marco Rabello and Henriques. H.R (2007). *Existence of solutions for impulsive partial neutral functional differential equations*, J.Math.Anal.Appl., **331**: 1135-1158. <https://doi.org/10.1016/j.jmaa.2006.09.043>
- [10] Wu. S.J and Meng. X.Z (2004). *Boundedness of nonlinear differential systems with impulsive effect on random moments*, Acta Mathematicae Applicatae Sinica., **20**(1): 147-154. <https://doi.org/10.1007/s10255-004-0157-z>
- [11] Mao. X (1997). *Stochastic Differential Equations and Applications*, M.Horwood, Chichester.
- [12] Da Prato. G and Zabczyk. J (1992). *Stochastic Equations in Infinite Dimensions*, Cambridge: Cambridge University Press. <https://doi.org/10.1017/CBO9781107295513>
- [13] Oksendal. B (2013). *Stochastic differential Equations: An introduction with Applications*, Springer Science and Business Media.
- [14] Yang. X and Zhu. Q (2014). *p-th moment exponential stability of stochastic partial differential equations with poisson jumps*, Asian J. Control, **16**: 1482-1491. <https://doi.org/10.1002/asjc.918>
- [15] Boudaoui. A, Caraballo. G. T and Ouahab. A (2017). *Stochastic differential equations with non instantaneous impulses driven by a fractional brownian motion*, Discrete contin. Dyn. Syst. B, **22**: 2521-2541. <https://doi.org/10.3934/dcdsb.2017084>
- [16] Djourdem. H and Bouteraa. N (2020). *Mild solution for a stochastic partial differential equation with noise*, WSEAS Trans. Syst., **19**: 246-256. <https://doi.org/10.37394/23202.2020.19.29>

- [17] Bouteraa. N, Mustafa Inc, Ali Akinlar. M and Bin-Mohsin. B (2021). *Mild solution for a fractional partial differential equation with noise*, Math Meth Appl Sci, **44**:5648-5662. <https://doi.org/10.1002/mma.7139>
- [18] Bouteraa. N, Mustafa Inc and Ali Akgul (2021). *Stability analysis of time-fractional differential equations with initial data*, Math Meth Appl Sci, **1-9**. <https://doi.org/10.1002/mma.7782>
- [19] Bianov. D. D and Simeonov. P. S (1989). *Systems with Impulse effect: Stability, Theory and Applications*.
- [20] Naghshtabrizi. P, Hespanha. J. P and Teel. A. R (2008). *Exponential stability of impulsive systems with application to uncertain sampled-data systems*, Systems Control Lett., **57(5)**:378-385. <https://doi.org/10.1016/j.sysconle.2007.10.009>
- [21] Sakthivel. R and Luo. J (2009). *Asymptotic stability of nonlinear impulsive stochastic differential equations*, Statist. Probab. Lett. **79(9)**:1219-1223. <https://doi.org/10.1016/j.spl.2009.01.011>
- [22] Pan. L and Cao. J (2011). *Exponential stability of impulsive stochastic functional differential equations*, J. Math. Appl., **382**: 672-685. <https://doi.org/10.1016/j.jmaa.2011.04.084>
- [23] Cui. J and Yan. L (2011). *Existence result for fractional neutral stochastic integro-differential equations with delay*, Journal of Physics A:Mathematical and Theoretical. **44**. <https://doi.org/10.1088/1751-8113/44/33/335201>
- [24] Mao. X (2007). *Stability of Stochastic integro-differential equations*, Stochastic Analysis and applications, **18**:1005-1017. <https://doi.org/10.1080/07362990008809708>
- [25] Li. S, Shu. L, Shu. X.B and Xu. F (2019). *Existence and Hyers-Ulam stability of random impulsive stochastic functional differential equations with finite delays*, An International Journal of Probability and Stochastic Processes. <https://doi.org/10.1080/17442508.2018.1551400>
- [26] Wang. T and Wu. S (2008). *Random impulsive model for stock prices and its applications for insurers*, Master thesis(in Chinese), Shanghai, East China Normal University.
- [27] El-Borai. M (2006). *On some stochastic fractional integro-differential equations*, Advances in dynamical systems and applications, **1**: 49-57.
- [28] Ahmed. M and El-Borai. M (2018). *Hilfer fractional stochastic integro-differential equations*, Applied mathematics and computation, **331**: 182-189. <https://doi.org/10.1016/j.amc.2018.03.009>
- [29] Guo. Y, Zhu. Q and Wang. F (2020). *Stability analysis of impulsive stochastic functional differential equations*, Commun Nonlinear Sci Number Simulat. **82**. <https://doi.org/10.1016/j.cnsns.2019.105013>
- [30] Liu. J and Xu. W (2020). *An averaging result for impulsive fractional neutral stochastic differential equations*, Applied Mathematics Letters. <https://doi.org/10.1016/j.aml.2020.106892>
- [31] Jose. S. A, Yukunthorn. W, Naploes Valdes. J. E and Leiva. H (2020). *Some existence, uniqueness and stability results of nonlocal random impulsive integro-differential equations*, Applied Mathematics E-Notes, **20**: 481-492. <http://www.math.nthu.edu.tw/amen/>
- [32] Guo. D. J (2003). *Multiple positive solutions for first order nonlinear integro-differential equations in Banach spaces*, Nonlinear Anal., **53**: 183-195. [https://doi.org/10.1016/S0362-546X\(02\)00165-7](https://doi.org/10.1016/S0362-546X(02)00165-7)
- [33] Guo. D. J (2006). *Extremal solutions for th-order impulsive integro-differential equations on the half-line in Banach spaces*, Nonlinear Analysis, **65(3)**:677-696. <https://doi.org/10.1016/j.na.2005.09.032>