



Relationship of Soilborne Mycoflora of Cassava Growing Fields to Incidence of Postharvest Rots of Cassava Tubers in Sokoto, Nigeria

¹Maccido Ibrahim and ²Kasimu Shehu

¹Department of Biology, Shehu Shagari College of Education, Sokoto, Nigeria; ²Department of Biological Sciences, Usmanu Danfodiyo University, Sokoto, Nigeria. Corresponding author : bmaccido88@gmail.com

Received : September 18, 2014

Accepted : November 17, 2014

Abstract - In this study the fungi associated with cassava growing fields in Sokoto were isolated and identified using soil dilution technique. A total of 215 fungal colonies from 9 fungal species were isolated from soil samples of different cassava fields between the month of June and August, 2012. It was observed that soil samples from Lambara recorded the highest number (64) of fungal species followed by Damba and the least number (44) of fungal species was observed in Wamakko. The fungi isolated were *Alternaria species*, *Aspergillus niger*, *Aspergillus fumigatus*, *Cylindrocarpum lichenicola*, *Fusarium oxysporum*, *Geotrichum candidum*, *Mucor hiemalis*, *Rhizopus oryzae* and *Scopulariopsis candida*. The highest percentage frequency of occurrence was observed in *Aspergillus niger* (39.5%) seconded by *Fusarium oxysporum* (18.2%) and the least was seen in *Rhizopus oryzae* (2.3%). The pathogenicity test indicated that all the fungal isolates were pathogenic on cassava tubers. The fungus *M. hiemalis* is the most pathogenic followed by *F. oxysporum* and the least was recorded by *R. oryzae*. Therefore, it would be concluded that there is relationship between soilborne fungi and incidence of postharvest rots of cassava tubers.

Keywords : Soilborne mycoflora; Cassava fields; Rots; Cassava tubers

Introduction

Fungi being ubiquitous organisms occur in all types of habitats and are the most adaptable organisms. The soil is one of the most important habitats for microorganisms like fungi, bacteria, viruses, nematodes etc. They are found in large numbers in soil usually between one to ten million microorganisms are present per gramme of soil (Ishaq and Khan, 2011). Many soil fungi are potential pathogens to both plants and animals (Moalaei *et al.*, 2006). Six pathogenic fungi were isolated from yam harvested soil in Ibadan. These fungi include *Penicillium oxalicum*, *P. cyclopium*, *A. niger*, *R. nigrican*, *Fusarium sp* and *Mucor circinelloides* (Odebode, 2002). Pandey *et al.* (2001) found species of *Penicillium* and *Trichoderma* from soils of tea growing locations in India. Similarly Karaoglu and Ulker (2006) shown that *Mycellia sterilla*, *Gliocladium virens* and *Trichoderma* were isolated from tea growing areas in Turkey. The findings of Moallaei *et al.* (2006) revealed that *A. flavus*, *Fusarium oxysporum*, *Anixiopsis stercoraria*, *Arthroderma guniculi*, *Reniospora flavissima*, *Chrysosporium keratinophilum*, *Trichophyton vanbreuseghemii*, *Geotrichum candidum*, *Microsporium gypseum* and *Penicillium* species were isolated from soil samples of forests and farm yards in Iran. Gaddeyya *et al.* (2012) reported *A. flavus*, *A. fumigatus*, *A. niger*, *A. nidulans*, *A. terreus*, *Penicillium chrysogenum*, *P. frequentans*, *P. funiculosum*, *Trichoderma viride*, *T. harzianum*, *Fusarium oxysporum*, *F. solani*, *Curvalaria clavata*, *C. lunata* and *Rhizopus stolonifer* from different crop fields at Salur Mandal, India.

Cassava (*Manihot esculentum* Crantz) is the most important root crop and source of energy for human consumption. Cassava is a major food crop for an estimated 500 million people in Africa (Hillocks *et al.*, 2012). Root crops are used in the preparation of major staples eaten in Nigeria and are estimated to contribute about 15% of total protein in daily diet of the average Nigerian (Ogunmodele, 1983). Cassava can be used in the production of starch, ethanol, confectionaries and animal feeds (FAO, 2001). Production of Cassava is hampered by diseases caused by fungi,

bacteria and viruses. Among serious fungal diseases root rots affected both young and old cassava plants (Msikita *et al.*, 2005). Cassava root rots are caused by a complex of soilborne pathogens which induced damages that eventually reduce the yield (Ambe, 1994). Okigbo *et al.* (2009a) isolated *Fusarium oxysporum*, *Botryodiplodia theobromae*, *Macrophina phaseolina*, *Penicillium oxalicum* and *Aspergillus niger* from rotted cassava tubers. Msikita *et al.* (2005) reported *Nattrassia mangifera*, *Macrophina phaseolina*, *Botryodiplodia theobromae*, *Fusarium sp.* and *Pythium* from the root of cassava in Benin. The finding of Lozano and Nolt (2000) showed that *Sclerotium rolfsii*, *Roselina necatrix*, *Fusarium moniliformae*, *F. oxysporum*, *Phytophthora erythroseptica*, *P. cryptogea*, *P. drechsleri*, *P. nicotina* were responsible for dry and wet tuber rot of cassava. Salami and Akintokun (2008) found that *Lasiodiplodia theobromae*, *Macrophina phaseolina*, *Rhizopus stolonifer*, and *Fusarium pallidoroseum* were the fungal pathogens causing postharvest rot of cassava in Southwestern Nigeria.

Furthermore, Kolawole *et al.* (2009) isolated from rotten cassava *Aspergillus niger*, *A. flavus*, *R. stolonifer*, *Mucor racemosus* and *F. oxysporum*. While *Phytophthora palmivora*, *Fusarium solani* and *Sclerotium rolfsii* were isolated from infected cassava tubers (Johnson *et al.*, 2002). Banitol *et al.* (2010) confirmed the presence of *B. theobromae*, *Sclerotium rolfsii*, *Fusarium sp.* and *Pythium sp.* in diseased roots and stems of cassava in ecozones of Togo. Studies carried out by Fokunang *et al.* (2002) and Udoudoh (2011) revealed that *Colletotrichum gloeosporioides*, *Aspergillus sp.* and *Penicillium sp.* were associated with postharvest storage and spoilage of cassava tubers. *Botryodiplodia acerina* has been reported to cause postharvest rot of cassava tubers (Amadioha and Markson, 2007). Snodow (1992) reported that *A. flavus*, *M. hiemalis*, *Rhizopus oryzae*, *B. theobromae* and *Fusarium solani* caused fungal rot of cassava tubers. This paper is aimed at identifying fungi associated with cassava fields in Sokoto and their relationship to postharvest deterioration of cassava.

Materials and Methods

Study area

Sokoto state is located on longitude 3 – 9° east and latitude 10 - 14° north at an altitude of 398m (NMA, Sokoto 2009). The area has an annual average temperature of 27 °C. The temperature reaches peak in April at 40 °C and drops to 18 °C in December to January (Kowal and Knabe, 2002). The area has average annual rainfall of 550mm. The rainfall start in June, reaches peak in August and end in early September (MOI, 2008; NMA, 2009; SARDA,2009). The relative humidity is about 16-55.5% during dry season but can rise up to 81% during rainy season (SARDA ; NMA, 2009).

Collection of Soil Samples

Soil samples were taken from 5 different cassava fields in 5 locations in each of the collection areas. The collection areas are Damba in Illela local government, Raka in Tangaza local government, Wamakko in Wamakko local government and Lambara in Shagari local government areas of Sokoto State.

The soil samples were collected monthly (June to August) from the layer with depth not exceeding 0-9cm. The soil samples were transferred immediately to the laboratory in sterile polythene bags.

Isolation of fungi from soil sample

Soil dilution plate method was used for the isolation of fungi. One gramme of soil sample was suspended in 10ml distilled water in test tubes. The soil samples were shaken and serially diluted to 10⁶ samples. One milliliter of each sample was pipetted onto sabouraud dextrose agar and incubated at 28 °C for 72 hours. Characteristic colony types were subcultured to obtained pure cultures of the isolates (Odebode, 2002).

Identification of soil fungi

The isolated fungi were identified based on the isolate colonial characteristics (colour and texture) and microscopic features such as nature of mycelium, type of fruiting bodies and the spore structure. The isolates were identified by reference to Kora *et al.* (2005).

Pathogenicity test

Fresh and healthy tubers of cassava were washed with tap water and surface sterilized with 70% ethanol. Cylindrical cores were removed from the tubers by means of sterile 5mm cork borer. 4mm agar discs containing 7 days old cultures of the isolates were introduced singly into the holes and sealed with sterile vaseline. Controls were set up as described except that the inocula consist of uninoculated potato dextrose agar blocks. All treated tubers were put singly in sterile polythene bags and incubated at 28 °C for 12 days. Three replicate tubers were cut through and examined for rot at 3 days intervals (Udo *et al.*, 2000).

Statistical analysis

The number of colonies per plate in 1g of soil was calculated. The percentage frequency of each isolate was calculated using the following formula:

$$\% \text{ frequency} = \frac{\text{Total no. of individual species}}{\text{Total no. of all species}} \times 100$$

The results of pathogenicity test were analysed using one-way analysis of variance (ANOVA) via instat 3a statistical package and least significance difference was used for mean separation at $P < 0.05$.

Results and Discussion

The results show that the number of fungal species was found high in soil samples of cassava fields from Lambara (60) seconded by Damba (56) the least was recorded by Wamakko (44). The fungi isolated include *Alternaria species*, *Aspergillus fumigatus*, *Aspergillus niger*, *Cylindrocarpon lichenicola*, *Fusarium oxysporum*, *Geotrichum candidum*, *Mucor hiemalis*, *Rhizopus oryzae* and *Scopulariopsis candida*. The highest frequency of occurrence was observed in *A. niger* (39.5%) followed by *Fusarium oxysporum* (18.2%) and least percentage frequency of occurrence was recorded by *R. oryzae* (2.3%). It was observed that *A. niger* occurred in almost all the months between June to August. While *F. oxysporum* and *Alternaria species* occurred more frequently in the humid months of July and August. The results were shown in Table 1.

Similarly Ogunmwoyi *et al.* (2008) they found *A. niger* as dominant fungi among fungi isolated from soil samples from Obafemi Awolowo University Ife. Also Sharma (2010) and Ishaq and Khann (2011) reported that *A. niger* and *A. fumigatus* were dominant among fungal species isolated from soil samples from Katao and agricultural fields at Ramgahn, India. The ability of *A. niger* to dominate other fungal species could be linked to its high sporulating capacity and ability to produce toxins which prevent the growth of other fungi (Gaddeyya *et al.*, (2012). Studies carried out by Pandey *et al* (2001) and Koraoglu and Ulker (2006) indicated that species of *Penicillium*, *Trichoderma*, *Aspergillus* and *Fusarium* were isolated from soil of tea growing areas in India and Turkey. The occurrence of these fungal species in the soil was influenced by some factors such as temperature, pH, organic contents and moisture (Okigbo, 2003 Gaddeyya *et al.*, 2012).

The results of the study indicates that cassava fields from Lambara has the highest frequency of fungal species followed by Damba and the least was recorded by Wamakko. The monthly (June – August) distribution of fungal species showed the highest frequency of fungal species occurred in August (85) and the least frequency of occurrence was observed in June (52). The distribution of the fungal species was found to be influenced by environmental factors such as temperature and moisture (Malik and Singh, 2004). At the inception of rainfall there was an increase in the population of the fungi. The fungi isolated during the month of June include *A. niger*, *A. fumigatus* and *C. lichenicola*. While fungi such as *F. oxysporum*, *M. hiemalis* and *Alternaria species* were found to be dominant in the humid months of July and August. The increase in percentage of fungi during humid months could be linked to the increase in the soil pH which affects availability of soil nutrients (Odebo, 2002).

Table 1. Fungi isolated from cassava fields

Organisms	June				July Collection Areas				August				Frequency (%)	
	DB	RK	WM	LM	DB	RK	WM	LM	DB	RK	WM	LM		
<i>Alternaria species</i>	2	-	-	-	3	-	1	1	8	3	1	-	19	8.8
<i>Aspergillus fumigatus</i>	3	-	-	5	3	-	-	-	3	-	-	-	14	6.5
<i>Aspergillus niger</i>	9	6	9	6	5	7	5	6	8	6	10	8	85	39.5
<i>Cylindricarpon lichenicola</i>	-	-	-	4	-	-	-	2	-	-	-	-	6	2.8
<i>Fusarium oxysporum</i>	-	-	-	1	-	7	6	5	-	11	-	9	39	18.2
<i>Geotrichum candidum</i>	-	2	-	-	-	-	3	-	-	2	-	-	7	3.3
<i>Mucor hiemalis</i>	-	2	-	-	8	2	2	9	-	5	4	2	34	15.8
<i>Rhizopus oryzae</i>	-	-	1	-	-	-	2	-	1	-	-	1	5	2.3
<i>Scopulariopsis candida</i>	-	2	-	-	1	-	-	-	2	-	-	1	6	2.8
<i>Total</i>	14	12	10	16	20	16	19	23	22	27	15	21	215	100

Key : DB = Damba ; RK = Raka ; WM : Wamakko; LM = Lambara

Table 2. Pathogenicity of fungi isolated from cassava fields on cassava tubers (Diameter of rot in mm).

Mean of the determination with three replications \pm Standard Error. Least Significant Difference (LSD).

Values with the same alphabet along column are not significantly different at (P<0.05).

Isolates	Incubation Period			
	3 days	6 days	9 days	12 days
<i>Cylindricarpon Lichenicola</i>	2.5 \pm 0.20g	4.5 \pm 0.64d	8.1 \pm 1.10f	11.3 \pm 1.00g
<i>Alternariasppecies</i>	5.0 \pm 0.12e	8.0 \pm 0.36c	14.0 \pm 0.29d	19.0 \pm 0.40e
<i>Aspergillus fumigatus</i>	6.0 \pm 0.12d	14.0 \pm 0.21b	24.0 \pm 1.27c	29.0 \pm 0.15d
<i>Aspergillus niger</i>	7.0 \pm 0.25c	15.0 \pm 0.16a	28.0 \pm 0.66b	36.0 \pm 0.68c
<i>Rhizopus oryzae</i>	2.3 \pm 0.31g	4.5 \pm 0.61d	7.1 \pm 0.89f	10.3 \pm 0.90g
<i>Fusarium oxysporum</i>	8.0 \pm 0.32b	16.0 \pm 0.74a	28.0 \pm 0.26b	38.0 \pm 0.72b
<i>Geotrichum candidum</i>	4.0 \pm 0.15f	7.0 \pm 0.36c	11.0 \pm 0.40e	16.0 \pm 0.47f
<i>Mucor hiemalis</i>	9.0 \pm 0.21a	16.0 \pm 0.38a	33.0 \pm 0.64a	48.0 \pm 0.36a
<i>Scopulariopsis candida</i>	3.0 \pm 0.20g	5.0 \pm 0.79d	12.0 \pm 0.31e	18.0 \pm 0.36e
Control	0.0 \pm 0.0h	0.0 \pm 0.0e	0.0 \pm 0.0g	0.0 \pm 0.0h
LSD(0.05)	0.75	1.66	1.75	1.39

The results of pathogenesis were shown in Table 2. The results indicate that all the fungi isolated from soil samples of cassava fields were found to be pathogenic on cassava tubers. The pathogens showed variation in nature and extent of rots. The fungus *M. hiemalis* is the most pathogenic followed by *F. oxysporum* and *R. oryzae* was the least pathogenic fungus.

It was observed that there is relationship between the soilborne fungi and incidence of post harvest rot of cassava tubers. The soil fungi got to the soil through infected plant parts and thus serve as source of inocula in the soil. These fungal pathogens overwinter in the soil and when cassava plants develop tubers and the tubers are wounded they become infected by the pathogens.

Conclusion

This study reveals that 9 soilborne fungal species were associated with cassava fields in Sokoto. It is apparent that there is relationship between the soilborne fungi and incidence of post harvest roots rot of cassava. Therefore, it would be recommended that cassava fields should be allowed to fallow for some years, alternate host plants should not be allowed to grow in the fields, healthy cassava cuttings should be used for planting to avoid reintroduction of pathogens and the use of resistant planting materials to reduce incidence of rots.

Acknowledgements

The authors are thankful to Mal.Umar Dambuwa of mycology laboratory Usmanu Danfodiyo University, Sokoto for allowing us use their laboratory facilities and his technical assistance during the conduct of the research work.

References

- Amadioha, A.C. and Markson, A.A. (2007). Postharvest control of cassava tuber rot by *Butryodiplodia acerina* using extracts of plant origin. Archives of Phytopathology and Plant Protection, 40(5): 359-366.
- Ambe, J.T. (1994). Effects of harvesting time on cassava fresh root yield in Cameroon. Discovery and Innovation, 6 (3): 315-317
- Banitol, A., Kpemoua, K.E., Bissang, B. and Wydra, K. (2010). Assessment of cassava root and stem rots in ecozones of Togo and evaluation of the pathogen virulence. Pakistan Journal of Botany, 42 (3): 2059-2068.
- FAO (2001). Food Agricultural Organization. The Global Cassava development strategy and implementation plan Vol. 1 FAO, Rome.
- Fokunang, C.N. , Dixon, A. G. O. Ikotun, T., Akem, C. N. A. and Tebe, E. (2002). Rapid screening methods of cassava cultivars for resistance to *Colletotrichum gloeosporoides f. s. manihotis*. Phytopathology, 150: 6- 12
- Gaddeyeyya, G., Hihaiika, P.S, Bharathi, P. and Kuma P.K..R (2012). Isolation and identification of soil mycoflora in different crop fields at Sahur Mandal. Advance in Applied Science Research, 3(4): 2020-2026.
- Hillocks, R.J., Thresh, J.M. and Belloti, A.C. (2012). Cassava, biology, production and utilization. CAB International: 43 - 48
- Ishaq, F. and Khan, A. (2011). Isolation, identification and bacterial strains found in organic and inorganic soil of different agricultural fields. Research in Science and Technology, 3 (11): 3336.
- Johnson, I., Palaniswami, A., Babus, S. and Nandakumar, R.(2002). Relative importance of fungal species associated with cassava tuber rot. International Journal of Tropical Plant Diseases, 18(112): 69-78.
- Karaoglu, S.A. and Ulker, S. (2006). Isolation, identification and seasonal distribution of soilborne fungi in Tea growing Areas of Iyidere-Ikiz dere vicinity, Turkey. Journal of Basic Microbiology, 46 (3): 208-218

- Kolawole, O. M., Adeyemi, B. J., Kayode, R. M. O. and Ajibola, T. B. (2009). The drying effect of colour light frequencies on the nutrients and microbial composition of cassava. *Africa Journal of Agricultural Science*, 4(3): 171-177.
- Kora, C., Medonald, M.R. and Boland, G.I. (2005). Occurrence of fungal pathogens on carrot wooden boxes used for storage. *Plant Pathology*, 54:666-670.
- Kowal, J.M. and Knabe, D.T. (2002). An agroclimatology atlas of the Northern States of Nigeria. Ahmadu Bello University Press, Zaria, Nigeria .
- Lozano, J. C. and Nolt, B. (2000). Disease of cassava (*Manihot esculentum* Crantz) In: Common Names for Plant Diseases. American Pathological Society: 3-38.
- Malik, V.K. and Singh, S.(2004). Effect of temperature and relative humidity on teliospore germination in *Ustilago hordei*. *Journal. Mycology and Plants Pathology*, 34:410-411
- Moallaei, H., Zain, F., Pihet, M., Mahamoudi, M. and Hasemi, J. (2006). Isolation of Keratinophilic fungi from soil samples of Forest and Farm Yards. *Iranian Journal of Public Health*, 35 (4) 62-69
- Ministry of Information (MOI), Sokoto, (2008), Nigeria Diary; 12.
- Msikita, W., Bissang, B., Jates, B.D., Gaimey, H. and Wilkson, H.T. (2005). Prevalence and severity of *Natrassia Magniferae* root and stem rot pathogens of cassava in Benin. *American Pathological Society*, 89 (1): 89-96.
- Nigerian Metereological Agency (NMA ,2009). Sultan Abubakar III International Airport, Sokoto. 1-2.
- Odebode, A.C. (2002). Survival of yam rot inducing fungi isolated from yam harvested soil in Ibadan, southwestern, Nigeria . *Nigerian Journal of Basic and Applied Science*, 11: 109 166.
- Ogunwonyi, I., Igbenesa, N., Ayegoro. O. E. and Odjadjare, E.E. (2008). Microbial analysis different top soil samples of selected sites in Obafemi Awolowo University , Nigeria *Science Research and Essay*, 156(4): 377 – 382
- Ogunmodele, B.K (1983).Demand and availability of food in Nigeria. *In: Nutritional and Food Policy in Nigeria*. Atinwo, T. and Akinyele L. (eds). Published by National Institute of Policy and Strategic Studies, Nigeria.
- Okigbo R.N. (2003). Fungi associated with peels of postharvest Yams in storage. *Global Journal of Pure and Applied Science*, 9 (1): 19-23.
- Okigbo, R. N. , Ramesh, P. and Achusi, C. T. (2009). Postharvest deterioration of cassava and its control using extracts of *Azadirachta indica* and *Aframomum meloqueta*. *Journal of Chemistry*, 6(4) :1274 - 1280
- Pandey, A., Soicol, C.R., Rodirgueleon, J.A. and Nigram, P. (2001). Solid state fermentation in *Biotechnology Fundamentals and Applications*. Asia Tech. Publishers Inc. New Delhi.
- SARDA (2009). Sokoto Agricultural and Rural Development Authority; 1-2
- Salami, A. O. and Akintokun, A. K. (2008). Postharvest Enzymatic Activities of Healthy and Infected Cassava (*Manihot esculentum* Crantz) Tubers. *Emirate Journal of Food and Agriculture*, 20(1): 1-17.
- Sharma, K. and Malish, P. (2010). Isolation of soil mycolflora from Yumthang valley. *Sikkim Laboratory to Land*, 2 (1):42-45.
- Snodow, A.I (1992). *Colour Atlas of Postharvest Diseases and Disorders of Fruits and Vegetables* CRC Press, Bota Raton.
- Udo, S.K. Maidunagu, B.E. and Isemug C.D. (2001). Inhibition of growth and sporulation of fungal pathogens of sweet potato and yam by garlic extract. *Nigerian Journal of Botany*, 4:31- 35.
- Udoudoli. P.J. (2011). Postharvest storage and sprilage of cassava tubers (*Manihot* spp.) in Ikot-Ekperere, Akwa Ibom State. *Nigerian Journal of Environmental Issue and Agriculture in Developing Countries*, 3 (2): 34 – 38.