

BREEDING FIELD CROPS FOR ORNAMENTAL PURPOSE: A CASE IN *Jatropha* spp.

Narathid Muakrong¹⁾, Chamnanr Phetcharat¹⁾, Patcharin Tanya²⁾ and Peerasak Srinives^{2*)}

¹⁾ Plant Breeding Program, Faculty of Agriculture at Kamphaeng Saen, Kasetsart University
Kamphaeng Saen, Nakhon Pathom 73140, Thailand

²⁾ Department of Agronomy, Faculty of Agriculture at Kamphaeng Saen, Kasetsart University
Kamphaeng Saen, Nakhon Pathom 73140, Thailand

^{*)} Corresponding author E-mail: agrpss@yahoo.com

Received: September 9, 2014 / Accepted: October 27, 2014

ABSTRACT

Populations of F₁, F₂ and BC₁F₁ were formed from an interspecific cross between green-flowered *Jatropha curcas* and red-flowered *J. integerrima* (dwarf type) to observe for segregation in flower colors and to select for ornamental segregants. The crossing to produce F₁ and BC₁F₁ was successful only when *J. curcas* was used as the female parent. The suitable time for crossing was between 08:00 – 13:00 o'clock with fruit setting rate of 65-78%. The F₁ segregated into white- and pink-flowered plants. While flowers of the F₂ population segregated into nine different colors, viz. light green, green, white, very light pink, light pink, pink, deep pink, very deep pink, and red. When *J. curcas* was backcrossed by a pink-flowered F₁, its BC₁F₁ progenies segregated into green, light green, white, very light pink and light pink flowers. While another set of BC₁F₁ progenies obtained from a white-flowered F₁ parent segregated into green, light green, and white. The segregating progenies in both sets showed smaller plants than *Jatropha* and bore colorful flowers suitable for ornamental purpose. Six plants were chosen and registered at Kasetsart University and the Thai Department of Agriculture as commercial cultivars, 'Kamphaeng Saen 1' to 'Kamphaeng Saen 6'.

Keywords: crossing durations flower color, ornamental *Jatropha*, interspecific cross

INTRODUCTION

Ornamental plants are plants grown for decoration in vase, hanging, and garden landscape for pleasure and to improve the environment. The demand for ornamental plants follows the country's economic growth and the quality of life. The market of ornamental plants

depends on satisfaction of the buyers, thus kinds and varieties of plants have to be novel in response to the market demand.

Breeding field crops for ornamental purpose is a new idea to develop unique ornamental plants such as red castor bean (Arslan, 2012), multi-colored and spineless safflower (Bradley *et al.*, 1999 and Golkar *et al.*, 2010). In pineapple, ornamental characters include brightly-colored flower head on a long stalk, small decorative fruit and long stalk, deep red leaf and a small clumping habit (Sanewski, 2009 and de Souza *et al.*, 2009). In sunflower, the objective in improvement for commercial cut flowers are medium flower size and long shelf-life, while for pot plant is dwarf plant type.

Jatropha (*Jatropha curcas* L.) is a shrub from which its seed oil is normally used for biodiesel production (Kumar *et al.*, 2008). *Jatropha* also has a potential for ornamental purpose through interspecific cross with related species; especially with *J. integerrima*, an ornamental shrub, hard wood, frost tolerance and resistant to root rot diseases and leaf eating caterpillars (Lakshminarayana and Sujatha, 2001). Crossing among the two species is usually successful but the F₁ plants set only few seeds (Sujatha and Prabakaran, 2003; Parthiban *et al.*, 2009 and Dhillon *et al.*, 2009) and thus it is difficult to advance the selection process to the next generation. In this study, we produced populations of F₁, F₂ and BC₁F₁ from a cross of *J. curcas* and *J. integerrima* and selected individual plants based on flower color and plant type for ornamental purpose.

MATERIALS AND METHODS

Jatropha or physic nut (*J. curcas*) and dwarf peregrina (*J. integerrima*) were propagated by cuttings and sown in plastic bags filled with

Accredited SK No.: 81/DIKTI/Kep/2011

<http://dx.doi.org/10.17503/Agrivita-2014-36-3-229-234>

commercial soil during the rainy season of 2009. At 3 months old, the seedlings were transplanted to a crossing block in the field of Agronomy Department, Kasetsart University, Kamphaeng Saen, Nakhon Pathom, Thailand; using a spacing of 1m × 1.5m. Six months after transplanting, the plants were flowering and thus pollinated at various time durations in both direct (*J. curcas* as female) and reciprocal crosses.

J. curcas has green corollas in both male and female flowers setting on the same florescence of raceme type, while *J. integerrima* has deep red corollas in both male and female flowers. The male flowers of each species were emasculated before anthesis and covered with pollinator-proof bag. Pollination was attempted in five time durations, viz. 06:00 – 07:00, 08:00 – 09:00, 10:00 – 11:00, 12:00 – 13:00, 14:00 – 15:00, and 16:00 – 17:00 o'clock. The pollen grain of each species was collected at the time of anthesis and dusted on female flowers of the opposite species. In each time duration, pollination was performed on 4 inflorescences, each treated as a replicate. Each inflorescence was trimmed leaving only 5 female flowers. The same practice was done for both direct and reciprocal crosses. Fruit setting rate and mature seed number per fruit were recorded. The parental and hybrid seeds were sown in plastic bags filled with commercial soil and kept in a nursery at 50% sun light intensity for 2 months, then transplanted to the field. At flowering, plants showing prolific flowers with distinct colors were chosen and grafted onto jatropha stocks grown in pots of 30 cm in diameter. The pots were filled with commercial soil and kept under the same nursery during December 2011 to August 2012. Horticultural characters were observed on the F₁ and parental plants at the age of around 8 months old, which was the common blooming time of both species. Each plant was recorded for sizes of canopy and leaf, as well as characters of raceme and flower. Once the *F*-test indicated a difference among the genotypes, their means were compared using Least Significant Difference (LSD). The number of F₂ and BC₁F₁ plants segregating in different flower colors were also recorded.

RESULTS AND DISCUSSION

The suitable duration for interspecific pollination between *J. curcas* and *J. integerrima*

was 08:00–13:00 o'clock (Table 1). Crossing was successful only when *J. curcas* was used as the female parent, while the reciprocal cross failed to set fruit. This can be explained based on the flower behaviors of both species. Their female flowers are fully blooming in the morning around 7:30-8:00 AM, while pollen from the male flowers is shedding around 8:00-8:30 AM and gradually wilting in afternoon of the same day. Thus pollinations are more successful in the morning.

Table 1. Mean and percentage of fruits set from interspecific hybridization between *J. curcas* and *J. integerrima* at different time durations. Crossing was successful only when *J. curcas* was used as the female parent

Duration (o'clock)	Mean fruits set ±SD (%) ^a
06.00 - 07.00	5.50 ± 1.36 (27.50)
08.00 - 09.00	14.50 ± 1.16 (72.50)
10.00 - 11.00	15.75 ± 1.28 (78.75)
12.00 - 13.00	13.00 ± 1.16 (65.00)
14.00 - 15.00	8.50 ± 1.32 (42.50)
16.00 - 17.00	4.50 ± 1.92 (22.50)
F-test	**
LSD	2.97
CV (%)	27.15

Remarks: ** Significant at P ≤ 0.01, ^a average form four replications

Interspecific hybrid seeds were obtained only when *J. curcas* was used as the receiver. Our result corresponds with the earlier report of Sujatha and Prabakaran (2003). They observed that when *J. integerima* was the female parent, although the pollen tube of *J. curcas* germinated entering into the ovule, there was no fertilization, however. At 8 months after transplanting, the F₁ plants were different from their parents in all morphological characters (Table 2).

The F₁ plants showed exceptionally high heterosis in number of inflorescences per plant, giving the average number of more than 27 inflorescences per plant. Although the number of flowers per inflorescence of the F₁ hybrid fell between both plants, the total number of flowers were many more than the high parent (*J. integerrima*). Flower colors in the F₁ segregated into five white-flowered plants and four pink-flowered plants.

Table 2. Mean \pm SD of horticultural characters of F₁ hybrid and their parents at 8 months old

Genotypes	Canopy height (cm)	Canopy width (cm)	Leaf length (cm)	Leaf width (cm)	Length of flower stalk (cm)	No. inflorescences /plant	No. female flowers/infl orescence	No. male flowers/ inflorescence
<i>J. curcas</i>	189.44 \pm 19.03	80.67 \pm 9.49	13.69 \pm 0.66	12.38 \pm 0.27	12.86 \pm 0.36	6.67 \pm 2.22	11.12 \pm 2.12	114.51 \pm 12.93
<i>J. integerrima</i>	48.78 \pm 7.69	31.00 \pm 3.45	4.83 \pm 0.33	4.54 \pm 0.13	16.08 \pm 0.46	14.44 \pm 1.99	4.87 \pm 0.53	30.09 \pm 6.24
F ₁ hybrid	127.78 \pm 8.36	45.56 \pm 7.53	11.62 \pm 0.16	11.34 \pm 0.23	13.92 \pm 0.70	27.56 \pm 5.64	7.70 \pm 0.76	69.92 \pm 14.53
F-test	**	**	**	**	**	**	**	**
LSD	15.26	8.99	0.45	0.25	0.62	4.57	1.59	14.05
CV (%)	9.49	12.63	3.92	2.06	3.28	21.37	15.32	14.91

Remarks: ** Significantly different between genotypes at P \leq 0.01



Figure 1. Variation of flower colors in an F₂ population derived from selfing of a pink F₁ plant: A) green, B) light green, C) white, D) very light pink, E) light pink, F) pink, G) deep pink, H) very deep pink, and I) red

The F₂ plants obtained from selfing of an F₁ plant with pink flowers segregated into nine color patterns (Figure 1), viz. 39 green : 83 light green : 88 white : 65 very light pink : 48 light pink : 28 pink : 21 deep pink : 12 very deep pink : 7 red-flowered plants. The backcross progenies of *J. curcas* and a pink-flowered F₁ segregated into 22 green : 28 light green : 31 white : 20 very light pink : 11 light pink. While backcross between *J. curcas* and a white-flowered F₁ plant segregated into 27 green : 22 light green : 19 white (Figure 2).

The number of plants segregated for flower colors in this study was similar to those reported by Sujatha and Prabakaran (2003). They observed flower colors in a BC₁F₁ population derived from interspecific cross between *J. curcas* and *J. integerrima* and postulated that the colors were controlled by three loci of genes; *G*, *R* and *r* for green and red patterns, and white flowers, respectively. They explained that gene at locus *G* was inherited from *J. curcas*, while *R* and *r* were from *J. integerrima*. There were 2 or more alleles with both additive

and dominance effects. Unfortunately, our segregating populations did not show any possible Mendelian segregation ratios. Thus we assumed that flower colors in jatropha is quantitatively inherited (see also Fig. 2), and the darker colors are resulted from accumulation of more alleles controlling red pigmentation. Yet, there are also other important characters to be considered in selection for ornamental jatropha such as plant type, number of flowers per inflorescence, number of inflorescences per plant, leaf shape, propagation potential and disease resistance.

We finally selected 6 ornamental jatropha plants from the F₂ population based on their unique flower colors and high number of flowers per plant. They were registered at Kasetsart University and the Thai Department of Agriculture under the names 'Kamphaeng Saen 1' to 'Kamphaeng Saen 6'. The cultivars were made available to the public since early 2013. All varieties can be easily propagated by cutting or grafting (Muakrong et al., 2014).




Figure 2. Histograms of F_2 and BC_1F_1 plants segregated in different flower colors. (A) F_2 derived from pink-flowered F_1 plant, (B) BC_1F_1 from *J. curcas* and pink-flowered F_1 and (C) BC_1F_1 from *J. curcas* and white-flowered F_1

CONCLUSIONS AND SUGGESTIONS

Interspecific hybridization between *J. curcas* and *J. integerrima* was successful only when *J. curcas* was used as female parent, with the suitable crossing duration spans from late morning to early afternoon. Their progenies showed high variation in flower colors, particularly in the F₂ population which could be classified into nine color groups. To maximize this genetic variation, jatropha breeding for ornamental purpose should be conducted the in F₂ population as in improvement of most other plants. The selected plants can be vegetatively propagated as clones for further tests as well as for commercial purpose.

ACKNOWLEDGEMENT

We thank Office of the Higher Education Commission of Thailand for providing the senior author with the Strategic Scholarships Fellowships Frontier Research Networks (specific for southern region of Thailand). A part of this research was supported by the Chair Professor Project (project no. P-11-00599) of Thailand's National Science and Technology Development Agency.

REFERENCES

- Arslan, M., 2012. Evolution of genetic similarity and agronomic traits of castor bean populations naturally grown in the eastern Mediterranean region of Turkey. *Biotechnol. & Biotechnol. Eq.* 26:3089–3093.
- Bradley, V. L., R. L. Guenther, R. C. Johnson and R. M. Hannan. 1999. Evaluation of saf-flower germplasm for ornamental use. ASHS Press, Alexandria, USA. 433–435 pp.
- de Souza, F.V., J.R.S. Cabral, E.H. de Souza, O.S.N. Santos and F.R. Ferreira. 2009. Evaluation of F₁ hybrids between *Ananas comosus* var. ananassoides and *Ananas comosus* var. erectifolius. *Acta Hort* 822:79–84.
- Dhillon, R.S., M.S. Hooda, M. Jattan, V. Chawla, M. Bhardwaj and S.C. Goyal. 2009. Development and molecular characterization of interspecific hybrids of *Jatropha curcas* × *J. integerrima*. *IJBT.* 8:384–390.
- Golkar, P., A. Arzani, and A. M. Rezaei. 2010. Inheritance of flower colour and spinelessness in safflower (*Carthamus tinctorius* L.). *J. Genet.* 89:259–262
- Kumar, A. and S. Sharma. 2008. An evaluation of multipurpose oil seed crop for industrial uses (*Jatropha curcas* L.): a review. *Ind Crops Prod* 28:1–10.
- Muakrong, N., P. Tanya and P. Srinives. 2014. 'Kamphaeng Saen 1', 'Kamphaeng Saen 2', and 'Kamphaeng Saen 3': new ornamental *Jatropha* cultivars derived through an interspecific cross. *HortScience.* 49:1083-1085.
- Lakshminarayana, M. and M. Sujatha. 2001. Screening of *Jatropha* species against the major defoliators of castor (*Ricinus communis* L.). *J. Oilseeds Res.* 18:228–230.
- Parthiban, K.T., S. Kumar, R. Thiyagarajan, P. Subbulakshmi, V. Vennila and S. Govinda. 2009. Hybrid progenies in *Jatropha* – a new development. *Curr Sci India.* 96:15–23.
- Sanewski, G.M. 2009. Breeding Ananas for the cut flower and garden markets. *Acta Hort.* 822:71–78.
- Sujatha, M. and A.J. Prabakaran. 2003. New ornamental *Jatropha* hybrids through interspecific hybridization. *Genet Resour Crop Evol.* 50:75–80.