

THE MINIMUM NUMBER OF VALVES FOR DIATOM IDENTIFICATION IN RAWAPENING LAKE, CENTRAL JAVA

TRI RETNANINGSIH SOEPROBOWATI^{1,2*}, S. DJALAL TANDJUNG³, SUTIKNO⁴,
SUWARNO HADISUSANTO³, and PETER GELL⁵

¹*Department of Biology, Faculty of Sciences and Mathematics, Universitas Diponegoro, Semarang 50275, Indonesia*

²*School of Postgraduate Studies, Universitas Diponegoro, Semarang 50275, Indonesia*

³*Faculty of Biology, Universitas Gadjah Mada, Yogyakarta 55281, Indonesia*

⁴*Faculty of Geography, Universitas Gadjah Mada, Yogyakarta 55281, Indonesia*

⁵*Water Research Network, Federation University Australia, Ballarat, Australia*

Received: 21 April 2015/Accepted: 30 June 2016

ABSTRACT

Technical challenges in using diatoms for paleolimnological work are the identification and enumeration of diatom valves. Variations exist in the minimum number of valves to identify, ranging from 100 to 700 valves of the dominant species. This task can be very time consuming, particularly when the diatom valves are not abundant. This research was conducted to determine the minimum number of valves to be identified in the diatom assemblages from Rawapening Lake, Central Java, Indonesia. Based on the 314 samples obtained from Rawapening Lake, the diatom efficiency rose above 0.85 upon the minimum count of 300 valves. The number of diatom species identified remained stable after the minimum of 300 valves. Therefore, the minimum number of diatom valves identified to represent the assemblage for paleolimnological analysis was 300.

Keywords: Diatom analysis, eutrophic, Indonesia, paleolimnology, Rawapening

INTRODUCTION

Diatoms are the common name of microalgae belong to Bacillariophyte that have wide distribution from the ocean, freshwater to humid land and part of aquatic food webs. The uniquely ornamentated silica cell wall remain undisturbed whenever fossilized. That is why diatoms are good tool for paleolimnology.

Diatom undoubtedly have potential as bioindicators for water quality changes due to their well preserved walls in sediments. Being primary producers, diatoms play significant roles in food webs. Diatoms are distributed worldwide in saline or freshwaters, have short life cycles and are responsive to environmental changes. Diatoms can live across ecological gradients, are easily sampled and can be analyzed at low cost (Dixit *et al.* 1992; Gell *et al.* 2007; Reid & Ogden 2009).

Diatoms community are responsive to environmental changes such as pH, water depth, nutrients, salinity, and also the current condition of the environment. Diatom assemblages are often specific to particular habitats, therefore diatom fossils can be used to characterize those habitats. For that reason, diatoms have been widely used to investigate the status of lakes and can explain about pollution control, water quality monitoring programs and the paleoecology of the lakes. Palaeolimnological studies offer an opportunity to understand the past environmental conditions (Bere 2014). Paleolimnological techniques can reveal long-term perturbations and transitions of lake ecosystems (Kattal *et al.* 2016).

To understand limnological change, diatom valves are often well preserved in sediments allowing the inference of water quality over time. However, Indonesia does not have standard methods for diatom analysis. Even in the temperate region countries, the protocol for identifying the number of valves differs, which may be not suitable if implemented in Indonesia.

* Corresponding author: trsoeprobowati@live.undip.ac.id

The analysis of diatoms consists of three steps, i.e. digestion to separate diatom valves from the sediment; sample mounting-slide preparation; and identification-enumeration. The number of valves identified in the enumeration step differs across regions. Battarbee (1986), identified 300 – 600 diatom valves in England, and many studies used Battarbee's standard. However, Bate and Newall (2002) suggested 200 valves as an adequate number to characterize the diatom assemblages. These studies are focused mainly on temperate systems. Tropical systems are very diverse which may influence diatom assemblages. Therefore, it is important to determine an adequate count size in tropical diatom assemblages to optimize the efficacy of the research effort. The lower standard number i.e. 100 valves (Round 1993) when being implemented in Indonesia, was still time consuming due to species diversity despite lower population size (Soeprbowati *et al.* 2005).

The objective of this research was to determine the minimum valves count in the identification-enumeration step of diatom analysis for paleolimnological studies.

MATERIALS AND METHODS

Rawapening Lake was chosen as a study site because the lake is small with eutrophication problems similar to other Indonesian lakes. Rawapening Lake is one of the 15 Indonesian national priority lakes in 2010-2014 (ME 2010). Rawapening Lake was chosen as a pilot project for 'Save Indonesian Lake' as it sufficiently represents the eutrophication and sedimentation problems of Indonesian lakes (ME 2011). Radical action is required to overcome lake degradation problems comprising, one clear program for action, substantial funding, and strong institutional collaboration (Soeprbowati 2015a). Paleolimnological study was among the priority programs, named Gerakan Penyelamatan Danau (Germadan) Rawapening, to save Rawapening Lake (ME 2011).

Rawapening Lake is located at 45 km south of Semarang and about 9 km east of Salatiga. Rawapening is surrounded by five volcanoes i.e. Telomojo (1,895 m asl), Butak (1,000 m asl), Balak (700 m asl), Payung (600 m asl), and Rong (600 m asl). There are four districts around the lake i.e.:

Tuntang, Bawen, Ambarawa and Banyubiru. About 17 villages are situated around the lake side and their agricultural areas are frequently subjected to flooding. Rawapening Lake, situated at about 400 m asl, is about 4 km long and 2.5 km wide, with slightly sloped (7%) sides. In the 1970s its maximum capacity was 65 million m³ and the minimum was 25 million m³. However, there has been a clear trend of reducing volume, which has affected its capacity to generate hydropower electricity (Soeprbowati *et al.* 2012b).

Echounding of the lake revealed extensive areas of less than 2 m depth; there were also three deeper depocentres in the lake's west, each 18 m deep (Soeprbowati 2012). Sediment cores of different lengths were collected from four research sites across Rawapening Lake. The longest sediment core (63 cm, *As*) was obtained from the Asinan site. The Panjang (*Pj*) and Tuntang (*Tg*) cores were 36.5 cm and 35.5 cm long, respectively. The Dangkel (*Dk*) core was 29 cm (Fig. 1).

Tg site was represent an outlet of Rawapening Lake, *As* site was near an inlet that come from a settlement and so represents a settlement catchment area and *Pj* site was near an inlet passing through agricultural area, thereby representing an agricultural catchment area. *Dk* site was located relatively close to the middle of the lake and so represents the lake body. Those 4 sites were determined to sufficiently represent Rawapening Lake. Diatoms were sliced every 0.5 cm based on the modified version of Battarbee *et al.* (2001). The first step of extraction is intended to separate diatom valves from organic material. Basically, depending on the type of sediment, strong acid may be used to digest sediment. This study applied 10% HCl followed by 10% H₂O₂ to digest sediment samples. In the preparation process, a mountant with refraction index of 1.7 is required. Hyrax was used in this research and the silicious striae were clearly seen under the microscope, expediting the identification process. Identification of diatom species was carried out by referencing the diatom samples to the standard texts (Kramer & Lange-Bertalot 2004a, 2004b, 2004c, 2010) and by referencing the samples to the diatom collections held at Universitas Diponegoro, Semarang, Indonesia and at the Federation University Australia, Ballarat, Australia. A total of 600 valves were counted for each sample with totals tallied at steps of 100 (i.e.




Figure 1 Study sites in Rawapening Lake for collecting sediment core samples

Notes: *As* = Asinan Site (63 cm) represents settlement catchment area, *Pj* = Panjang Site (35.5 cm) represents agricultural catchment area, *Tg* = Tuntang Site (36.5 cm) represents outlet of Rawapening, *Dk* = Dangkel Site (29 cm) represents the lake body

100, 200, 300, 400, 500, 600). Species accumulation curves were applied to identify the minimal number of valves to be counted to achieve maximum efficiency, which was calculated using the formula (Bates & Newall 2002):

$$\text{Maximum efficiency} = 1 - \frac{\text{number of species}}{\text{number of individuals}}$$

Maximum efficiency reflected the probability of new species to be found at each identification

step (Pappas & Stoermer 1996; Bates & Newall 2002).

RESULTS AND DISCUSSION

Three steps commonly used in diatom analysis consist of extraction, preparation, and identification. In the identification process, the minimal number of valves identified varied among researchers. Counting the valves of




Figure 2 Maximum efficiency counts of diatoms from the upper 1 cm of sediment cores from 4 study sites

Notes: *As* = Asinan Site (63 cm) represents settlement catchment area, *Pj* = Panjang Site (35.5 cm) represents agricultural catchment area, *Tg* = Tuntang Site (36.5 cm) represents outlet of Rawapening, *Dk* = Dangkel Site (29 cm) represents the lake body

diatoms is important to provide data about the ecological condition in the past.

This study analyzed 314 samples, compared valve counts at 100, 200, 300, 400, 500, and 600 valves, found that the uppermost (1 cm) sediment samples from each of four sites. The counting results showed maximum efficiency of 0.87–0.93 (Fig. 2). The number of diatom species identified remained stable after 300 valves while the maximum efficiency rose to above 0.85. New finding from this research is that 300 valves is the most efficient number of valves for paleolimnological assessment.

The number of diatom species identified from 4 sites were different. The highest number of diatom species was found in *Dk* site having 42 species in the counts of 200 valves which remained stable in the counts of 300 through to 600 valves (Fig. 3). *Dk* was situated close to the middle of the lake, having a water depth of 2 m with sediment of peat mud. This might be correlated with numerous *Ennotia* species found only in the *Dk* site indicating its acid condition (Soeprbowati *et al.* 2012b). Typically, when the euphotic depth is shallow, the number of aquatic plants attached to the substrate decline, causing nutrients and sediments to be further released into the water column. This drives the increasing numbers of phytoplankton to continually exploit

the light that is confined to the surface waters (Reid *et al.* 2007). This condition is shown in the diatom record of the Rawapening Lake sediments that are dominated by planktonic forms thriving in eutrophic, turbid and alkaline waters. The odd situation happened in *Tg* site. Thirty diatom species were found in the count of 200–600 valves. The R^2 of 0.42 for *Tg* site indicated that there was a low correlation between the number of species with the number of valves counted. In *Tg* site, large diatom species might adapt well to the moving outlet water. *As* site showed 22 diatom species in 400 valves and remained stable. *Pj* site had the lowest number of species (14 species) in the count of 200 valves and increased to 17 species in the count of 500 valves.

Statistically, comparing the results of analysis of variance, the diatom species numbers found in each sediment layer were significantly different between counts of 100, 200, 300, 400, 500, and 600. Further Least Significant Difference (LSD) analysis showed that results from the count of 500 valves were significantly different from the count of 400 and 600 valves. There was also significant difference between the diatom species number for the count of 400 and 600 valves. The increasing numbers of diatom species in the count of 600 valves were probably due to contaminant species. This was based on the fact




Figure 3 Number of diatom species identified from the upper 1 cm of sediment cores from 4 research sites

Notes: *As* = Asinan Site (63 cm) represents settlement catchment area, *Pj* = Panjang Site (35.5 cm) represents agricultural catchment area, *Tg* = Tuntang Site (36.5 cm) represents outlet of Rawapening, *Dk* = Dangkel Site (29 cm) represents the lake body

that there had been no or low increase in the numbers of diatom species with low populations from the 4 sites of Rawapening Lake (Fig. 3). The maximum efficiency of those 4 sites also indicated that the count of 400 – 600 valves had maximum efficiency above 0.9 (Fig. 2). Analysis of variance supported this result, there is no significant difference. Hence, the minimum count of 300 valves having efficiency more than 0.85, was appropriate for the paleolimnological analysis of a tropical lake, with specific reference to the eutrophic Rawapening Lake.

In England, Battarbee *et al.* (1986; 2001) proposed counting 300 - 600 valves for routine analysis, but Round (1993) determined that 100 valves were sufficient for dominant species. In Sweden, Gothe *et al.* (2013) identified diatoms until they found at least 400 valves. In Finland, Soininen & Kononen (2004) identified 250 – 500 valves. In France, Morin *et al.* (2008) identified 300 valves. In Australia Fluin *et al.* (2010) identified diatoms until they found between 300 – 540 valves, whereas Grundell *et al.* (2012) identified 200 valves. In Canada, Koster *et al.* (2005) and Pienitz *et al.* (2006) identified as many as 500 valves. In Mexico, a minimum of 500 valves were identified (Siqueiros-Beltrones *et al.* 2005). In Uganda, Mills (2009) counted until 300 - 500

valves were found. In America, Kireta *et al.* (2012) identified 100 valves in samples with sparse diatoms. In India, it is recommended to count 400 valves (Karthick *et al.* 2010). For fossil diatoms, the count should be different, because some diatom species may have been dissolved over a period of time or diatom samples may contain diatom species from previous period of time.

In Indonesia, the minimum number of valves recommended by Round (100 valves) had been implemented, but this is less efficient and less effective, particularly given Round (1993) stipulation that 100 valves of the dominant species had to be counted (Soeprbowati *et al.* 2005, 2012a). In this study, there were significant differences between the counts of 100, and 200 and 300, while the count of 500 valves was not significantly different from 400 and 600. These results were similar to Battarbee's (1986) statement that there were marked differences between the count of 100 and 200 valves, while there was little difference between the count of 400 and 500. For this reason, he recommended that a count of 300 to 600 may be used for routine analysis purposes. Based on this research, it was recommended that a count of a minimum 300 valves might be used for paleolimnological analysis in Rawapening Lake.

The counting of diatom valves is to produce a semi-quantitative approach for ecological analysis. Therefore, it is very important to determine the minimum number of valves to be counted to get a reliable approach to gauge the relative species composition at sampling sites (Karthick *et al.* 2010). The minimum total number of valves to be counted for each sample varies depending on the purpose of the assessment and the need to produce statistically sound results. Comparing the count of 200 and 600 valves, there were percentage differences of 1.89. The count of 300 and 600 valves provided 1.85 percentage differences which suggested that the count of 300 valves was sufficient for the calculation of diatom species.

Results of comparing the maximum efficiency of counting 100, 200, 300, 400, 500, and 600 valves showed that the minimum number of valves that should be identified was 300, since its maximum efficiency was more than 0.85. Maximum efficiency is considered to sufficiently represent diatom species numbers because the formula to calculate maximum efficiency includes the number of individual valves. Therefore, maximum efficiency can be used to determine the minimum valves to be identified in diatom analysis (Bates & Newall 2002).

In Australia, the minimum number of 200 valves had more than 80% efficiency, and this number was deemed suitable to be used in water quality monitoring programs (Bates & Newall 2002). There was no influence on the diatom index at counts of valves 300 or above (Prygiel *et al.* 2002).

A European Protocol for diatom enumeration, DALES (*Diatoms for Assessing Lake Ecological Status*, version 1.0 2004), determined that at least 300 valves should be identified, especially for non planktonic taxa. When the abundance of one taxon was more than one third of all individual valves, the protocol recommended to increase the sample size until a minimum of 200 valves of non planktonic diatom are found. New species found in the count above 400 valves were determined to be contaminant species. The presence of these species may cause bias for further analysis. Therefore, species found to be less than 5% of valves were considered to be rare species and were not included in the data analysis. Dominant species are considered to provide more evidence of the water quality than the rare species.

Based on the diatom efficiency and diatom species found from Rawapening Lake, a minimum count of 300 valves was sufficient for paleolimnological analysis, which is lower than the recommended valves count for temperate or polar areas. This recommendation might be related to the year round warm temperature in the tropics which increases the diatom species diversity.

A minimum count of 300 diatom valves was implemented to reconstruct the environmental condition of Rawapening Lake since the 1960s. The dominance of *Fragilaria capucina* Desm., *Luticola goeppertiana* (Bleisch) Mann, *Mayamaea atomus* (Kutzing) Lange-Bertalot, *Navicula radiosa* Kutzing, *Nitzschia palea* (Kutzing) W. Smith and in As site, *Tryblionella apiculata* Gregory, reflected eutrophic, but clear waters in 1967-1974. The presence of *Eunotia pectinalis* (Kutzing) Rabenhorst var. *undulata* (Ralfs) Rabh suggested neutral to slightly acid conditions in 1967-1974 and the appearance of *Fragilaria capucina* Desm., *Gomphonema gracilis* Ehr. and *Navicula radiosa* Kutzing suggested changes of water pH to alkaline conditions. An increase in epiphytic *Gomphonema* spp. in 1974-1983 marked an increase in aquatic macrophyte plants, perhaps in response to high nutrient levels. This change was followed promptly by the increasing numbers of acidophilous *Eunotia* spp. reflecting high organic production. A transition to a diatom community dominated by planktonic forms occurred around 1983. This community was initially dominated by more clear water, oligotrophic species such as *Discostella stelligera* (Cleve and Grunow) Houk and Klee and *Aulacoseira distans* (Ehrenberg) Simonsen, but transitions happened in 1990 to be dominated by *A. granulata* (Ehrenberg) Simonsen and ultimately *Aulacoseira ambigua* (Grunow) Simonsen. This was interpreted as a shift to a turbid water phase that contained beneficial phytoplankton, at the expense of benthic or epiphytic taxa requiring clear water. The dominance of *A. granulata* (Ehrenberg) Simonsen since the 1990s indicated that the lake experienced hypertrophic conditions with pH>9. Although a high proportion of the taxa in Rawapening Lake sediments were not represented in the European data set, Rawapening Lake experienced hypertrophic condition with pH>9 as indicated by the dominance of *Aulacoseira granulata* (Ehrenberg) Simonsen (Soeprbowati *et al.* 2012b).

However, when the identification was done using counts of less than 300 valves, some predominant species such as *Aulacoseira ambigua*, *Cyclotella meneghiniana*, *Gomphonema gracillius*, *Synedra ulna* were not found. The predominance of *Synedra* from 1967 to the present indicated that Rawapening Lake had been fresh and meso-eutrophic throughout. *Synedra ulna* (Nitzsch) Ehrenberg is a tolerant species, found in Indonesian rivers and lakes with high organic content with total phosphorous content of 20 - 1,000 µg/L and pH of 5 – 9. The modern sampling of pH at Rawapening Lake revealed that pH in Rawapening Lake fluctuated. Goltenboth (1994) reported that the pH of Rawapening Lake was 7.4 ± 0.2 (dry season) and 7.3 ± 0.1 (wet season). EPA-ERC Undip (1999) reported that the pH of Rawapening Lake was 7.96 ± 0.42 . In 2004 and 2005, the pH of the inlet and lake tended to be neutral (7.04 ± 1.13), except in the site around the spring and floating island where the pH was 9.52 (Soeprbowati *et al.* 2005). During field work for this study (2008), pH increased up to 9.39 ± 2.51 . In recent study, those paleoreconstruction of ecological change in Rawapening proved a trend of increasing pH. Measurements of pH in June 2015 showed that pH of 14 sites in Rawapening Lake was 7.22 ± 2.45 (Soeprbowati 2015b).

CONCLUSIONS

Diatom maximum efficiency rose to above 0.85 at the minimum count of 300 valves. The number of diatom species identified remained stable after the minimum count of 300 valves. A minimum count of 300 valves was appropriate for the paleolimnological analysis of a tropical lake, with specific reference to the eutrophic Rawapening Lake.

ACKNOWLEDGEMENTS

Some parts of this article was produced as a part of project supported by Indonesian Higher Education through a Fundamental Research Grant to Universitas Diponegoro Number: 0160.0/023-04.2/XIII/2009, as stated in Rector Letter Number 180/SK/H7/18 March 2009 and contract of multiyears research number:

124A/H7.2/KP/18 March 2009. The recent survey of pH was supported by Competence Research Grant which is supported by the Directorate of Research and Community Services), Directorate General of Higher Education, the Ministry of Education and Culture, Year 2014, through DIPA UNIP Number: 023.04.2.189185/2014, 03 Maret 2014.

Thanks to Keely Mills and Rosie Grundell from the Federation University Australia, Ballarat, Australia for their assistance in the diatom analysis. Special thanks to Jafron Wasiq Hidayat and Kariyadi Baskoro for fieldworks assistance. Special thanks to Sri Widodo Agung Suedy from Universitas Diponegoro, Semarang, Indonesia for his permission in using pH data from Competence Research Grant. Deep gratitude goes to Nina Desianti from Laboratory of the Academy of Natural Sciences, Drexel University, USA for her assistance in data analysis.

REFERENCES

- Bates N, Newall P. 2002. Techniques for the use of diatoms in water quality assessment: how many valves?. In John J, editor. Proceeding of 15th International Diatom Symposium. p.153-60.
- Battarbee RW. 1986. Diatom analysis. In: Berglund, editor. Handbook of Holocene Paleocology and Paleohydrology. London (UK) John Wiley&Sons. p:527-70.
- Battarbee R, Jones VJ, Flower RJ, Cameron NG, Bennion H, Carvalho L, Juggins S. 2001. Diatoms. In: Smol, JP, Birks, HJB, Last, WM, editors. Tracking Environmental Change using Lake Sediments. Volume 3: Terrestrial, Algal and Silicious Indicators. Kluwer Academic Publishers. p.155-202.
- Bere T. 2014. Ecological preferences of benthic diatoms in a tropical river system in São Carlos-SP, Brazil. *Tropical Ecology* 55(1): 47-61.
- DALES (Diatoms for Assessing Lake Ecological Status). 2004. Enumeration Protocol, ver 1.0. <http://craticula.ncl.ac.uk/DALES/downloads> 1 April 2016.
- Dixit SS, Smol JP, Kingston JC, Charles DF. 1992. Diatoms: powerful indicators of environmental change. *Environ Sci Technol* 26(1):23-32. <http://pubs.acs.org/doi/abs/10.1021/es00025a002>.
- EPA-ERC Undip. 1999. Environmental Quality Index Development and Biological Indicator. 1st Ed. Research Institute Diponegoro University, Semarang, pp: 81.

- Fluin J, Tibby J, Gell, P. 2010. The palaeolimnological record from Lake Cullulleraine, Lower Murray River (south-east Australia): implications for understanding riverine histories. *J Paleolimnol* 43(2):309-22. <http://dx.doi.org/10.1007/s10933-009-9333-8>.
- Gell P, Tibby J, Little F, Baldwin D, Hancock G. 2007. The impact of regulation and salinisation on floodplain lakes: The lower River Murray, Australia. *Hydrobiologia* 591:135-46. <http://dx.doi.org/10.1007/s10750-007-0806-3>.
- Goltenboth F, Timotius KH. 1994. Danau Rawa Pening di Jawa Tengah, Indonesia (Rawapening Lake, Central Java, Indonesia). Satya Wacana University Press, Salatiga.
- Gothé E, Angler DG, Gottschalk S, Lofge S, Sandin L. 2013. The Influence of Environmental, Biotic and Spatial Factors on Diatom Metacommunity Structure in Swedish Headwater Streams *PLoS ONE* 8(8): e72237. doi:10.1371/journal.pone.0072237.
- Grundell R, Gell P, Mills K, Zawadzki A. 2012. Interaction between a river and its wetland: evidence from the Murray River for spatial variability in diatom and radioisotope records. *J Paleolimnol* 47:205-19. <http://link.springer.com/article/10.1007/s10933-011-9572-3>.
- Kattel G, Gell P, Zawadzki A, Barry L. 2016. Palaeoecological evidence for sustained change in a shallow Murray River (Australia) floodplain lake: regime shift or press response? *Hydrobiologia* · September 2016 DOI: 10.1007/s10750-016-2970-9.
- Karthick B, Taylor JC, Mahesh MK, Ramachandra TV. 2010. Protocols for Collection, Preservation and Enumeration of Diatoms from Aquatic Habitats for Water Quality Monitoring in India. *The IUP Journal of Soil and Water Sciences*, 3(1):25-60.
- Kireta AR, Reavie ED, Sgro GV, Angradi TR, Bolgrien DW, Jicha TM, Hill BH. 2012. Assessing the condition of the Missouri, Ohio, and Upper Mississippi rivers (USA) using diatom-based indicators. *Hydrobiologia* 691:171–88. <http://link.springer.com/article/10.1007/s10750-012-1067-3#page-1>.
- Koster D, Pienitz R, Wolfe BB, Barry S, Foster DR, Dixit SS. 2005. Paleolimnological assessment of human-induced impacts on Walden Pond (Massachusetts, USA) using diatoms and stable isotopes. *Aquat Ecosys Health & Manage* 8(2):117-31. <http://www.tandfonline.com/doi/abs/10.1080/14634980590953743>.
- Kramer K, Lange-Bertalot Ha. 2004. *Susswasserflora Von Mitteleuropa*, Bd. 02/2: Bacillariophyceae: Teil 2: Bacillariophyceae, Epithemiaceae, Surirellaceae. Spectrum, Berlin.
- Kramer K, Lange-Bertalot Hb. 2004. *Susswasserflora Von Mitteleuropa*, Bd. 02/3: Bacillariophyceae: Teil3: Centrales, Fragillariaceae, Eunotiaceae. Spectrum, Berlin.
- Kramer K, Lange-Bertalot Hc. 2004. *Susswasserflora Von Mitteleuropa*, Bd. 02/4: Bacillariophyceae: Teil 4: Achnanthes S.L., Navicula Sstr. Spectrum, Berlin.
- Krammer K, Lange-Bertalot H. 2010. *Susswasserflora von Mitteleuropa*, Bd. 2/1: Bacillariophyceae, Teil 1: Naviculaceae. 1st Edn., Spektrum Akademischer Verlag, ISBN-10:3827426154, p: 882.
- ME (Ministry of Environment). 2010. 2010-2014 National Lake Priority Program. 1st Ed. The Indonesian Ministry of Environment, pp: 7.
- ME (Ministry of Environment). 2011. Gerakan Penyelamatan Danau (GERMADAN) Rawapening. 1st Ed. The Indonesian Ministry of Environment, pp: 93.
- Mills K. 2009. Ugandan Crater Lakes Limnology, Palaeolimnology and palaeoenvironmental history. Unpublished Ph.D., Loughborough University, England. <https://dspace.lboro.ac.uk/dspace-jspui/bitstream/2134/13219/3/Thesis-2009-Mills.pdf>.
- Morin S, Coste M, Delmas F. 2008. A comparison of specific growth rates of periphytic diatoms of varying cell size under laboratory and field conditions. *Hydrobiologia* 614:285-97. <http://link.springer.com/article/10.1007/s10750-008-9513-y>.
- Pappas JL, Stoermer EF. 1996. Quantitative method for determining a representative algae sample count. *Phycol* 32:693-6. <http://onlinelibrary.wiley.com/doi/10.1111/j.0022-3646.1996.00693.x/pdf>.
- Pienitz R; Robergem K, Vincent WF. 2006. Three hundred years of human-induced change in an urban lake: paleolimnological analysis of Lac Saint-Augustin, Quebec City, Canada. *Can J Bot* 84:303-20. <http://www.nrcresearchpress.com/doi/abs/10.1139/B05-152>.
- Prygiel J, Carpentier P, Almeida, Coste M, Druart JC, Ector L, ... Zydek N. 2002. Determination of the Diatom Index (IBD NF T 90-354): results of an intercalibration exercise. *J. App Phycol* 14, pp.27-39. <http://link.springer.com/article/10.1023/A:1015277207328>.
- Reid MA, Sayer CD, Kershaw AP, Heijnis H. 2007. Palaeolimnological evidence for submerged plant loss in a floodplain lake associated with accelerated catchment soil erosion (Murray River, Australia). *J. Paleolimnol* 38:191-208. <http://link.springer.com/article/10.1007/s10933-006-9067-9>.
- Reid MA, Ogden RW. 2009. Factors affecting diatom distribution in floodplain lakes of the Southeast Murray basin, Australia and implications for paleolimnological studies. *J Paleolimnol* 41:453-70. <http://link.springer.com/article/10.1007/s10933-008-9236-0>.

- Round FE. 1993. A review and methods for the use of epilithic diatoms for detecting and monitoring changes in river water quality. Methods for the examination of waters and associated materials. pp. 63. HMSO, London.
- Siqueiros-Beltrones, Lo´Pez-Fuerte FO, Ga´Rate-Liza´Raga, I. 2005. Structure of diatom assemblages living on prop roots of the red mangrove (*Rhizophora mangle*) from the west coast of Baja California Sur, Mexico. *Pacific Science* 59 (1):79–96. <https://muse.jhu.edu/article/176850/pdf>.
- Soininen J, Kononen K. 2004. Comparative study of monitoring South-Finnish Rivers and streams using diatom and macro invertebrate community structure. *Aquatic Ecol* 38:63-75. <http://link.springer.com/article/10.1023/B:AECO.0000021004.06965.bd#page-1>.
- Soeprbowati TR. 2012. Peta batimetri Danau Rawapening (Batimetric map of Rawapening). *J. Bioma* 14(2):78-84. <http://www.ejournal.undip.ac.id/index.php/bioma/article/view/9452/7616>.
- Soeprbowati TR. 2015a. Integrated Lake Basin Management for Save Indonesian Lake Movement. *Procedia Environ. Sci* 23(2015):368-74. <http://www.sciencedirect.com/science/article/pii/S1878029615000547>.
- Soeprbowati TR. 2015b. Bioindicator of water quality. Prosiding National Seminar of Biology, Program Magister Biologi Universitas Diponegoro, Semarang, 6 August 2015.
- Soeprbowati TR, Rahmanto WA, Hidayat JW, Baskoro K. 2005. Diatoms and present condition of Rawapening Lake. *Inter Sem Environ Chem Toxicol*, April 2005, INJECT Yogyakarta.
- Soeprbowati TR, Suedy SWA, Gell P. 2012a. Diatom stratigraphy of mangrove ecosystems on the Northern Coast of Central Java. *J Coast Dev* 15(2):197-208. <http://www.omicsonline.com/open-access/diatom-stratigraphy-of-mangrove-ecosystems-on-the-northern-coast-of-central-java-1410-5217-15-343.pdf>.
- Soeprbowati TR, Hadisusanto S, Gell P, Zawadski A. 2012b. The diatom stratigraphy of Rawapening Lake, implying eutrophication history. *Am J Environ Sci*(3): 334-344. <http://thescipub.com/PDF/ajessp.2012.334.344.pdf>.