



On d -antimagic labelings of plane graphs

Martin Bača^a, Ljiljana Branković^b, Marcela Lascsáková^a, Oudone Phanalasy^c,
Andrea Semaničová–Feňovčíková^a

^aDepartment of Applied Mathematics and Informatics, Technical University, Košice, Slovak Republic

^bSchool of Electrical Engineering and Computer Science, The University of Newcastle, Australia

^cDepartment of Mathematics, National University of Laos, Vientiane, Laos

martin.baca@tuke.sk, ljiljana.brankovic@newcastle.edu.au, marcela.lascsakova@tuke.sk,
oudone.phanalasy@newcastle.edu.au, andrea.fenovcikova@tuke.sk

Abstract

The paper deals with the problem of labeling the vertices and edges of a plane graph in such a way that the labels of the vertices and edges surrounding that face add up to a weight of that face. A labeling of a plane graph is called d -antimagic if for every positive integer s , the s -sided face weights form an arithmetic progression with a difference d . Such a labeling is called *super* if the smallest possible labels appear on the vertices.

In the paper we examine the existence of such labelings for several families of plane graphs.

Keywords: plane graph, d -antimagic labeling, super d -antimagic labeling

Mathematics Subject Classification : 05C78

1. Introduction

Let $G = (V, E, F)$ be a finite connected plane graph without loops and multiple edges, where V , E and F are its vertex set, edge set and face set, respectively. Let $|V(G)| = p$, $|E(G)| = q$ and $|F(G)| = r$ be the number of the vertices, the edges and the faces, respectively.

A labeling of type $(1, 1, 1)$ assigns labels from the set $\{1, 2, \dots, p+q+r\}$ to the vertices, edges and faces of a plane graph G in such a way that each vertex, edge and face receives exactly one label and each number is used exactly once as a label.

Received: 31 March 2011, Revised: 28 July 2011, Accepted: 21 November 2012.

A labeling of type $(1, 1, 0)$ is a bijection from the set $\{1, 2, \dots, p + q\}$ to the vertices and edges of a graph G .

The *weight of a face* under a labeling is the sum of labels (if present) carried by that face and the edges and vertices on its boundary.

A labeling of a plane graph G is called d -antimagic if for every positive integer s the set of s -sided face weights is $W_s = \{a_s, a_s + d, a_s + 2d, \dots, a_s + (r_s - 1)d\}$ for some integers a_s and $d \geq 0$, where r_s is the number of s -sided faces. We allow different sets W_s for different s .

If $d = 0$ then Ko-Wei Lih in [16] called such labeling *magic*. Ko-Wei Lih [16] described magic (0-antimagic) labelings of type $(1, 1, 0)$ for the wheels, the friendship graphs and the prisms. The magic labelings of type $(1, 1, 1)$ for the grid graphs and the honeycomb are given in [2] and [3], respectively.

The concept of the d -antimagic labeling of the plane graphs was defined in [10], where it was also proved that the prism D_n has d -antimagic labelings of type $(1, 1, 1)$ for $d \in \{2, 3, 4, 6\}$ and $n \equiv 3 \pmod{4}$. The d -antimagic labelings of type $(1, 1, 1)$ for the hexagonal planar maps, the generalized Petersen graph $P(n, 2)$ and the grids can be found in [5], [7] and [8], respectively. Lin *et al.* in [17] showed that prism D_n , $n \geq 3$, admits d -antimagic labelings of type $(1, 1, 1)$ for $d \in \{2, 4, 5, 6\}$. The d -antimagic labelings of type $(1, 1, 1)$ for D_n and for several $d \geq 7$ are described in [19].

A d -antimagic labeling is called *super* if the smallest possible labels appear on the vertices. The super d -antimagic labelings of type $(1, 1, 1)$ for antiprisms and for $d \in \{0, 1, 2, 3, 4, 5, 6\}$ are described in [4], and for disjoint union of prisms and for $d \in \{0, 1, 2, 3, 4, 5\}$ are given in [1]. The existence of super d -antimagic labelings of type $(1, 1, 1)$ for disconnected plane graphs and for plane graphs containing a special Hamilton path is examined in [6] and [12], respectively.

In this paper we examine the existence of super d -antimagic labelings of type $(1, 1, 0)$ for several families of plane graphs. To label the vertices and edges of plane graphs we will use an edge-antimagic vertex labeling and an edge-antimagic total labeling.

Simanjuntak, Bertault and Miller in [18] define an (a, d) -edge-antimagic vertex labeling of a (p, q) -graph $G = (V, E)$ as an injective mapping $\beta : V(G) \rightarrow \{1, 2, \dots, p\}$ such that the set of edge-weights $\{\beta(u) + \beta(v) : uv \in E(G)\}$ is $\{a, a + d, a + 2d, \dots, a + (q - 1)d\}$ for two non-negative integers a and d . A bijection $\alpha : V(G) \cup E(G) \rightarrow \{1, 2, \dots, p + q\}$ is called an (a, d) -edge-antimagic total labeling of G if the edge-weights $\{\alpha(u) + \alpha(uv) + \alpha(v) : uv \in E(G)\}$ form an arithmetic sequence starting at a and having a common difference d , where $a > 0$ and $d \geq 0$ are two fixed integers. An (a, d) -edge-antimagic total labeling is a natural extension of a notion of *magic valuation* defined by Kotzig and Rosa in [15].

An (a, d) -edge-antimagic total labeling is called *super* if the smallest possible labels appear on the vertices. A super (a, d) -edge-antimagic total labeling is a natural extension of a notion of *super edge-magic labeling* defined by Enomoto *et al.* in [13].

More comprehensive information on magic valuations and (a, d) -edge-antimagic total labelings can be found in [11], [14] and [20], respectively.

2. Edge-antimagic labelings of paths

Let P_n be the path on n vertices. It is known (see [9]), that P_n is super (a, d) -edge-antimagic total if and only if $d \leq 3$. We denote the vertices of P_n by v_1, v_2, \dots, v_n and describe these labelings $\alpha_d^n : V(P_n) \cup E(P_n) \rightarrow \{1, 2, \dots, 2n - 1\}$ in the following way.

a) The super $(2n + \lceil \frac{n}{2} \rceil + 1, 0)$ -edge-antimagic total labeling α_0^n of P_n :

$$\alpha_0^n(v_i) = \begin{cases} \frac{i+1}{2} & \text{for } i \equiv 1 \pmod{2} \text{ and } 1 \leq i \leq n, \\ \lceil \frac{n}{2} \rceil + \frac{i}{2} & \text{for } i \equiv 0 \pmod{2} \text{ and } 2 \leq i \leq n, \end{cases}$$

$$\alpha_0^n(v_i v_{i+1}) = 2n - i \quad \text{for } i = 1, 2, \dots, n - 1.$$

The common weight for all edges of P_n is

$$w_{\alpha_0^n}(v_i v_{i+1}) = 2n + \lceil \frac{n}{2} \rceil + 1 = C_{\alpha_0, 0}^n, \quad i = 1, 2, \dots, n - 1.$$

b) The super $(2n + 2, 1)$ -edge-antimagic total labeling α_1^n of P_n :

$$\alpha_1^n(v_i) = i \quad \text{for } i = 1, 2, \dots, n,$$

$$\alpha_1^n(v_i v_{i+1}) = 2n - i \quad \text{for } i = 1, 2, \dots, n - 1.$$

The set of edge-weights of P_n consists of the consecutive integers

$$\{w_{\alpha_1^n}(v_i v_{i+1}) = 2n + 1 + i = C_{\alpha_1, 1}^n + i : i = 1, 2, \dots, n - 1\}.$$

c) The super $(n + \lceil \frac{n}{2} \rceil + 3, 2)$ -edge-antimagic total labeling α_2^n of P_n :

$$\alpha_2^n(v_i) = \begin{cases} \frac{i+1}{2} & \text{for } i \equiv 1 \pmod{2}, \\ \lceil \frac{n}{2} \rceil + \frac{i}{2} & \text{for } i \equiv 0 \pmod{2}, \end{cases}$$

$$\alpha_2^n(v_i v_{i+1}) = n + i \quad \text{for } i = 1, 2, \dots, n - 1.$$

The edge-weights of P_n constitute the arithmetic progression of difference 2:

$$\{w_{\alpha_2^n}(v_i v_{i+1}) = n + \lceil \frac{n}{2} \rceil + 1 + 2i = C_{\alpha_2, 2}^n + 2i : i = 1, 2, \dots, n - 1\}.$$

d) The super $(n + 4, 3)$ -edge-antimagic total labeling α_3^n of P_n :

$$\alpha_3^n(v_i) = i \quad \text{for } i = 1, 2, \dots, n,$$

$$\alpha_3^n(v_i v_{i+1}) = n + i \quad \text{for } i = 1, 2, \dots, n - 1.$$

The edge-weights of P_n constitute the arithmetic progression of difference 3:

$$\{w_{\alpha_3^n}(v_i v_{i+1}) = n + 1 + 3i = C_{\alpha_3, 3}^n + 3i : i = 1, 2, \dots, n - 1\}.$$

Now, we define the super (a, d) -edge-antimagic total labelings of P_n also for negative differences d in the following way

$$\begin{aligned}\alpha_{-k}^n(v_i) &= \alpha_k^n(v_{n+1-i}) & \text{for } i = 1, 2, \dots, n, \\ \alpha_{-k}^n(v_i v_{i+1}) &= \alpha_k^n(v_{n+1-i} v_{n-i}) & \text{for } i = 1, 2, \dots, n-1,\end{aligned}$$

where $k = 0, 1, 2, 3$.

In this paper we will use also (a, d) -edge-antimagic vertex labelings of P_n for two differences $d = 1$ and $d = 2$. These labelings $\beta_d^n : V(P_n) \rightarrow \{1, 2, \dots, n\}$ we define in the following way:

e) The $(\lceil \frac{n}{2} \rceil + 2, 1)$ -edge-antimagic vertex labeling β_1^n of P_n :

$$\beta_1^n(v_i) = \begin{cases} \frac{i+1}{2} & \text{for } i \equiv 1 \pmod{2}, \\ \lceil \frac{n}{2} \rceil + \frac{i}{2} & \text{for } i \equiv 0 \pmod{2}. \end{cases}$$

The set of edge-weights of P_n consists of the consecutive integers

$$\{w_{\beta_1^n}(v_i v_{i+1}) = \lceil \frac{n}{2} \rceil + 1 + i = C_{\beta,1}^n + i : i = 1, 2, \dots, n-1\}.$$

f) The $(3, 2)$ -edge-antimagic vertex labeling β_2^n of P_n :

$$\beta_2^n(v_i) = i \quad \text{for } i = 1, 2, \dots, n.$$

The edge-weights of P_n constitute the arithmetic progression of difference 2:

$$\{w_{\beta_2^n}(v_i v_{i+1}) = 1 + 2i = C_{\beta,2}^n + 2i : i = 1, 2, \dots, n-1\}.$$

The (a, d) -edge-antimagic vertex labelings of P_n for d negative we define as follows

$$\beta_{-l}^n(v_i) = \beta_l^n(v_{n+1-i}) \quad \text{for } i = 1, 2, \dots, n,$$

where $l = 1, 2$.

3. Partitions with determined differences

For construction of vertex and edge labelings of plane graphs we will use the partitions of a set of integers with determined differences.

Let n, k, d and i be positive integers. We will consider the partition $\mathcal{P}_{k,d}^n$ of the set $\{1, 2, \dots, kn\}$ into $n, n \geq 2, k$ -tuples such that the difference between the sum of the numbers in the $(i+1)$ th k -tuple and the sum of the numbers in the i th k -tuple is always equal to the constant d , where $i = 1, 2, \dots, n-1$. Thus they form an arithmetic sequence with the difference d . By the symbol $\mathcal{P}_{k,d}(i)$ we denote the i th k -tuple in the partition with the difference d , where $i = 1, 2, \dots, n$.

Let $\sum \mathcal{P}_{k,d}^n(i)$ be the sum of the numbers in $\mathcal{P}_{k,d}^n(i)$. Evidently $\sum \mathcal{P}_{k,d}^n(i+1) - \sum \mathcal{P}_{k,d}^n(i) = d$. It is obvious that if there exists a partition of the set $\{1, 2, \dots, kn\}$ with the difference d , there also

exists a partition with the difference $-d$. By the notation $\mathcal{P}_{k,d}^n(i) \oplus c$ we mean that we add the constant c to every number in $\mathcal{P}_{k,d}^n(i)$.

If $k = 1$ then only the following partition of the set $\{1, 2, \dots, n\}$ is possible

$$\mathcal{P}_{1,1}^n(i) = \{i\} \quad \text{for } i = 1, 2, \dots, n.$$

If $k = 2$ then we have several partitions of the set $\{1, 2, \dots, 2n\}$. Let us define the partitions into 2-tuples in the following way:

$$\begin{aligned} \mathcal{P}_{2,0}^n(i) &= \{i, 2n+1-i\}, \\ \sum \mathcal{P}_{2,0}^n(i) &= 2n+1, & \text{for } i = 1, 2, \dots, n. \\ \mathcal{P}_{2,2}^n(i) &= \{i, n+i\}, \\ \sum \mathcal{P}_{2,2}^n(i) &= n+2i, & \text{for } i = 1, 2, \dots, n. \\ \mathcal{P}_{2,4}^n(i) &= \{2i-1, 2i\}, \\ \sum \mathcal{P}_{2,4}^n(i) &= 4i-1, & \text{for } i = 1, 2, \dots, n. \end{aligned}$$

Moreover, for $3 \leq n \equiv 1 \pmod{2}$

$$\begin{aligned} \mathcal{P}_{2,1}^n(i) &= \begin{cases} \left\{ \frac{n+1}{2} + \frac{i-1}{2}, n+1 + \frac{i-1}{2} \right\} & \text{for } i \equiv 1 \pmod{2}, \\ \left\{ \frac{i}{2}, n + \frac{n+1}{2} + \frac{i}{2} \right\} & \text{for } i \equiv 0 \pmod{2}, \end{cases} \\ \sum \mathcal{P}_{2,1}^n(i) &= n + \frac{n+1}{2} + i, & \text{for } i = 1, 2, \dots, n. \end{aligned}$$

Note that we are also able to obtain the partitions into 2-tuples $\mathcal{P}_{2,0}^n(i)$ and $\mathcal{P}_{2,2}^n(i)$ as $\mathcal{P}_{1,s}^n(i) \cup (\mathcal{P}_{1,t}^n(i) \oplus n)$, where $s, t = \pm 1$. We can use this idea to construct the other partitions. More precisely,

$$\mathcal{P}_{k,d}^n(i) = \mathcal{P}_{l,s}^n(i) \cup (\mathcal{P}_{m,t}^n(i) \oplus ln),$$

where $k = l + m$.

For example, we are able to obtain $\mathcal{P}_{3,d}^n(i)$ from the partitions $\mathcal{P}_{1,s}^n(i)$, $s = \pm 1$ and $\mathcal{P}_{2,t}^n(i)$, $t = 0, \pm 2, \pm 4$ and also $t = \pm 1$ for n odd. It means, $\mathcal{P}_{3,d}^n$ exists for $d = \pm 1, \pm 3, \pm 5$ and if $n \equiv 1 \pmod{2}$ also for $d = 0, \pm 2$. Moreover, we are able to construct $\mathcal{P}_{3,9}^n$ in the following way

$$\begin{aligned} \mathcal{P}_{3,9}^n(i) &= \{3(i-1)+1, 3(i-1)+2, 3(i-1)+3\}, \\ \sum \mathcal{P}_{3,9}^n(i) &= 9i-3, \quad \text{for } i = 1, 2, \dots, n. \end{aligned}$$

Thus $\mathcal{P}_{3,d}^n$ exists for $d = \pm 1, \pm 3, \pm 5, \pm 9$ and if $n \equiv 1 \pmod{2}$ also for $d = 0, \pm 2$.

For the partition into 4-tuples we can use the following fact

$$\mathcal{P}_{4,d}^n(i) = \mathcal{P}_{l,s}^n(i) \cup (\mathcal{P}_{m,t}^n(i) \oplus ln),$$

where $l = 3, m = 1$ or $l = 2, m = 2$. Also

$$\begin{aligned}\mathcal{P}_{4,16}^n(i) &= \{4(i-1) + 1, 4(i-1) + 2, 4(i-1) + 3, 4(i-1) + 4\}, \\ \sum \mathcal{P}_{4,16}^n(i) &= 16i - 6, \quad \text{for } i = 1, 2, \dots, n.\end{aligned}$$

Thus $\mathcal{P}_{4,d}^n$ exists for $d = 0, \pm 2, \pm 4, \pm 6, \pm 8, \pm 10, \pm 16$ and if $n \equiv 1 \pmod{2}$ also for $d = \pm 1, \pm 3, \pm 5$.

Let us note that each of the defined partition $\mathcal{P}_{k,d}^n$ has the property that

$$\sum \mathcal{P}_{k,d}^n(i) = C_{k,d}^n + di,$$

where $C_{k,d}^n$ is a constant depending on the parameters k and d .

4. d -antimagic labelings for certain families of plane graphs

In this section, we shall use the edge-antimagic labelings of paths P_n and the partitions of the set $\{1, 2, \dots, kn\}$ with determined differences described in the previous two sections to examine the existence of a super d -antimagic labeling for several families of plane graphs.

The *friendship graph* F_n is a set of n triangles having a common central vertex, say v , and otherwise disjoint. The friendship graph F_n has $2n$ vertices of degree 2, say v_i, u_i for $i = 1, 2, \dots, n$, and $3n$ edges, say $v_i v, u_i v, v_i u_i$ for $i = 1, 2, \dots, n$. Let us define the 3-sided face $f_i, i = 1, 2, \dots, n$, as the face bounded by the edges $vv_i, v_i u_i$ and $u_i v$ and let f be the external unbounded face.

Theorem 4.1. *The friendship graph $F_n, n \geq 2$, has a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{1, 3, 5, 7, 9, 11, 13\}$.*

Moreover, if $n \equiv 1 \pmod{2}$ then the graph F_n also admits a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 2, 4, 6, 8, 10\}$.

Proof. We define the bijection $g_1 : V(F_n) \cup E(F_n) \rightarrow \{1, 2, \dots, 5n + 1\}$ as follows:

$$\begin{aligned}\{g_1(v_i), g_1(u_i)\} &= \mathcal{P}_{2,k}^n(i), & i = 1, 2, \dots, n, \\ g_1(v) &= 2n + 1, \\ \{g_1(v_i v), g_1(v_i u_i), g_1(u_i v)\} &= \mathcal{P}_{3,l}^n(i) \oplus (2n + 1) & i = 1, 2, \dots, n,\end{aligned}$$

where $k = 0, \pm 2, \pm 4$ or for $n \equiv 1 \pmod{2}$ also $k = \pm 1$, and $l = \pm 1, \pm 3, \pm 5, \pm 9$ or for $n \equiv 1 \pmod{2}$ also $l = 0, \pm 2$.

It is not difficult to check that the vertices are labeled by the smallest possible numbers $1, 2, \dots, 2n + 1$. Moreover, for the weight of the face $f_i, i = 1, 2, \dots, n$, we obtain

$$\begin{aligned}w_{g_1}(f_i) &= (g_1(v_i) + g_1(u_i)) + g_1(v) + (g_1(v_i v) + g_1(v_i u_i) + g_1(u_i v)) \\ &= \sum \mathcal{P}_{2,k}^n(i) + (2n + 1) + \sum (\mathcal{P}_{3,l}^n(i) \oplus (2n + 1)) \\ &= (C_{2,k}^n + ki) + (2n + 1) + (C_{3,l}^n + li + 3(2n + 1)) \\ &= C_{2,k}^n + C_{3,l}^n + 4(2n + 1) + (k + l)i.\end{aligned}$$

As $k = 0, \pm 2, \pm 4$ or for $n \equiv 1 \pmod{2}$ also $k = \pm 1$, and $l = \pm 1, \pm 3, \pm 5, \pm 9$ or for $n \equiv 1 \pmod{2}$ also $l = 0, \pm 2$, we obtain that $k + l \in \{1, 3, 5, 7, 9, 11, 13\}$ and for n odd we also get $k + l \in \{0, 2, 4, 6, 8, 10\}$. Thus under the labeling g_1 the weights of the 3-sided faces form an arithmetic sequence with the desired difference, which completes the proof. \square

If we replace every edge $v_i u_i, i = 1, 2, \dots, n$, of the friendship graph F_n by a path of length two with vertices v_i, w_i, u_i , then we obtain a graph, say B_n , with the vertex set $V(B_n) = \{v_i, w_i, u_i, v : i = 1, 2, \dots, n\}$ and the edge set $E(B_n) = \{v_i v, u_i v, v_i w_i, w_i u_i : i = 1, 2, \dots, n\}$. Let us define the 4-sided face $f_i, i = 1, 2, \dots, n$, as the face bounded by the edges $vv_i, v_i w_i, w_i u_i$ and $u_i v$ and let f be the external unbounded face.

Theorem 4.2. *The graph $B_n, n \geq 2$, has a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{1, 3, 5, \dots, 21, 25\}$.*

Moreover, if $n \equiv 1 \pmod{2}$ then the graph B_n also admits a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 2, 4, \dots, 18\}$.

Proof. We define the bijection $g_2 : V(B_n) \cup E(B_n) \rightarrow \{1, 2, \dots, 7n + 1\}$ in the following way:

$$\begin{aligned} \{g_2(v_i), g_2(w_i), g_2(u_i)\} &= \mathcal{P}_{3,k}^n(i), & i = 1, 2, \dots, n, \\ g_2(v) &= 3n + 1, \\ \{g_2(v_i v), g_2(v_i w_i), g_2(w_i u_i), g_2(u_i v)\} &= \mathcal{P}_{4,l}^n(i) \oplus (3n + 1), & i = 1, 2, \dots, n. \end{aligned}$$

It is not difficult to see that the vertices are labeled by the numbers $1, 2, \dots, 3n + 1$. Moreover, for the weight of the face $f_i, i = 1, 2, \dots, n$, we have

$$\begin{aligned} w_{g_2}(f_i) &= (g_2(v_i) + g_2(w_i) + g_2(u_i)) + g_2(v) \\ &\quad + (g_2(v_i v) + g_2(v_i w_i) + g_2(w_i u_i) + g_2(u_i v)) \\ &= \sum \mathcal{P}_{3,k}^n(i) + (3n + 1) + \sum (\mathcal{P}_{4,l}^n(i) \oplus (3n + 1)) \\ &= (C_{3,k}^n + ki) + (3n + 1) + (C_{4,l}^n + li + 4(3n + 1)) \\ &= C_{3,k}^n + C_{4,l}^n + 5(3n + 1) + (k + l)i, \end{aligned}$$

where $k = \pm 1, \pm 3, \pm 5, \pm 9$ or for $n \equiv 1 \pmod{2}$ also $k = 0, \pm 2$, and $l = 0, \pm 2, \pm 4, \pm 6, \pm 8, \pm 10, \pm 16$ or for $n \equiv 1 \pmod{2}$ also $l = \pm 1, \pm 3, \pm 5$. It means that g_2 is a super d -antimagic labeling of type $(1, 1, 0)$ of B_n , for $d = 1, 3, 5, \dots, 21, 25$ and if $n \equiv 1 \pmod{2}$ then $d = 0, 2, 4, \dots, 18$. \square

A *triangular snake* E_n is a triangular cactus whose block-cutpoint graph is a path, i.e. E_n is obtained from a path v_1, v_2, \dots, v_{n+1} by joining v_i and v_{i+1} to a new vertex u_i , for $i = 1, 2, \dots, n$. Let f_i be the 3-sided face, $i = 1, 2, \dots, n$, bounded by the edges $v_i u_i, u_i v_{i+1}$ and $v_i v_{i+1}$. We denote the external unbounded face by the symbol f .

Theorem 4.3. *The graph $E_n, n \geq 2$, has a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 1, 2, \dots, 12\}$.*

Proof. Define the bijection $g_3 : V(E_n) \cup E(E_n) \rightarrow \{1, 2, \dots, 5n + 1\}$ as follows:

$$\begin{aligned} g_3(v_i) &= \alpha_k^{n+1}(v_i), & i &= 1, 2, \dots, n+1, \\ g_3(u_i) &= \alpha_k^{n+1}(v_i v_{i+1}), & i &= 1, 2, \dots, n, \\ \{g_3(v_i u_i), g_3(u_i v_{i+1}), g_3(v_{i+1} v_i)\} &= \mathcal{P}_{3,l}^n(i) \oplus (2n+1), & i &= 1, 2, \dots, n. \end{aligned}$$

The labeling g_3 assigns the numbers $1, 2, \dots, 2n+1$ to the vertices of the graph E_n . For the weight of the face $f_i, i = 1, 2, \dots, n$, we have

$$\begin{aligned} w_{g_3}(f_i) &= (g_3(v_i) + g_3(u_i) + g_3(v_{i+1})) + (g_3(v_i u_i) + g_3(u_i v_{i+1}) + g_3(v_{i+1} v_i)) \\ &= w_{\alpha_k^{n+1}}(v_i v_{i+1}) + \sum \mathcal{P}_{3,l}^n(i) \oplus (2n+1) \\ &= (C_{\alpha,k}^{n+1} + ki) + (C_{3,l}^n + li + 3(2n+1)) \\ &= C_{\alpha,k}^{n+1} + C_{3,l}^n + 3(2n+1) + (k+l)i, \end{aligned}$$

where $k = 0, \pm 1, \pm 2, \pm 3$ and $l = \pm 1, \pm 3, \pm 5, \pm 9$, moreover for $n \equiv 1 \pmod{2}$ also $l = 0, \pm 2$. Analogously as in the proof of the previous theorem we obtain that for $d \in \{0, 1, 2, \dots, 12\}$ the bijection g_3 is a super d -antimagic labeling of type $(1, 1, 0)$ of the graph E_n . \square

If we replace every edge $v_i v_{i+1}, i = 1, 2, \dots, n$, of the triangular snake E_n by a path of length two with vertices v_i, w_i, v_{i+1} , then we obtain a graph, say G_n , with the vertex set $V(G_n) = \{v_1, v_2, \dots, v_{n+1}, u_1, u_2, \dots, u_n, w_1, w_2, \dots, w_n\}$ and the edge set $E(G_n) = \{v_i u_i, u_i v_{i+1}, v_i w_i, w_i v_{i+1} : i = 1, 2, \dots, n\}$. Let us define the 4-sided face $f_i, i = 1, 2, \dots, n$, as the face bounded by the edges $v_i u_i, u_i v_{i+1}, v_i w_i$ and $w_i v_{i+1}$ and let f be the external unbounded face.

Theorem 4.4. *The graph $G_n, n \geq 2$, has a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 1, 2, \dots, 22\}$.*

Proof. Define the bijection $g_4 : V(G_n) \cup E(G_n) \rightarrow \{1, 2, \dots, 7n + 1\}$ in the following way:

$$\begin{aligned} g_4(v_i) &= \beta_k^{n+1}(v_i), & i &= 1, 2, \dots, n+1, \\ \{g_4(u_i), g_4(w_i)\} &= \mathcal{P}_{2,l}^n(i) \oplus (n+1), & i &= 1, 2, \dots, n, \\ \{g_4(v_i u_i), g_4(u_i v_{i+1}), g_4(v_i w_i), g_4(w_i v_{i+1})\} &= \mathcal{P}_{4,m}^n(i) \oplus (3n+1), & i &= 1, 2, \dots, n. \end{aligned}$$

It is easy to verify that the labeling g_4 assigns integers $1, 2, \dots, 3n+1$ to the vertices. By direct computation we obtain that the weight of the face $f_i, i = 1, 2, \dots, n$, admits a value

$$\begin{aligned} w_{g_4}(f_i) &= (g_4(v_i) + g_4(v_{i+1})) + (g_4(u_i) + g_4(w_i)) \\ &\quad + (g_4(v_i u_i) + g_4(u_i v_{i+1}) + g_4(v_i w_i) + g_4(w_i v_{i+1})) \\ &= w_{\beta_k^{n+1}}(v_i v_{i+1}) + \sum \mathcal{P}_{2,l}^n(i) \oplus (n+1) + \sum \mathcal{P}_{4,m}^n(i) \oplus (3n+1) \\ &= (C_{\beta,k}^{n+1} + ki) + (C_{2,l}^n + li + 2(n+1)) + (C_{4,m}^n + mi + 4(3n+1)) \\ &= C_{\beta,k}^{n+1} + C_{2,l}^n + C_{4,m}^n + 14n + 6 + (k+l+m)i, \end{aligned}$$

where $k = \pm 1, \pm 2, l = 0, \pm 2, \pm 4$ and $m = 0, \pm 2, \pm 4, \pm 6, \pm 8, \pm 10, \pm 16$. After some manipulations we obtain that there exists a super d -antimagic labeling of the graph G_n for every difference $d \in \{0, 1, 2, \dots, 22\}$. \square

Let ladder L_n be a Cartesian product $L_n \simeq P_n \times P_2$ of a path on n vertices with a path on two vertices. Let $V(L_n) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n\}$ be the vertex set and $E(L_n) = \{v_i v_{i+1}, u_i u_{i+1} : i = 1, 2, \dots, n-1\} \cup \{v_i u_i : i = 1, 2, \dots, n\}$ be the edge set of ladder. Let us define the 4-sided face $f_i, i = 1, 2, \dots, n$, as the face bounded by the edges $v_i v_{i+1}, v_{i+1} u_{i+1}, u_i u_{i+1}$ and $v_i u_i$ and let f be the external unbounded face.

Theorem 4.5. *The ladder $L_n \simeq P_n \times P_2, n \geq 3$, admits a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 1, 2, \dots, 10\}$.*

Proof. Construct the bijective function $g_5 : V(L_n) \cup E(L_n) \rightarrow \{1, 2, \dots, 5n-2\}$ as follows:

$$\begin{aligned} g_5(v_i) &= \beta_k^n(v_i), & i &= 1, 2, \dots, n, \\ g_5(u_i) &= \beta_l^n(v_i) + n, & i &= 1, 2, \dots, n, \\ g_5(v_i u_i) &= \beta_m^n(v_i) + 2n, & i &= 1, 2, \dots, n, \\ \{g_5(v_i v_{i+1}), g_5(u_i u_{i+1})\} &= \mathcal{P}_{2,t}^{n-1}(i) \oplus (3n), & i &= 1, 2, \dots, n-1. \end{aligned}$$

It is a routine procedure to verify that the vertices are labeled by the smallest possible numbers $1, 2, \dots, 2n$. Moreover, for the weight of the face $f_i, i = 1, 2, \dots, n-1$, we obtain

$$\begin{aligned} w_{g_5}(f_i) &= (g_5(v_i) + g_5(v_{i+1})) + (g_5(u_i) + g_5(u_{i+1})) + (g_5(v_i u_i) + g_5(v_{i+1} u_{i+1})) \\ &\quad + (g_5(v_i v_{i+1}) + g_5(u_i u_{i+1})) \\ &= w_{\beta_k^n}(v_i v_{i+1}) + (w_{\beta_l^n}(v_i v_{i+1}) + 2n) + (w_{\beta_m^n}(v_i v_{i+1}) + 4n) \\ &\quad + \sum \mathcal{P}_{2,t}^{n-1}(i) \oplus (3n) \\ &= (C_{\beta,k}^n + ki) + (C_{\beta,l}^n + li + 2n) + (C_{\beta,m}^n + mi + 4n) \\ &\quad + (C_{2,t}^{n-1} + ti + 6n) \\ &= C_{\beta,k}^n + C_{\beta,l}^n + C_{\beta,m}^n + C_{2,t}^{n-1} + 12n + (k + l + m + t)i. \end{aligned}$$

As $k = \pm 1, \pm 2, l = \pm 1, \pm 2, m = \pm 1, \pm 2$ and $t = 0, \pm 2, \pm 4$ we obtain that $k + l + m + t \in \{0, 1, 2, 3, \dots, 10\}$. This completes the proof. \square

Another variation of a ladder graph is specified as follows. A ladder $\mathbb{L}_n, n \geq 2$, is a graph obtained by completing the ladder $L_n \simeq P_n \times P_2$ by $n-1$ edges such that $V(\mathbb{L}_n) = \{v_1, v_2, \dots, v_{2n}\}$ is the vertex set and $E(\mathbb{L}_n) = \{v_1 v_2, v_2 v_3, \dots, v_{2n-1} v_{2n}, v_1 v_3, v_2 v_4, \dots, v_{2n-2} v_{2n}\}$ is the edge set.

Theorem 4.6. *The graph $\mathbb{L}_n, n \geq 2$, admits a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 1, 2, \dots, 6\}$.*

Proof. Construct the bijective function $g_6 : V(\mathbb{L}_n) \cup E(\mathbb{L}_n) \rightarrow \{1, 2, \dots, 6n-1\}$ in the following way:

$$\begin{aligned}
 g_6(v_i) &= i, & i &= 1, 2, \dots, 2n, \\
 g_6(v_i v_{i+1}) &= \beta_k^{2n-1}(v_i) + 2n, & i &= 1, 2, \dots, 2n-1, \\
 g_6(v_i v_{i+2}) &= \mathcal{P}_{1,l}^{2n-2}(i) \oplus (4n-1), & i &= 1, 2, \dots, 2n-2.
 \end{aligned}$$

The reader can easily verify that the labeling g_6 assigns integers $1, 2, \dots, 2n$ to the vertices and a weight of the face f_i , $i = 1, 2, \dots, 2n-2$, is

$$\begin{aligned}
 w_{g_6}(f_i) &= (g_6(v_i) + g_6(v_{i+1}) + g_6(v_{i+2})) + (g_6(v_i v_{i+1}) + g_6(v_{i+1} v_{i+2})) \\
 &\quad + g_6(v_i v_{i+2}) \\
 &= (i + (i+1) + (i+2)) + (w_{\beta_k^{2n-1}}(v_i v_{i+1}) + 4n) \\
 &\quad + \sum \mathcal{P}_{1,l}^{2n-2}(i) \oplus (4n-1) \\
 &= (3 + 3i) + (C_{\beta,k}^{2n-1} + ki + 4n) + (C_{1,l}^{2n-2} + li + (4n-1)) \\
 &= C_{\beta,k}^{2n-1} + C_{1,l}^{2n-2} + 8n + 2 + (3 + k + l)i.
 \end{aligned}$$

Since $k = \pm 1, \pm 2$, and $l = \pm 1$ we are able to show that $3 + k + l \in \{0, 1, 2, \dots, 6\}$. This implies that the labeling g_6 is a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 1, 2, \dots, 6\}$ of the graph \mathbb{L}_n . \square

If we replace every edge $v_i v_{i+1}$ (respectively, every edge $u_i u_{i+1}$), $i = 1, 2, \dots, n-1$, of the ladder $L_n \simeq P_n \times P_2$ by a path of length two with vertices v_i, w_i, v_{i+1} (respectively, u_i, w_{n-1+i}, u_{i+1}) then we obtain a graph, say H_n , with the vertex set $V(H_n) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n, w_1, w_2, \dots, w_{2n-2}\}$ and the edge set $E(H_n) = \{v_i w_i, w_i v_{i+1}, u_i w_{n-1+i}, w_{n-1+i} u_{i+1} : i = 1, 2, \dots, n-1\} \cup \{v_i u_i : i = 1, 2, \dots, n\}$.

Let us define the 6-sided face f_i , $i = 1, 2, \dots, n-1$, as the face bounded by the edges $v_i w_i, w_i v_{i+1}, v_{i+1} u_{i+1}, u_{i+1} w_{n-1+i}, w_{n-1+i} u_i, u_i v_i$ and let f be the external unbounded face.

Theorem 4.7. *The graph H_n , $n \geq 2$, admits a super d -antimagic labeling of type $(1, 1, 0)$ for $d \in \{0, 1, 2, \dots, 26\}$.*

Proof. We define the bijection $g_7 : V(H_n) \cup E(H_n) \rightarrow \{1, 2, \dots, 9n-2\}$ in the following way:

$$\begin{aligned}
 g_7(v_i) &= \beta_k^n(v_i), & i &= 1, 2, \dots, n, \\
 g_7(u_i) &= \beta_l^n(v_i), & i &= 1, 2, \dots, n, \\
 \{g_7(w_i), g_7(w_{n-1+i})\} &= \mathcal{P}_{2,m}^{n-1}(i) \oplus (2n), & i &= 1, 2, \dots, n-1, \\
 g_7(v_i u_i) &= \beta_t^n(v_i) + 4n-2, & i &= 1, 2, \dots, n, \\
 \{g_7(v_i w_i), g_7(w_i v_{i+1}), g_7(u_i w_{n-1+i}), g_7(w_{n-1+i} u_{i+1})\} \\
 &= \mathcal{P}_{4,s}^{n-1}(i) \oplus (5n-2), & i &= 1, 2, \dots, n-1.
 \end{aligned}$$

It is easy to see that the vertices of H_n are labeled by the smallest possible integers $1, 2, \dots, 4n-2$. For the weight of the face f_i , $i = 1, 2, \dots, n-1$, we get

$$\begin{aligned}
 w_{g_7}(f_i) &= (g_7(v_i) + g_7(v_{i+1})) + (g_7(u_i) + g_7(u_{i+1})) + (g_7(w_i) + g_7(w_{n-1+i})) \\
 &\quad + (g_7(v_i u_i) + g_7(v_{i+1} u_{i+1})) + (g_7(v_i w_i) + g_7(w_i v_{i+1}) + g_7(u_i w_{n-1+i})) \\
 &\quad + g_7(w_{n-1+i} u_{i+1})) \\
 &= w_{\beta_k^n}(v_i v_{i+1}) + (w_{\beta_l^n}(v_i v_{i+1}) + 2n) + \sum \mathcal{P}_{2,m}^{n-1}(i) \oplus (2n) \\
 &\quad + (w_{\beta_t^n}(v_i v_{i+1}) + 8n - 4) + \sum \mathcal{P}_{4,s}^{n-1}(i) \oplus (5n - 2) \\
 &= (C_{\beta,k}^m + ki) + (C_{\beta,l}^m + li + 2n) + (C_{2,m}^{m-1} + mi + 2(2n)) \\
 &\quad + (C_{\beta,t}^m + ti + 8n - 4) + (C_{4,s}^{m-1} + si + 4(5n - 2)) \\
 &= C_{\beta,k}^m + C_{\beta,l}^m + C_{\beta,t}^m + C_{2,m}^{m-1} + C_{4,s}^{m-1} + 34n - 12 + (k + l + m + t + s)i.
 \end{aligned}$$

Since $k = \pm 1, \pm 2$, $l = \pm 1, \pm 2$, $m = 0, \pm 2, \pm 4$, $t = \pm 1, \pm 2$ and $s = 0, \pm 2, \pm 4, \pm 6, \pm 8, \pm 10, \pm 16$ we get that $k + l + m + t + s \in \{0, 1, 2, \dots, 26\}$. Thus g_7 is the required super d -antimagic labeling of type $(1, 1, 0)$ of the graph H_n . This completes the proof. \square

5. Concluding Remarks

In the foregoing section we studied the super d -antimagic labelings for the seven families of plane graphs. We have shown that there exist super d -antimagic labelings of type $(1, 1, 0)$ for these graphs for wide variety of difference d . We conclude with the following open problems.

Problem 1. Find the upper bound for the feasible values of the difference d which makes a super d -antimagic labelings of type $(1, 1, 0)$ possible for the studied families of plane graphs.

Problem 2. Find other feasible values of the difference d and the corresponding super d -antimagic labelings of type $(1, 1, 0)$ for the studied families of plane graphs.

Acknowledgement

The work was supported by Slovak VEGA Grant 1/0130/12.

- [1] G. Ali, M. Bača, F. Bashir and A. Semaničová-Feňovčíková, On face antimagic labelings of disjoint union of prisms, *Utilitas Math.* **85** (2011), 97–112.
- [2] M. Bača, On magic labelings of grid graphs, *Ars Combin.* **33** (1992), 295–299.
- [3] M. Bača, On magic labelings of honeycomb, *Discrete Math.* **105** (1992), 305–311.
- [4] M. Bača, F. Bashir and A. Semaničová, Face antimagic labelings of antiprisms, *Utilitas Math.* **84** (2011), 209–224.
- [5] M. Bača, E.T. Baskoro, S. Jendroľ and M. Miller, Antimagic labelings of hexagonal planar maps, *Utilitas Math.* **66** (2004), 231–238.

- [6] M. Bača, L. Brankovic and A. Semaničová-Feňovčíková, Labelings of plane graphs containing Hamilton path, *Acta Math. Sinica - English Series* **27** (4) (2011), 701–714.
- [7] M. Bača, S. Jendroň, M. Miller and J. Ryan, Antimagic labelings of generalized Petersen graphs that are plane, *Ars Combin.* **73** (2004), 115–128.
- [8] M. Bača, Y. Lin and M. Miller, Antimagic labelings of grids, *Utilitas Math.* **72** (2007), 65–75.
- [9] M. Bača, Y. Lin, F.A. Muntaner-Batle, Super edge-antimagic labelings of the path-like trees, *Utilitas Math.* **73** (2007), 117–128.
- [10] M. Bača and M. Miller, On d -antimagic labelings of type $(1, 1, 1)$ for prisms, *J. Combin. Math. Combin. Comput.* **44** (2003), 199–207.
- [11] M. Bača and M. Miller, *Super Edge-Antimagic Graphs: A Wealth of Problems and Some Solutions*, BrownWalker Press, Boca Raton, Florida, 2008.
- [12] M. Bača, M. Miller, O. Phanalasy and A. Semaničová-Feňovčíková, Super d -antimagic labelings of disconnected plane graphs, *Acta Math. Sinica - English Series* **26** (12) (2010), 2283–2294.
- [13] H. Enomoto, A.S. Lladó, T. Nakamigawa and G. Ringel, Super edge-magic graphs, *SUT J. Math.* **34** (1998), 105–109.
- [14] J. Gallian, A dynamic survey of graph labeling. *The Electronic J. Combin.* **17** (2010), #DS6.
- [15] A. Kotzig and A. Rosa, Magic valuations of finite graphs, *Canad. Math. Bull.* **13** (1970), 451–461.
- [16] Ko-Wei Lih, On magic and consecutive labelings of plane graphs, *Utilitas Math.* **24** (1983), 165–197.
- [17] Y. Lin, Slamin, M. Bača and M. Miller, On d -antimagic labelings of prisms, *Ars Combin.* **72** (2004), 65–76.
- [18] R. Simanjuntak, F. Bertault and M. Miller, Two new (a, d) -antimagic graph labelings. *Proc. of Eleventh Australasian Workshop on Combinatorial Algorithms* (2000), 179–189.
- [19] K.A. Sugeng, M. Miller, Y. Lin and M. Bača, Face antimagic labelings of prisms, *Utilitas Math.* **71** (2006), 269–286.
- [20] W.D. Wallis, *Magic Graphs*, Birkhäuser, Boston - Basel - Berlin, 2001.