



## Original Article

# In-vitro Antioxidant activity, Biosafety, Nociceptive and Anti-inflammatory potential of Acetone Polyherbal (ELNA) Extract in mice

MAC DONALD Idu Chike U. Onuigbo, BENJAMIN O. Gabriel;

Phytomedicine Unit, Department of Plant Biology and Biotechnology, PMB 1154, Benin City, Edo State, Nigeria.

## ARTICLE INFOR

## Article history:

Received 25 July 2021

Revised 08 September 2021

Accepted 20 December 2021

## Keywords:

Biosafety,  
Nociceptive,  
Anti-inflammatory,  
In-vitro Antioxidant,  
Polyherbal Elna.

## ABSTRACT

ELNA (*Moringa oleifera*, *Crateva religiosa* and *Curcuma longa*) is a polyherbal formulation comprising of *Moringa oleifera*, *Crateva religiosa* and *Curcuma longa*. It is used traditionally in the treatment of inflammation, gastro-intestinal infections, hypertension and immune compromised diseases. This study investigate the *in-vitro* anti-oxidant capacity, biosafety, nociceptive and anti-inflammatory potential of polyherbal ELNA acetone extract. *In-vitro* anti-oxidant study was done on the extract using a standard protocol of 1, 1, Diphenyl 2, picrylhydrazyl (DPPH) radical scavenging activity. Toxicological profiling was done using standard methods. Acetic acid, hot plate and egg albumin models were designed for the nociceptive and anti-inflammatory properties of the extract. DPPH scavenging property of the extract showed significant increase at graded concentration. Acute toxicity study of ELNA acetone extract revealed no toxic effect, with absent mortality and less adverse effect ( $LD_{50} > 5000$  mg/kg body weight). The sub-chronic toxicity study of the acetone extract for 28 days, showed no significant difference ( $p = 0.05$ ) in the organ to body weight ratio. The haematological indexes indicated no significant different except at 400 mg/kg that elicited slight significant ( $p < 0.05$ ) increase in the platelet. The results showed that graded doses of the extract at (400, 800 and 1200 mg/kg body weight) exhibited significant ( $p < 0.05$ ) decrease in peripheral and central pain also a decrease in the paw edema volume of inflammation. This was achieved in dose dependent manner. The result of this study established a pharmacological evidence for the traditional use of ELNA as an analgesic and anti-inflammatory agent, it also present information on the anti-oxidant properties and toxicity profile of the formulation.

© 2021 Faculty of Natural Sciences and Life, University of Echahid Hamma Lakhdar. All rights reserved

## 1. Introduction

The use of herbs as medicine is the oldest form of healthcare known to humanity and has been used in all cultures throughout history [1]. Early humans recognized their dependence on nature for a healthy life and since that time they have depended on the diversity of plant resources for food, clothing, shelter and medicine to cure myriads of ailments. Led by instinct, taste and experience, primitive men and women treated illness by using plants, animal parts and minerals that were not part of their usual diet. Polyherbal preparations are herbal preparation of multiple herbs. Plant formulation and combined extracts of plants are used as drug of choice rather than individual drugs [2], this could be linked to the fact that the individual plants parts making up the polyherbal formulation could act in synergy or have a potentiating effect. Several works have been reported on the effectiveness of polyherbal formulations. In a work on the development and evaluation of analgesic polyherbal formulation containing

some indigenous medicinal plants, the formulation was found to have a significant ( $p < 0.05$ ) analgesic activity in a dose dependent manner [3].

*Moringa oleifera* or Moringa leaves contain several bio-active compounds that exert direct effect on blood pressure thus these can be used for stabilizing blood pressure. Moringa compounds leading to blood pressure lowering effect includes nitrile, mustard oil glycosides and thiocarbamate glycosides present in Moringa leaves [4]. The  $\beta$ -sitosterol present in them is responsible for cholesterol lowering effect [5]. In addition to earlier mentioned bradycardia effect of Moringa leaves, all parts of Moringa are reported with somewhat cardiac and circulatory stimulant activity. Root bark of Moringa contains alkaloid moringinine which acts as cardiac stimulant through its effect on sympathetic nervous system [6].

*Crateva religiosa* is a much branched deciduous tree belonging to the family Capparidaceae, commonly called as Varuna [7]. The leaves were used in combination with other plants by

\* Corresponding author : Benjamin O. Gabriel. Tel.: +2348050607009; +2348037373057; +2347032695752

E-mail address: mcdonald@uniben.edu; chike.onuigbo@gmail.com; alexthemain076@gmail.com

Peer review under responsibility of University of Echahid Hamma Lakhdar. © 2021 University of Echahid Hamma Lakhdar. All rights reserved.

doi : <http://dx.doi.org/10.5281/zenodo.5810172>.

traditional healers and local populations as analgesic, antispasmodic, antimalarial and antidiarrheal. It possessed antimicrobial activity against *Candida albicans*, *Candida tropicalis*, *Candida krusei*, *Cryptococcus marinus* and *Aspergillus niger*, hepatitis, edema, ascites, urinary stones and arthritis [8,9].

*Curcuma longa* is commonly known as turmeric is a perennial member of the Zingiberaceae family. The rhizome contains active compound curcumin which is responsible for the yellowish colour. Curcumin is a lipophilic polyphenol that is nearly insoluble in water but is quite stable in the acidic pH of the stomach. Turmeric is reported to reduce the uptake of cholesterol from the gut and increase the high-density lipids (HDL) cholesterol and decrease low-density lipids (LDL) type. It can also inhibit the peroxidation of serum LDL, which leads to atherosclerotic lesions. Thus, turmeric can prevent coronary problems and heart diseases [10]. This study elicited the biosafety profiling of the polyherbal extract with its therapeutic effect being validated of its nociceptive, anti-inflammatory and anti-oxidant properties.

## 2. Materials and Methods

### 2.1. Plant Collection

ELNA consist of three herbal leaves (*Moringa oleifera*, *Crateva religiosa* and *Curcuma longa*) and it was obtained from an herbal therapist in Lagos. The plant was identified and authenticated by Dr. O. Timothy a plant Taxonomist in the Department of Plant Biology and Biotechnology University of Benin, Benin City with a voucher number GW-E174.

### 2.2. Plant Preparation

The various plant was shade dried, pulverized and prepared into extract with a formulation involved the combination of three plants leaves in their dried powdered form at different proportions. It was extracted in acetone solvent system, using Soxhlet Extractor. The filtrate was dried freeze dryer. Sample was stored in a refrigerator for further use.

### 2.3. DPPH Scavenging Activity

A solution of 0.1 mM DPPH in methanol was prepared, and 1.0 ml of this solution was mixed with 3.0 ml of extract in methanol containing 0.01 – 0.2 mg/ml of the extract. The reaction mixture was vortexed thoroughly and left in the dark at room temperature for 30 min. The absorbance of the mixture was measured spectrophotometrically at 517 nm. Ascorbic acid was used as standard. The DPPH radical scavenging activity was calculated according to the following equation:

$$\% \text{ Inhibition of DPPH} = \frac{(A_{\text{control}} - A_{\text{sample}})}{A_{\text{control}}} \times 100$$

Where A sample and A control are absorbance of sample and control respectively. Decrease in the absorbance of the DPPH solution indicates an increase in DPPH radical scavenging activity.

### 2.4. Experimental Animals

Healthy albino mice weighing between 13-25 g obtained from the Animal House of University of Jos, Plateau State and were kept in the Animal facility of National Institute for Pharmaceutical Research and Development, Idu Industrial Layout, Abuja and kept for about 2 weeks for acclimatization. They were housed in groups of 5 in clean dry cages 34x47x18 cm with soft wood shavings as bedding and maintained in well ventilated animal house with 12 hour light, 12 hour dark cycle. Water and standard pellet feed were given *ad-libitum* for the duration of the study. Procedure of this study was revealed by the ethical committee of Life Sciences, University of Benin, and ethical number LS020172 was issued by the committee.

### 2.5. Acute Toxicity Study

Twelve mice (12) were randomly divided into three groups (n=3) and each group was kept in a separate cage. The first group was given 10 mg/kg of the extract, the second group given with 100 mg/kg of the extract, third group received 1000 mg/kg of the extract and the control group. Four to Twenty four hours later after no death was recorded, phase 2 of the protocol involved three (3) mice administered with 1500, 2900 and 5000 mg/kg of the extract. All were observed after 24 hours all through 14 days using a method described by Lorke[11].

### 2.6. Sub-chronic Toxicity Study

Two four albino mice of male sex weighing 20 to 25 g were randomly divided into four (4) groups (n=6), each group was put in a separate cage and labelled appropriately. There were administered with graded doses of oral route of the extracts daily for 28. Graded doses of the extract (400, 800 and 1200 mg/kg body weight acetone extract of ELNA and the control group received 0.5 ml/kg body weight of distilled water. Mice were weighed every weekly for the duration of twenty eight days. Animals were anaesthetized and sacrificed, Blood sample was collected via cardiac puncture into EDTA bottle. Organs weight was recorded.

### 2.7. Acetic acid-induced pain

Twenty five albino mice weighing 13-25 g were randomly selected into groups of five (5) (n=5) after acclimatization. Two methods were adopted to induce pain in mice described by Nakamura *et al.*[12]. The writhing response was elicited by intraperitoneal injection of 0.75 % acetic acid at 0.1 ml/10 g body weight. Test substances and control vehicle were intraperitoneally injected into the mice 30 min before acetic acid and the number of writhes was observed and recorded for 15 min beginning from 5 min onset of acetic acid injection.

### Hot plate induced pain

Twenty five animals were randomly divided into five groups (n=5). Including untreated control, 100 mg/kg of aspirin and graded doses of acetone extract at 400, 800 and 1200 mg/kg

body weight). Treatment was administered across the groups 16 hr after fasted and were introduced on hot plate (maintained at 55°C) after acclimatization. Reaction time is characterized by jumping off or licking of paws) to determine thermal stimuli triggering central pain was noted at 30, 60, 90 and 120 min post extract administration. The mean of the latencies of the animals on the hot plate was determined [13].

## 2.8. Egg Albumin induced inflammation

The study was determined by adopting the techniques described by Winter *et al.* [14]; Akah and Nwabe [15]. Twenty five Swiss mice were randomly grouped into five (n=5) and treated as follows: Negative control administered with distilled water alone. Reference drug given 10 mg/kg body weight Indomethacin, test groups received graded doses of 400, 800 and 1200 mg/kg of acetone extract of ELNA. Oedema was induced by injecting mice with 0.05 ml of new raw egg albumin in the left hind paw after 30 min of post drug administration. Oedema size was evaluated by using Vernier's Calliper, with results taken at 30 minutes intervals (0, 30, 60, 90, 120 min) after albumin administration.

## 2.9. Statistical Analysis

Data were analysed using one-way analysis of variances (ANOVA). Group means were compared using turkey multiple comparison test and dunett post-test using a Graph pad prism instant software. Values were represented as mean standard error of mean  $\pm$  standard error of mean ( $P < 0.05$ ) were considered significant.

## 3. Results and Discussion

The aqueous and acetone extract showed a significant ( $p < 0.05$ ) dose dependent activity, but the aqueous extract had better activity than the acetone extract.

After 48 hours, no deaths or signs of toxicity was observed in the mice treated with the different doses (10, 100, 1000, 1500, 2900, 5000 mg/kg) of both aqueous and acetone extract of ELNA as show in Table 1. Throughout the duration of the experiment, the mice did not show any observable signs of toxicity or morbidity as they looked bright, were feeding well and their faeces looked normal.

Tables 2 shows the changes in body weight of mice treated with different doses (400 mg/kg, 800 mg/kg and 1200 mg/kg) of acetone extract of ELNA. Within each group there was no significant ( $p < 0.05$ ) increase in body weight for both extract.

There was no significant ( $p < 0.05$ ) difference in the percentage weight gain of the organs (heart, lungs, kidneys, liver and spleen) of the mice in acetone extracts of ELNA when compared to the control (Tables 3).

The acetone extracts of ELNA significantly ( $p < 0.05$ ) decreased the number of acetic acid induced writhes in mice at highest dose of 1200 mg/kg. At lower doses of the extracts showed no significant activity ( $p > 0.05$ ). The activity was also noted to be dose dependent (Table 4).

Table 6 revealed that 1200 mg/kg of acetone extract increased latency period significantly ( $p < 0.05$ ) at 60 mins. The effect is in dose dependent increase of the latency period for each time with no significant ( $p > 0.05$ ).

The results show that acetone extract caused inhibition of egg albumin induced oedema in mice over a period of 3 hrs. These effects were dose and time dependent for all doses but showed significant different ( $P < 0.05$ ) at doses of 1200 mg/kg of acetone extract and indomethacin respectively at 120, 150 and 180 min (Tables 7).




Figure 1: Effect of acetone extract of ELNA on DPPH scavenging activity

Table 1: Effect of acute toxicity of acetone extract of ELNA in mice

| Treatment | Dose (mg/kg) | No of mortality | No of Mortality and adverse effect |
|-----------|--------------|-----------------|------------------------------------|
| ELNA      | 10           | 0/3             | Absent                             |
| ELNA      | 100          | 0/3             | Absent                             |
| ELNA      | 1000         | 0/3             | Absent                             |
| ELNA      | 1500         | 0/1             | Absent                             |
| ELNA      | 2900         | 0/1             | Absent                             |
| ELNA      | 5000         | 0/1             | Absent                             |
| Control   | Dw           | 0/3             | Absent                             |

DW= distilled water

Table 2: Effect of acetone extract of ELNA on body weight of mice.

| Treatment/dosesmg/kg |      | Means $\pm$ SEM body weight of mice (g) |                  |                  |                  |
|----------------------|------|---|------------------|------------------|------------------|
|                      |      | Day 1                                   | Day 10           | Day 19           | Day 28           |
| Control              | Dw   | 21.00 $\pm$ 2.85                        | 20.83 $\pm$ 2.88 | 21.50 $\pm$ 2.94 | 21.67 $\pm$ 2.99 |
| ELNA                 | 400  | 21.50 $\pm$ 2.47                        | 19.33 $\pm$ 2.31 | 19.17 $\pm$ 2.27 | 19.17 $\pm$ 2.27 |
| ELNA                 | 800  | 21.17 $\pm$ 2.39                        | 19.00 $\pm$ 2.07 | 20.00 $\pm$ 1.98 | 19.83 $\pm$ 1.54 |
| ELNA                 | 1200 | 21.17 $\pm$ 2.24                        | 20.33 $\pm$ 1.76 | 20.17 $\pm$ 1.87 | 18.83 $\pm$ 1.92 |

Mean $\pm$ SEM;  $p > 0.05$ , n=5.

Table 3: Effect of ELNA acetone extract on percentage organ body weight of mice.

| Organs/Treatment | Means $\pm$ SEM Control Dw | Means $\pm$ SEM 400 mg/kg ELNA | Means $\pm$ SEM 800 mg/kg ELNA | Means $\pm$ SEM 1200 mg/kg ELNA |
|------------------|----------------------------|--------------------------------|--------------------------------|---------------------------------|
| Heart            | 0.52 $\pm$ 0.61            | 0.94 $\pm$ 0.19                | 0.53 $\pm$ 0.06                | 0.55 $\pm$ 0.04                 |
| Lungs            | 1.09 $\pm$ 0.14            | 0.91 $\pm$ 0.05                | 0.10 $\pm$ 0.06                | 0.95 $\pm$ 0.03                 |
| Kidney           | 1.42 $\pm$ 0.13            | 1.57 $\pm$ 0.13                | 1.67 $\pm$ 0.23                | 1.38 $\pm$ 0.09                 |
| Liver            | 4.93 $\pm$ 0.39            | 3.72 $\pm$ 0.94                | 5.77 $\pm$ 0.23                | 5.00 $\pm$ 0.17                 |
| Spleen           | 0.60 $\pm$ 0.04            | 0.90 $\pm$ 0.11                | 0.65 $\pm$ 0.10                | 0.56 $\pm$ 0.07                 |

Mean $\pm$ SEM;  $P > 0.05$ . n=5.

Table 4: Effect of acetone extract of ELNA on the haematological parameters of mice.

| Parameters                    | Means $\pm$ SEM Control alone | Means $\pm$ SEM 400 mg/kg ELNA | Means $\pm$ SEM 800 mg/kg ELNA | Means $\pm$ SEM 1200 mg/kg ELNA |
|-------------------------------|-------------------------------|--------------------------------|--------------------------------|---------------------------------|
| WBC $\times 10^3/\mu\text{l}$ | 5.15 $\pm$ 0.92               | 9.900 $\pm$ 1.42               | 7.62 $\pm$ 2.08                | 9.24 $\pm$ 0.88                 |
| RBC $\times 10^6/\mu\text{l}$ | 8.87 $\pm$ 0.50               | 9.320 $\pm$ 0.39               | 8.58 $\pm$ 0.44                | 10.07 $\pm$ 0.07                |
| HB g/dl                       | 14.30 $\pm$ 0.61              | 13.220 $\pm$ 0.50              | 12.93 $\pm$ 0.61               | 14.68 $\pm$ 0.39                |
| PCV %                         | 47.75 $\pm$ 1.71              | 48.020 $\pm$ 1.47              | 43.37 $\pm$ 2.29               | 52.12 $\pm$ 0.67                |

|                          |              |                            |               |               |
|--------------------------|--------------|----------------------------|---------------|---------------|
| MCV fl                   | 56.37±5.56   | 52.200±0.67                | 50.57±0.44    | 51.53±0.59    |
| MCH pg                   | 17.03±0.47   | 15.400±0.18                | 15.12±0.28    | 14.07±0.40    |
| MCHC g/dl                | 29.98±0.47   | 29.450±0.26                | 29.88±0.41    | 28.50±0.76    |
| PLT ×10 <sup>6</sup> /μl | 216.80±97.54 | 907.20±148.10 <sup>b</sup> | 703.80±246.70 | 771.70±157.70 |
| LYM %                    | 64.48±8.12   | 61.800±8.99                | 85.36±2.79    | 75.44±4.48    |
| MXD %                    | 7.61±1.81    | 5.600±1.44                 | 3.04±0.53     | 6.56±1.87     |
| NEUT %                   | 2852±6.58    | 18.16±3.05                 | 11.20±2.63    | 32.40±8.00    |

n=5; means ±SEM; \**p*<0.05

Table 5: Effect of acetone and aqueous extract of ELNA on acetic acid induced writhes.

| Dose    | Doses mg/kg | Mean±SEM number of Writhes |
|---------|-------------|----------------------------|
| Control | Dw          | 47.60±1.89                 |
| ELNA    | 400         | 43.60±2.42                 |
| ELNA    | 800         | 34.80±4.21                 |
| ELNA    | 1200        | 29.60±5.71 <sup>b</sup>    |
| Aspirin | 100         | 24.80±1.24 <sup>c</sup>    |

Mean±SEM; <sup>b</sup>*p*<0.05; <sup>c</sup>*p*<0.01, n=5. Dw----- distilled water.

Table 6: Effect of ELNA acetone extract on hot plate induced nociceptive in mice.

| Mean±SEM latency period of the animals on the Hot Plate (seconds) |                  |                |                |                         |                   |
|---|------------------|----------------|----------------|-------------------------|-------------------|
| Time Interval (mins)  | Control Dw alone | ELNA 400 mg/kg | ELNA 800 mg/kg | ELNA 1200 mg/kg         | Aspirin 100 mg/kg |
| 30  | 12.15±0.75       | 13.10±0.60     | 13.38±2.00     | 14.90±1.12              | 12.32±1.24        |
| 60  | 11.60±1.10       | 9.84±1.56      | 15.94±1.14     | 16.24±1.36 <sup>b</sup> | 15.14±0.75        |
| 90  | 9.76±1.02        | 9.96±1.24      | 10.64±1.20     | 11.68±1.34              | 13.78±1.27        |
| 120   | 10.32±1.66       | 9.78±0.70      | 11.98±0.96     | 9.78±0.70               | 9.20±1.70         |

Mean±SEM; \**P*<0.05, n=5.

Table 7: Effect of acetone extract of ELNA on egg albumin induced right hind paw Inflammation in mice.

| Time (mins) | Mean±SEM Control Dw alone | Mean±SEM 400 mg/kg ELNA | Mean±SEM 800 mg/kg ELNA | Mean±SEM 1200 mg/kg ELNA | Mean±SEM 100 mg/kg Indomethacin |
|-------------|---------------------------|-------------------------|-------------------------|--------------------------|---------------------------------|
| 30          | 0.10±0.01                 | 0.13±0.08               | 0.12±0.01               | 0.13±0.01                | 0.12±0.01                       |
| 60          | 0.12±0.01                 | 0.11±0.18               | 0.11±0.02               | 0.10±0.01                | 0.08±0.01                       |
| 90          | 0.12±0.01                 | 0.10±0.01               | 0.09±0.01               | 0.09±0.00                | 0.08±0.01                       |
| 120         | 0.11±0.01                 | 0.09±0.01               | 0.09±0.00               | 0.10±0.02 <sup>a</sup>   | 0.06±0.06 <sup>c</sup>          |
| 150         | 0.10±0.01                 | 0.08±0.01               | 0.09±0.01               | 0.07±0.00 <sup>b</sup>   | 0.04±0.01 <sup>c</sup>          |
| 180         | 0.09±0.01                 | 0.08±0.02               | 0.07±0.01               | 0.04±0.01 <sup>b</sup>   | 0.01±0.00 <sup>c</sup>          |

Mean±SEM; <sup>a</sup>, <sup>b</sup>, <sup>c</sup>*P*<0.05, n=5.

Low levels antioxidant or inhibition of antioxidant enzymes causes oxidative stress and possible damage or kill cells [16]. Free radicals are also responsible for oxidative processes. This study showed DPPH scavenging property of acetone extract of ELNA in graded concentration, significantly inhibits oxidative and promote scavenging effect of the extract when compared with ascorbic acid (Figure 1). This study is in line with Siddhuraju and Becker [17] on antioxidant properties of various solvent extracts of total phenolic constituents from three different agro-climatic origins of drumstick tree *Moringa oleifera* leaves. *In-vitro* anti-oxidant evaluation of extract showed significant ( $p < 0.05$ ) increase in scavenging DPPH radical in concentration dependent manner, had a better inhibitory activity. This concurred with the work of Sies [18]; Singh *et al.* [19].

Acute toxicity study may serve as the basis for classification and labelling, provide initial information on the mode of toxic action of a substance, dose of a new compound, help in dose determination in animal studies and determine  $LD_{50}$  that provide many indices of potential types of drug activity [20]. Substances with  $LD_{50}$  values higher than 5000 mg/kg by oral route are regarded as being safe or practically less toxic. This is in line with the report of Adamu *et al.* [21] with the effect of Aqueous and methanol stem bark extract of *Maerua angolensis* on acute and sub-acute inflammations. Present acute toxicity study, showed absent mortality or signs of toxicity observed after 48 hours all through 14 days of a single and highest doses of acetone extract ELNA (Table 1), this indicated that the extract was safe with wide range of tolerance. Studies from Chainani-Wu [22]; Tripathy *et al.* [23]; Awodele *et al.* [24] carried out on acute toxicity of individual herbs that makes up the herbal formulation with wide safety margin ( $LD > 5000$  mg/kg) concurred with present study. These justified the reason why the formulation was also safe. General behaviour and body weight are one of the critical parameters for the evaluation of first signs of toxicity [25]. In the assessment of the sub-chronic toxicity study of ELNA extract, the body weight of the mice were taken at alternate days throughout 28 days. There was no appreciable weight gain or weight loss, as the body weight showed no significant different ( $p < 0.05$ ). The mean percentage organ body weight (heart, lungs, liver, kidneys, spleen) of the mice was not significant ( $p > 0.05$ ) across the tested groups of the extract after 28 days (Table 4), though a slight increase in the group administered with the extract. Haematological parameters were relatively stable and no significant increase or decrease in the extract, with slight increase ( $p < 0.05$ ) in 400 mg/kg of platelet count when compared with the control. Awodele *et al.* [24] reported a similar response on the toxicological evaluation of aqueous leaf extract of *Moringa oleifera* Lam. (Moringaceae).

Herbal therapist outlined ELNA usefulness as a potent anti-oxidant, anti-inflammatory and analgesic agent [26]. The analgesic activity of acetone extract of ELNA was demonstrated using acetic acid writhing (peripherally acting) and hot plate (centrally acting) model showed no significant different ( $p < 0.05$ ) when compared with the untreated control. Similar results was reported from Gene *et al.* [27] showed *Heterotheca inuloides* anti-inflammatory and analgesic effects. One of the possible mechanisms through which ELNA extract elicited its action via inhibition of the releases of prostaglandins PGE2 and PGF2 $\alpha$ , triggered by acetic acid. Study from Tanko *et al.* [28] showed similar nociceptive and anti-inflammatory mechanism of action derived from aqueous leaves extract of *Ocimum gratissimum* (Labiatae) in rodents. This is also implicated as a mediator of inflammation [29]. Increase level of prostaglandin within the peritoneal cavity enhanced inflammatory pain via increased capillary permeability [30]. Prostaglandin synthesis is possibly inhibited through peripheral mechanism of pain by ELNA extract as nociceptive and anti-inflammatory effect to reduce the number of writhes rendering analgesic effect when compared with the control (Table 5). The report of Ferdous *et al.* [31] showed the nociceptive activity of aqueous extract of *Ficus racemosa* Linn. The acetone extract of ELNA showed significant ( $p < 0.01$ ) decreased in writhes at (32.20 and 29.60) respectively specifically at 1200 mg/kg. Generally, acetone extract demonstrated a viable result in dose dependent effect. Hot plate model for assaying nociceptive effects of acetone extract of ELNA on central pain effective in against pain [32]. The acetone extract of ELNA showed a significance ( $p < 0.05$ ) decrease at 60 mins for 1200 mg/kg body weight as it aid in reducing mean heat latency period as shown in Table 6. Aspirin displayed analgesic effect with significant decrease in inflammation when compared with untreated control. Based on the effectiveness of the extract on hot plate test, a central mechanism of action could be implicated.

Inflammatory processes involved mediated chemicals such as prostaglandins, histamine and serotonin. Egg albumin-induced oedema from this present study results stimulated the release of histamine and serotonin [33]. The anti-inflammatory properties of acetone extract of ELNA elicited significant decrease ( $p < 0.05$ ) of paw edema at 1200 mg/kg for 120 mins, 150 mins and 180 mins respectively. The extracts under investigation showed significant reduction in inflammation triggered by fresh egg-albumin against the progressive increase in mice paw circumference of the untreated control. Graded doses of the extract suppressed the increase in mice paw oedema in a dose dependent manner two to three hours after inducing inflammation as shown in Table 7. This implies that the extracts will be useful in the management of inflammatory pain. Similar report of Wannang *et al.* [34] whose analgesic and anti-inflammatory

activity of the aqueous leaf extract of *Solanum nigrum* Linn (Solanaceae) in Rat.

#### 4. Conclusion

Acetone extract of ELNA possesses a wide range of biosafety with anti-oxidant properties, analgesic and anti-inflammatory, which validated the ethnomedicinal claim by herbal therapist that the formulation is safe with LD<sub>50</sub> greater than 5000 mg/kg. They were also safe after 28 days of oral administration because there was no residual effect of the prolonged administration.

#### Acknowledgements

Our earnest gratitude goes to Mr. Dialect and Mr. Ehigie of the Department of Haematology University of Benin Teaching Hospital, Mr. Odega Kelvin and Mrs. Queen Okoro of the Department of Morbid Anatomy, University of Benin Teaching Hospital the slides preparation and interpretation. Dr. Agu for the antioxidant assays in the Department of Medical Biochemistry, University of Benin.

#### Conflict of interests

No conflict of interest

#### References

1. Barnes J, Anderson LA, Phillipson JD. Herbal Medicine. 3rd Edition, Pharmaceutical Press, London. 2007, pp. 1-23.
2. Chandraprakash D, Swarnali D. Anti-diabetic herbal drugs and polyherbal formulation used for diabetes: A review. *Journal of Phytopharmacology*, 2013, 2(3); 44-51.
3. Gomase P, Shire SP, Nazim S, Choudhari AB. Development and evaluation of polyherbal formulation for anti-inflammatory activity. *Journal of Natural Product and Plant Resource*, 2011, 1; 85-90.
4. Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: a food plant with multiple medicinal uses. *Phytotherapy Research*, 2007, 21; 17-25.
5. Ghasi S, Nwobodo E, Ofili JO. Hypocholesterolemic effects of crude extract of leaf of *Moringa oleifera* Lam in high-fat diet fed wistar rats. *Journal of Ethnopharmacology*, 2000, 69; 21-25.
6. Duke JA. Root bark of *Moringa* contains alkaloid moringinine which acts as cardiac stimulant through its effect on sympathetic nervous system. *Handbook of Nuts*. CRC Press, USA. 2001, pp. 214-217.
7. Williamson M. Major Herbs of Ayurveda. Churchill Livingston publication, *Elsevier Science*, 2002, Ltd, 116p.
8. Warriar PK. Indian medicinal plants: A compendium of 500 species. Arya V. S. (Ed.), Orient Publication, France, 1997, 4; 202.
9. Sahoo S, Mishra SK, Panda PK, Mishra SR, Ellaiah P, Dash SK. Antimycotic potential of *Crataeva religiosa* against some selected fungal pathogens. *Acta Poloniae Pharmaceutica - Drug Research*, 2008, 65(2); 245-247.
10. Georgia K, Victoria Y, Ashok K. Cardiovascular protection by curcumin; molecular aspect. *Indian Journal of Biochemistry and Biophysics*. 2012, 49; 306-315.
11. Lorke D. A new approach to practical acute toxicity testing. *Archives of Toxicology*, 1983, 54; 275-287.
12. Nakamura H, Shimoda A, Ishii K, Kadokawa T. Central and peripheral analgesic action of non-acidic non-steroidal anti-inflammatory drugs in mice and rats. *Archives of International Pharmacodynamics*, 1986, 282; 16–25.
13. Eddy NB, Touchberry CF, Lieberman IE. Synthetic analgesics, a methadone isomers and derivatives. *Journal of Pharmacology and Experimental Therapeutics*, 1950, 98; 121-137.
14. Winter ER, Risley EA, Nuss GV. Anti-inflammatory and anti-pyretic activities of indomethacin. *Journal of pharmacology*, 1963, 141; 369-376.
15. Akah P, Nwabie AI. Evaluation of Nigerian traditional medicines: Plants used for rheumatic disorder. *Journal of Ethnopharmacology*, 1994, 42; 179-182.
16. Iwu MM. Handbook of African Medicinal Plants. CRC Press, London, 1993, pp. 183-184.
17. Siddhuraju P, Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *Journal of Agricultural and Food Chemistry*. 2003, 51; 2144-2155.
18. Sies H. Oxidative stress: Oxidant and anti-oxidant. *Experimental Physiology*, 1997, 82(2); 291–295.
19. Singh BN, Singh BR, Singh RL, Prakash D, Dhakarey R, Upadhyay G, Singh HB. Oxidative DNA damage protective activity, antioxidant and anti-quorum sensing potentials of *Moringa oleifera*. *Food Chemistry and Toxicology*. 2009, 47; 1109-1116.
20. Ukwuani AN, Abubakar MG, Hassan SW, Agaie BM. Toxicological studies of hydromethanolic leaves extract of *Grewia crenata*. *International Journal of Pharmaceutical Science and Drug Research*, 2012, 4; 245–249.
21. Adamu A, Abdurrahman EM, Ibrahim H, Abubakar MS, Magaji MG, Yaro AH. Effect of Aqueous methanolic stem of bark extract of *Maerua angolensis* acute and sub-acute inflammations. *Nigerian Journal of Pharmaceutical Sciences*, 2007, 6(2); 1-6.
22. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: a component of tumeric (*Curcuma longa*). *Journal of Alternative and Complementary Medicine*, 2003, 9(1); 161–168.
23. Tripathy SA, Ashaa MD. Pradhana acute and chronic anti-inflammatory evaluation of *Crataeva religiosa* in rats. *International Journal of Pharmacy and Technology*, 2010, 2(4); 1270-1279.
24. Awodele O, Oreagba IA, Odoma S, da Silva JA, Osunkalu VO. Toxicological evaluation of the aqueous leaf extract of *Moringa oleifera* Lam. (Moringaceae). *Journal of Ethnopharmacology*, 2012, 139(2); 330-336.
25. Sireeratawong S, Urarat N, Supaporn V, Natthakarn C, Supachai S, Pennapa S, Pornthip T, Parunkul AC, Kanjana J. Acute and subchronic toxicity study of tud-rak-ka-sai-puu recipe in rats. *African Journal of Traditional, Complementary and Alternative Medicine*, 2013, 10(1); 142-148.
26. Park JJ, Kang M, Shin S, Choi E, Kwon S, Wee H, Nam B, Kaptchuk TJ. Unexplained infertility treated with acupuncture and herbal medicine in Korea. *Journal of Alternative and. Complementary Medicine*, 2010, 16(2); 193-198.

27. Gene RM, Segura L, Adzet T, Marin E, Iglesias J. *Heterotheca inuloides*: Anti-inflammatory and analgesic effects. *Journal of Ethnopharmacology*, 1998, 60; 157-162.
28. Tanko Y, Magaji G, Yerima M, Magaji A, Mohammed A. Antinociceptive and anti-inflammatory activities of aqueous leaves extract of *Ocimum gratissimum* (Labiata) in rodents. *African Journal of Traditional, Complimentary and Alternative Medicines*, 2008, 5(2); 141-146.
29. Santosh S, Sohan S, Anupama S, Devanand S, Manohar JP. Anti-inflammatory activity of an isolated flavonoid fraction from *Celosia argentea* Linn. *Journal of Medicinal Plants Research*, 2008, 2(3); 52-54.
30. Zakaria Z, Sulaiman M, Gopalan H, Ghani Z, Raden-Mohd R, Jais M, Abdullah F. Anti-nociceptive and anti-inflammatory properties of *Corchorus capsularis* leaves chloroform extract in experimental animals models. *Journal of the Pharmaceutical Society of Japan*, 2007, 127; 359-365.
31. Ferdous M, Rouf R, Shilpi J, Uddin S. Antinociceptive activity of aqueous extract of *Ficus racemosa* Linn. *Oriental Pharmacy and Experimental Medicine*, 2008, 8; 93-96.
32. Kumar Y, Vanitha K, Prakash K, Swetha M. A new facile and sensitive method for the estimation of tapentadol. *International Journal of Pharmaceutical Sciences and Drug Research*, 2014, 6(1); 82-84.
33. Akindele AJ, Adeyemi OO. Anti-inflammatory activity of the aqueous leaf extract of *Byrsocarpus coccineus*. *Fitoterapia*, 2007, 78; 25-28.
34. Wannang NN, Anuka JA, Kwanashie HO, Auta A. Analgesic and anti-inflammatory activity of the aqueous leaf extract of *Solanum nigrum* Linn (Solanaceae) in Rat. *Nigerian Journal of Pharmaceutical Research*, 2006, 5; 9-13.

### Recommended Citation

MacDonald I, Onuigbo C U, Gabriel B O. In-vitro Antioxidant activity, Biosafety, Nociceptive and Anti-inflammatory potential of Acetone Polyherbal (ELNA) Extract in mice. *Algerian Journal of Biosciences*. 2021, 02;02:059-066.

doi : <http://dx.doi.org/10.5281/zenodo.5810172>

Or

### Recommended Citation

MacDonald I, Onuigbo C U, Gabriel B O. In-vitro Antioxidant activity, Biosafety, Nociceptive and Anti-inflammatory potential of Acetone Polyherbal (ELNA) Extract in mice. *Alger. j. biosciences*. 2021, 02;02:059-066.

doi : <http://dx.doi.org/10.5281/zenodo.5810172>

