

How to Cite:

Chauhan, U., & Chauhan, E. S. (2022). Review on nutritional composition, bioactive compounds and therapeutic properties of *Hippophae rhamnoides* L. (Sea Buckthorn). *International Journal of Health Sciences*, 6(S3), 11620–11632. <https://doi.org/10.53730/ijhs.v6nS3.8773>

Review on nutritional composition, bioactive compounds and therapeutic properties of *Hippophae rhamnoides* L. (Sea Buckthorn)

Urvashi Chauhan

Junior Research Fellow, Dept. of Food Science and Nutrition, Banasthali Vidyapith, Rajasthan, India

Ekta Singh Chauhan*

Associate Professor, Dept. of Food Science and Nutrition, Banasthali Vidyapith, Rajasthan, India

Abstract---*Hippophae rhamnoides* L. (Sea Buckthorn) is a valuable multifunction plant largely distributed in Europe and Asia. In particular, the berries of sea buckthorn have been widely used as a food source like jam, juices, etc. The berry and seed of this plant are the main sources for its therapeutic and nutritional values. Thus, the plant has been extensively investigated for both of its nutritional and bioactive constituents. In general, the berries of this plant are rich in vitamins (C, E and K); while the ascorbic acid has been reported to be the primary vitamin, contributing to the acidic nature of the berries. Sea buckthorn berries also contain good amount of minerals like calcium and iron. On the other hand, the seed of sea buckthorn offers an excellent source of polyunsaturated fatty acids like palmitic acid, oleic acid and palmitoleic acid. Significant bioactive constituents in sea buckthorn including sitosterol, salicylic acid, triterpenoid, catechin, gallic acid, epigallocatechin and carotenoids help in combating many diseases like cardiovascular diseases (angina), inflammation, cancer, gastric ulcer and epidermis issues. This review compiled the applications of sea buckthorn including a database of its nutritional and bioactive compounds which ultimately contribute towards the health promoting properties. As a result, these beneficial attributes have made sea buckthorn desirable source for medicinal as well as for cosmetic products.

Keywords---sitosterol, palmitoleic acid, catechin, polyphenols, carotenoids.

Introduction

Hippophae rhamnoides L. commonly called sea buckthorn, sanddorn, seaberry or Siberian pineapple. It's a deciduous spiny shrub (Fig. 1) belongs to the Elaeagnaceae family (Singh et al, 2019). Sea buckthorn berries (Fig. 2) are of orange or yellow colour, circular shaped and about 3 to 8 mm in size. The detailed taxonomy of the plant is outlined in Table 1. The sea buckthorn plant naturally grows in north-western Europe, Central Asia and the northern Himalayas (Teleszko et al, 2015). While in India, it found in the zone of Ladakh, Himachal Pradesh, Uttarakhand and Sikkim. Sea buckthorn berries are mostly utilized for the local production of liquors, syrups and jams (Rop et al, 2014). All parts of this plant (especially berries and seeds) are rich in nutrients and bioactive constituents which are important for health enhancing factors. Therefore, sea buckthorn plant has been used traditionally as a remedy in many nations such as China, India, Romania, etc.



Fig 1: Sea buckthorn plant



Fig 2: Sea buckthorn berries

Table 1
Taxonomy of *Hippophae rhamnoides* L.

Kingdom	Plantae
Class	Dicotyledonae
Order	Elaeagnales
Family	Elaeagnaceae
Genus	<i>Hippophae</i>
Species	<i>H. rhamnoides</i>

Latest medicinal researches of sea buckthorn identified numerous health-promoting applications in vitro as well as in vivo (Wani et al, 2016). Therefore,

this review describes the valuable properties of *Hippophae rhamnoides* L. including its nutritional composition, bioactive compounds and health benefits.

Nutritional Properties

The nutritional attributes of sea buckthorn depend on the fruit's size, maturity, geographical region, species, climate and the method of extraction (Leskinen et al, 2010). The nutritional profile of sea buckthorn berries is briefly described in Table 2 and Table 3.

Carbohydrates

Carbohydrates are considered to be the main component in sea buckthorn berries. CHO present in sea buckthorn berries are majorly disaccharides and monosaccharides. The total amount of CHO in fresh weight is 0.48%–2.87% (Tkacz et al, 2019; Wu et al, 2016). However, the main monosaccharides are xylose, glucose, rhamnose and fructose, while the utmost representative disaccharides are sucrose and maltose.

Protein

According to Nazir and Bashir (2017), the protein in sea buckthorn berries is around 0.4% to 2.5% in fresh weight as well as the amount of free amino acid ranges in fresh weight vary from 0.77% to 2.19%. Hence, eighteen free amino acids identified in sea buckthorn and eight of these (phenylalanine, threonine, isoleucine, tryptophan, leucine, valine, lysine, and methionine) are important for health (Stobdan et al, 2013). Yet, the leading amino acid in sea buckthorn fruit pips are glutamic acid along with aspartic acid (Trineeva et al, 2015).

Lipids

Morphologically, sea buckthorn berries contain skin (8% w/w), pulp (68% w/w) and seeds (23% w/w) (García, 2019) and all consist good amount of oil. The lipid in sea buckthorn on fresh weight ranges from 1.2%–7.8%. In a study (Craciun, 2018) identified the amount of saturated, monounsaturated, and polyunsaturated fatty acids is 13.70%–42.68%, 40.73%–60.37% and 3.70%–24.62% (Craciun, 2018). However, (Dulf, 2012) evaluated the dominating fatty acids in berries pulp oil were palmitoleic (11-27%), palmitic (23-40%) and oleic (20-53%).

Table 2
Nutrient Content in Sea buckthorn Berries

Compounds			Content
Carbohydrates (%)			0.48–2.87
Protein (%)			0.4–2.5
	Free amino acid		0.77–2.19
		Aspartic acid *	0.43–55.68
		Glutamic acid*	11.76–16.48

Lipids (%)			1.20–7.8
	Fatty acid (%lipid)		8.8–11.1
	Saturated fatty acids (% Total fatty acid)		13.70–42.68
	Monounsaturated fatty acids (% Total fatty acid)		40.73–60.37
	Polyunsaturated fatty acids (% Total fatty acid)		3.70–24.62
		Palmitic acid (16:0)	23-40
		Oleic acid (18:1n-9)	20-53
		Palmitoleic (16:1n-7)	11-27

*Essential Amino Acid and % lipid mean data based on sea buckthorn oil extract

Vitamins and Minerals

Sea buckthorn is a good source of vitamins like thiamine, riboflavin, pyridoxine and folic (Lu et al, 2019). However, the other vitamins rich in sea buckthorn are vitamin C, E and K. According to Pop et. al (2015), amount of α -tocopherol in sea buckthorn in fresh weight is between 43-223 mg/kg in fresh. In a very recent study Wang et. al (2021) observed that in fresh weight sea buckthorn berries vitamin K is around 90-120mg/kg (Table 3). Moreover, vitamin C content in sea buckthorn is higher than the other fruits. The vitamin C content in sea buckthorn has an average of 7950 mg/kg in fresh weight (Hussain et al, 2014). The brief detail of vitamins and minerals in sea buckthorn berries are mentioned in Table 3. Additionally, twenty-four chemical elements are available in the juice of sea buckthorn such as, calcium, manganese, iron, boron, etc (Hussain et al, 2014). The calcium present in dried sea buckthorn is 724mg/kg similarly, the content in zinc (22.3mg/kg) as well as in copper (8.3 mg/kg) (Nour et al, 2021). Vaitkeviciene et. al (2019) studied there have also been other minerals available in sea buckthorn like, Iron (282 mg/kg DW), Manganese (16.30 mg/kg DW) and Boron (13mg/kg DW).

Table 3
Vitamins and Minerals Content in Sea buckthorn Berries

Compounds		Content
Vitamins FW (mg/kg)	Vitamin E	43-223
	Vitamin K	90-120
	Vitamin C	7950
Minerals DW (mg/kg)	Calcium (Ca)	724
	Zinc (Zn)	22.3
	Copper (Cu)	8.3
	Iron (Fe)	282
	Boron (B)	13

	Manganese (Mn)	16.30
--	----------------	-------

DW (Dry Weight); FW (Fresh Weight)

Bioactive Compounds

Bioactive constituents are the non-nutrient plant compounds which include health promoting properties as shown in Table 5. All the parts of sea buckthorn are rich source of large number of phytochemicals and antioxidants (Christaki, 2012).

Phytosterols

Sea buckthorn consist seventeen phytosterols. The phytosterol content in fresh weight sea buckthorn found to be 310–520 mg/kg (Hussain et al, 2014) and even higher than soybean oil. The dominating phytosterol content is sitosterol, which is between 160–330 mg/kg in fresh weight. The other phytosterols are mostly stigmastanol, campesterol, α -amyrin, and stigmastadienol.

Polyphenols

Recently Criste et. al (2020) identified that the phenolic compounds present in sea buckthorn berries and leaves, were aglycones and derivatives of cinnamic or quercetin acid. Total phenolic contents in sea buckthorn recalculated as gallic acid equivalents ranges from 32.93 to 1417 mg/ 100 g in fresh weight (Ivanisova et al, 2017; Rop et al, 2014). Flavonoids, tannins and phenolic acids are the major polyphenols in sea buckthorn (Guo et al, 2017).

- **Flavonoids**

Epicatechin, catechin, kaempferol and isorhamnetin are the major flavonoids in sea buckthorn (Fan et al, 2012). The flavonoid level in sea buckthorn berries is vary from 1680 - 8590 mg/kg FW (Heinaaho and Julkunen Tiitto, 2011) and a lot times higher than the concentration present in other high-flavonoid plants like cornelian cherry, European blackberry, , hawthorn, dog rose or blackthorn.

- **Tannins**

Tannins in sea buckthorn are of two types which are condensed and hydrolysable tannins. These are available in larger amount in sea buckthorn roots, green berries, seeds, stems and flowers (Michel et al, 2012).

- **Phenolic acids**

In the plant's immune system, phenolic acids considered to be the most essential compound (Daško et al, 2012). In sea buckthorn, gallic acid in free form is leading with variable concentration appeared in berries (16.9 mg/kg) and leaves (79 mg/kg). While, the other phenolic acids of sea buckthorn observed in minor concentration of ferulic acid, p-coumaric acid and caffeic acid (Bittová et al, 2014). Salicylic acid is the main phenolic acid present in the berries sea buckthorn and the content is between 21- 47 mg/kg depends on the variety. Aside from salicylic acid, sea buckthorn berries also contain gallic acid in minor content (1 - 4.6 mg/kg).

Carotenoids

Carotenoids are mostly found in the soft parts of the fruit, which give them a distinctive orange-yellowish colour. In sea buckthorn berries 15–55% of all compounds of this group is β -carotene (Teleszko et al, 2015). Since, sea buckthorn berries are high in carotenoids as well as up to eighteen different carotenoids observed in the oil of sea buckthorn with the carotenoids owning provitamin A activity (sintexanthin, β -cryptoxanthin, β -zeacarotene, β -carotene and γ -carotene) along with lutein. The oil of sea buckthorn has been a primary subject in terms for health benefits, due to its higher concentration of carotenoids especially zeaxanthin (Preedy, 2014). Pop et. al (2015) identified the total concentration of carotenoids in sea buckthorn which was cultivated in Romania, only on average 172 mg/kg in fresh weight which is equivalent to around 860 mg/kg dry weight of berries. While observing various species, Ranjith et. al (2016) observed that *H. rhamnoides* berries made higher concentration of carotenoids as compared to *H. salicifolia* and *H. tibetana*.

Table 4
Bioactive compounds in sea buckthorn

Compounds		Content
Total Phytosterol (mg/kg)		310–520
	Sitosterol (mg/kg)	160–330
Total Phenolic compounds (mg GAE/100 g)		32.93–1417
	Flavonoids (mg/kg)	1680–8590
	Phenolic acids (mg/ /kg)	16.9
Total Carotenoids (mg/kg)		172

Therapeutic Properties

Nutritional and bioactive compounds are rich in sea buckthorn as they may affect inflammatory disorders, carcinoma, CVD, gastric ulcer and epidermis problems.

Anti-inflammatory

Chinese and Tibetan natives have been widely using sea buckthorn as a medicine for treating inflammatory disorders. Many studies have showed that it's capable of promoting immunomodulatory and anti-inflammatory activities (Suryakumar and Gupta, 2011). The chief bioactive constituents in sea buckthorn which are responsible for curing inflammatory disorders are antioxidants like carotenoids, flavonoids, vitamin E and ascorbic acid (Sayegh et al, 2014). In addition, the oil of sea buckthorn seed is rich in omega-3 fatty acid which helps in promoting anti-inflammatory property as well as insulin-sensitizing (Fatima et al, 2012). A clinical test was performed by Pentelescu et. al (2014) targeting at acute gingivitis which implied that sea buckthorn maybe used for treating gingivitis through increased immune response and improved gingival tissue. A few presences of mitogens in sea buckthorn activate lymphocyte-proliferation (Patel et al, 2012).

Anti-carcinogenic

Polyphenols in sea buckthorn can prohibit the polyphenols mutation of carcinoma cells, like human breast cancer cells as well as hepatocellular carcinogenic cells (Wang et al, 2021). Especially, berries of sea buckthorn having a potential of anticancer activity. Chemo preventive potential of sea buckthorn was evaluated in mice by Padmavathi et. al (2005) and the results showed that *Hippophae* berry extract stimulated activities of both phase II and antioxidant enzymes in the mouse liver. Studies have also been found that sea buckthorn decreased tumor incidence of skin and forestomach papilloma genesis in mice. Yasukawa et. al (2009) revealed 3 phenolic compounds, epigallocatechin, gallic catechin, triterpenoid and catechin, ursolic acid from the active fraction of the 70% ethanol extract displayed significant anti-cancer activity.

Cardiovascular risk reduction

Reduction of CVD risk is one of the most essential health beneficial activities of sea buckthorn. Since, sea buckthorn is rich in flavonoids like aglycones, quercetin and isorhamnetin, many researchers have tested the effect of various edible parts of sea buckthorn like its berry, flavonoid extracts, oil and its derivatives for preventing the cardiovascular diseases (Xu et al, 2011). Many researches (Koyama et al, 2009; Larmo et al, 2009) have identified the health promoting effects of plant sterols, polyunsaturated fatty acids, antioxidants, and flavonoids which were available in berry's juice, leaves and oil of sea buckthorn.

Flavonoids in sea buckthorn flavonoids improve function of by lowering blood viscosity and scavenging free radicals. There are many other health beneficial effects of sea buckthorn on CVD as they help in lowering the blood sugar level, cholesterol level and pressure level (Xu et al, 2011). It contains good antioxidant activities that's why it could be used for improving blood pressure along with lipids, in order to prevent and treat CVD symptoms like angina, to reduce free radical levels and prevent atheroma too (Zakynthinos and Varzaka, 2015). Numerous studies have revealed that particularly oil and juice of sea buckthorn have a positive effect on cardiovascular functions as well as lowering the cholesterol levels (Bal et al, 2011). Therapeutic properties of sea buckthorn on CVD have been showed in various animal and clinical trials. For instance, patients took 10 mg sea buckthorn flavonoids extract thrice a day for six weeks revealed decreased cholesterol and improved cardiovascular system (Zakynthinos and Varzaka, 2015).

Gastroprotective

Berries and seeds of sea buckthorn have indicated a potential importance in treating the mucous membranes which include gastro-intestinal disorders, ulcers as well as vaginal problems (Bal et al, 2011). The anti-ulcerogenic effect of a hexane extract from *Hippophae rhamnoides* was tested on indomethacin and stress induced ulcer models. As a result, hexane extract from sea buckthorn was found to be active in preventing gastric injury (Suleyman et al, 2001).

Epidermis health

Traditional utilization of sea buckthorn oil helps in the healing of skin injuries and skin conditions which is also agreed with the data of modern clinical and preclinical studies (Suryakumar and Gupta, 2011). Sea buckthorn oil is effective in preventing and treating many skin diseases like Sjogren's syndrome, atopic dermatitis, and elasticity of skin because of the presence of unsaturated fatty acids, phytosterols tocopherols and carotenoids (Larmo et al, 2014). In their study (Marsiñach and Cuenca, 2019) revealed sea buckthorn oil may had clinical properties for skin conditions like atopic dermatitis, dry eye and acne due to the presence of omega-6 fatty acids (γ -Linolenic acid and Linoleic acid). Studies have been indicated that omega-7 (palmitoleic acid) was also available in the oil of sea buckthorn which is beneficial for wound healing and regeneration of skin in the epidermis (Zielińska and Nowak, 2017). Furthermore, sea buckthorn can also facilitate cell migration, eliminate damaged protein, remodel the granulation tissue and regulate the activity of some growth factors through increasing the expression of matrix metalloproteinase in all phases of wound recovery process (Edraki et al, 2014). Sea buckthorn oil is extensively used alone or in many other forms like topically applied for burns, infections, ulcerations and scalds (Chandra et al, 2018).

Table 5
Summary of Therapeutic Properties

Compounds	Health Benefits
Flavonoids, vitamin C and vitamin E	Promotes healthy immune system function.
	Supports healthy cardiovascular function as well as lowering the blood sugar and blood pressure levels.
	Assists in the process of healthy cellular rejuvenation.
	Fights cell-damaging free radicals.
Omega 3, 6 & 7	Promotes healthy skin.
	Supports health of cardiovascular system.
Carotenoids	Supports anti- inflammatory activity.
	Zeaxanthin promotes health of eyes.
Phenolic compounds (catechin, gallic acid, epigallocatechin and triterpenoid)	Promotes anti- carcinogenic properties.

Toxicity and Safety

Sub chronic toxicity and efficacy study of sea buckthorn oil revealed no significant changes in many haematological, physiological, biochemical (creatinine, uric acid, urea, ALT, ALP and AST) and histopathological parameters in 3 groups of New Zealand white rabbits (0.5, 1, and 1.5 ml/kg body weight) with no mortality in any of the treatment groups (Rashid et al, 2011). Zhao et. al (2017) identified that sea buckthorn oil extracted from whole berry through a supercritical carbon dioxide method during 90-days toxicity study in rats indicated no adverse effects after oral administration. Additionally, in a very latest

study by Wen et. al (2020) pregnant rats were treated with 4.68, 2.34, and 1.17 g/kg oil from gestation 7 to 16 days and no treatment-related maternal toxicity or embryotoxicity was found. However, the info about safety and toxicity of sea buckthorn or its derivatives for dietary consumption by humans is still very lacking. Overall, the current results implied that sea buckthorn is not toxic for potential human consumption in food or as a dietary supplement.

Conclusion

Sea buckthorn plant has been extensively used as a traditional remedy in many nations like China, Nepal, India, Pakistan, Romania, Germany, Russia, etc. The berries of sea buckthorn offer good amount of nutrients such as essential amino acid (aspartic and glutamic acid), monounsaturated fatty acid, vitamins (C, E, K) and minerals (calcium, iron, zinc). Also, the major bioactive compounds present in sea buckthorn are sitosterol, salicylic acid, carotenoids, catechin, kaempferol, isorhamnetin, etc. Due to its excellent source of nutrients and bioactive constituents the plant has various therapeutic attributes as they help in fighting against cancer, inflammation, cardiovascular, gastro-intestinal disorders and epidermis problems. Thus, sea buckthorn could be utilized as a promising source of functional and therapeutic food for the consumption of human. Although, a future scope is really required for exploring the mechanism of sea buckthorn nutritional and phytochemical constituents along with the incorporation of cost-effective food products.

References

- Bal LM, Meda V, Naik SN, Satya S. Sea buckthorn berries: A potential source of valuable nutrients for nutraceuticals and cosmoceuticals. *Food Research International*. 2011;44(7):1718-1727.
- Bittová M, Krejzová E, Roblová V, Kubáň P, Kubáň V. Monitoring of HPLC profiles of selected polyphenolic compounds in sea buckthorn (*Hippophaë rhamnoides* L.) plant parts during annual growth cycle and estimation of their antioxidant potential. *Open Chemistry*. 2014;12(11):1152-1161.
- Chandra S, Zafar R, Dwivedi P, Shinde LP, Prita B. Pharmacological and nutritional importance of sea buckthorn (*Hippophae*). *The Pharma Innovation*. 2018;7(5):258-263.
- Christaki E. *Hippophae rhamnoides* L. (Sea Buckthorn): a potential source of nutraceuticals. *Food Public Health*. 2012;2(3):69-72.
- Crăciun I. Comparative Study of Liposoluble Vitamins and Fatty Acids from Sea Buckthorn Oil, Wheat Germ Oil and Fish Oil. *Acta Universitatis Cibinensis, Series E: Food Technology*. 2018;22(2):85-90.
- Criste A, Urcan AC, Bunea A, Pripon Furtuna FR, Olah NK, Madden RH, Corcionivoschi N. Phytochemical composition and biological activity of berries and leaves from four Romanian sea buckthorn (*Hippophae rhamnoides* L.) varieties. *Molecules*. 2020;25(5):1170.
- Daško L, Drimal J, Klimeková M, Ůrgeová E. Response of organically and conventionally produced potatoes to a controlled attack of a pathogen. *Journal of Food and Nutrition Research*. 2012;51(2):89-95.

- Dulf FV. Fatty acids in berry lipids of six sea buckthorn (*Hippophae rhamnoides* L., subspecies *carpatica*) cultivars grown in Romania. *Chemistry Central Journal*. 2012;6(1):1-12.
- Edraki M, Akbarzadeh A, Hosseinzadeh M, Tanideh N, Salehi A, Koochi-Hosseiniabadi O. Healing effect of sea buckthorn, olive oil, and their mixture on full-thickness burn wounds. *Advances in skin & wound care*. 2014;27(7):317-323.
- Fan XM, Shen XL, Yan LL, Wang XC, Yin JT. The contents of quercetin, kaempferol and isorhamnetin in the fruit of *Hippophae rhamnoides* were determined by HPLC. *Chinese Journal of Hospital Pharmacy*. 2012; 32(17):1343–1346.
- Fatima T, Snyder CL, Schroeder WR, Cram D, Datla R, Wishart D, Weselake RJ, Krishna P. Fatty acid composition of developing sea buckthorn (*Hippophae rhamnoides* L.) berry and the transcriptome of the mature seed. *PloS one*. 2012;7(4):e34099.
- García VL. The omega 7 as a health strategy for the skin and mucous membranes. *EC Nutrition*. 2019; 14:484-489.
- Guo XF, Yang B, Cai W, Li D. Effect of sea buckthorn (*Hippophae rhamnoides* L.) on blood lipid profiles: a systematic review and meta-analysis from 11 independent randomized controlled trials. *Trends in Food Science & Technology*. 2017; 61:1-10.
- Heinaaho M, Julkunen-Tiitto R. Efficient extraction of flavonoids of sea buckthorn berries. *BioChemistry: An Indian Journal*. 2011; 5:83-87.
- Hussain M, Ali S, Awan S, Hussain M, Hussain I. Analysis of minerals and vitamins in sea buckthorn (*Hippophae rhamnoides*) pulp collected from Ghizer and Skardu districts of Gilgit-Baltistan. *International Journal of Bioscience*. 2014; 4:144-152.
- Ivanišová E, Mittuchová K, Mareček J, Frančáková H. Small Berries—Attractive Source of Bioactive Compounds for Consumers. *Agrobiodiversity for Improving Nutrition, Health and Life Quality*. 2017; 1:188-194.
- Koyama T, Taka A, Togashi H. Effects of a herbal medicine, *Hippophae rhamnoides*, on cardiovascular functions and coronary microvessels in the spontaneously hypertensive stroke-prone rat. *Clinical Hemorheology and Microcirculation*. 2009;41(1):17-26.
- Larmo PS, Yang B, Hurme SA, Alin JA, Kallio HP, Salminen EK, Tahvonon RL. Effect of a low dose of sea buckthorn berries on circulating concentrations of cholesterol, triacylglycerols, and flavonols in healthy adults. *European Journal of Nutrition*. 2009;48(5):277-282.
- Larmo PS, Yang B, Hyssälä J, Kallio HP, Erkkola R. Effects of sea buckthorn oil intake on vaginal atrophy in postmenopausal women: a randomized, double-blind, placebo-controlled study. *Maturitas*. 2014;79(3):316-321.
- Leskinen HM, Suomela JP, Yang B, Kallio HP. Regioisomer compositions of vaccenic and oleic acid containing triacylglycerols in sea buckthorn (*Hippophae rhamnoides*) pulp oils: influence of origin and weather conditions. *Journal of Agricultural and Food chemistry*. 2010;58(1):537-645.
- Lu SG, Lu J, X F. Wen. Review on plant resources distribution and nutrition application of sea buckthorn. *Soil and Water Conservation in China SWCC*. 2019; 7:45–49.
- Marsiñach MS, Cuenca AP. The impact of sea buckthorn oil fatty acids on human health. *Lipids in health and disease*. 2019;18(1):1-11.

- Michel T, Destandau E, Le Floch G, Lucchesi ME, Elfakir C. Antimicrobial, antioxidant and phytochemical investigations of sea buckthorn (*Hippophaë rhamnoides* L.) leaf, stem, root and seed. Food Chemistry. 2012;131(3):754-760.
- Nazir F, Bashir M. Chemical and antioxidant properties of Sea buckthorn (*Hippophae rhamnoides*). The Pharma Innovation Journal. 2017;6(12):173-176.
- Nour V, Panaite TD, Corbu AR, Ropota M, Turcu RP. Nutritional and bioactive compounds in dried sea-buckthorn pomace. Erwerbs-Obstbau. 2021;63(1):91-98.
- Padmavathi B, Upreti M, Singh V, Rao AR, Singh RP, Rath PC. Chemoprevention by *Hippophae rhamnoides*: effects on tumorigenesis, phase II and antioxidant enzymes, and IRF-1 transcription factor. Nutrition and Cancer. 2005;51(1):59-67.
- Patel CA, Divakar K, Santani D, Solanki HK, Thakkar JH. Remedial prospective of *Hippophae rhamnoides* Linn. (sea buckthorn). International Scholarly Research Notices.2012.
- Pentelescu C, Pop DM, Colceriu L, Deleanu A. Use of Sea-buckthorn (*Hippophae rhamnoides*) in the Treatment of Acute Gingivitis. Medicine and Pharmacy Reports. 2012;85(2):254-259.
- Pop EA, Diaconeasa ZM, Fetea F, Bunea A, Dulf F, Pintea A, Socaciu C. Carotenoids, tocopherols and antioxidant activity of lipophilic extracts from sea buckthorn berries (*Hippophae rhamnoides*), apricot pulp and apricot kernel (*Prunus armeniaca*). Bulletin UASVM Food Science and Technology. 2015;72(2):14-21.
- Pop RM, Weesepeol Y, Socaciu C, Pintea A, Vincken JP, Gruppen H. Carotenoid composition of berries and leaves from six Romanian sea buckthorn (*Hippophae rhamnoides* L.) varieties. Food Chemistry. 2014; 147:1-9.
- Preedy VR, editor. Handbook of Nutrition, Diet, and the Eye. Academic Press. 2014.
- Rop O, Ercişli S, Mlcek J, Jurikova T, Hoza I. Antioxidant and radical scavenging activities in fruits of 6 sea buckthorn (*Hippophae rhamnoides* L.) cultivars. Turkish Journal of Agriculture and Forestry. 2014;38(2):224-232.
- Ranjith A, Kumar KS, Venugopalan VV, Arumughan C, Sawhney RC, Singh V. Fatty acids, tocopherols, and carotenoids in pulp oil of three sea buckthorn species (*Hippophae rhamnoides*, *H. salicifolia*, and *H. tibetana*) grown in the Indian Himalayas. Journal of the American Oil Chemists Society. 2006;83(4):359-364.
- Rashid A, Raisuddin A, Kumar ND, Abhinav J, Amit T, Gaurav M, Aseem B. Safety and efficacy study of intramuscularly administered sea buckthorn (*Hippophae rhamnoides* L.) oil as depot formulation. International Journal of Drug Development & Research. 2011;3(3):356-365.
- Sayegh M, Miglio C, Ray S. Potential cardiovascular implications of Sea Buckthorn berry consumption in humans. International Journal of Food Sciences and Nutrition. 2014;65(5):521-528.
- Singh IP, Ahmad F, Gore DD, Tikoo K, Bansal A, Jachak SM, Jena G. Therapeutic potential of seabuckthorn: a patent review (2000-2018). Expert Opinion on Therapeutic Patents. 2019;29(9):733-744.
- Stobdan T, Korekar G, B Srivastava R. Nutritional attributes and health application of seabuckthorn (*Hippophae rhamnoides* L.)-A review. Current Nutrition & Food Science. 2013;9(2):151-165.

- Süleyman H, Demirezer LÖ, Büyükokuroglu ME, Akcay MF, Gepdiremen A, Banoglu ZN, Göçer F. Antiulcerogenic effect of *Hippophae rhamnoides* L. *Phytotherapy Research*. 2001;15(7):625-627.
- Suryakumar G, Gupta A. Medicinal and therapeutic potential of Sea buckthorn (*Hippophae rhamnoides* L.). *Journal of Ethnopharmacology*. 2011;138: 268-278.
- Teleszko M, Wojdyło A, Rudzinska M, Oszmianski J, Golis T. Analysis of lipophilic and hydrophilic bioactive compounds content in sea buckthorn (*Hippophae rhamnoides* L.) berries. *Journal of Agricultural and Food Chemistry*. 2015;63(16):4120-4129.
- Tkacz K, Wojdyło A, Turkiewicz IP, Bobak Ł, Nowicka P. Anti-oxidant and anti-enzymatic activities of sea buckthorn (*Hippophaë rhamnoides* L.) fruits modulated by chemical components. *Antioxidants*. 2019;8(12):618.
- Trineeva OV, Safonova EF, Sinkevich AV, Slivkin AI. Assay of amino acids in medicinal plants by TLC (using stinging nettle leaves and common sea buckthorn fruits as examples). *Pharmaceutical Chemistry Journal*. 2015;49(5):323-328.
- Vaitkeviciene N, Danilcenko H, Mazeika R, Jariene E, Vaitkeviciene N, Kulaitiene J, Hallmann E, Blinstrubien A. Comparison of mineral and fatty acid composition of wild and cultivated sea buckthorn berries from Lithuania. *Journal of Elementology*. 2019;24(3):1101–1113.
- Wang K, Xu Z, Liao X. Bioactive compounds, health benefits and functional food products of sea buckthorn: A review. *Critical Reviews in Food Science and Nutrition*. 2021:1-22.
- Wani TA, Wani SM, Ahmad M, Ahmad M, Gani A, Masoodi FA. Bioactive profile, health benefits and safety evaluation of sea buckthorn (*Hippophae rhamnoides* L.): A review. *Cogent Food & Agriculture*. 2016;2(1):1128519.
- Wen P, Zhao P, Qin G, Tang S, Li B, Zhang J, Peng L. Genotoxicity and teratogenicity of seabuckthorn (*Hippophae rhamnoides* L.) berry oil. *Drug and Chemical Toxicology*. 2020;43(4):391-377.
- Wu ZJ, Ruan CJ, Li H, Yang ML, Xie MM, Xiao TT. Compositions of Soluble Sugars and Organic Acids in Berries of 12 Sea-buckthorn Cultivars. *Journal of Northwest Forestry University*. 2016; 4:106-112.
- Xu YJ, Kaur M, Dhillon RS, Tappia PS, Dhalla NS. Health benefits of sea buckthorn for the prevention of cardiovascular diseases. *Journal of functional foods*. 2011;3(1):2-12
- Yasukawa K, Kitanaka S, Kawata K, Goto K. Anti-tumor promoters phenolics and triterpenoid from *Hippophae rhamnoides*. *Fitoterapia*. 2009;80(3):164-167.
- Zakynthinos G, Varzakas T. *Hippophae rhamnoides*: safety and nutrition. *Current Research in Nutrition and Food Science Journal*. 2015;3(2):89-97.
- Zhao P, Wang S, Liang C, Wang Y, Wen P, Wang F, Qin G. Acute and subchronic toxicity studies of seabuckthorn (*Hippophae rhamnoides* L.) oil in rodents. *Regulatory Toxicology and Pharmacology*. 2017; 91:50-57.
- Zielińska A, Nowak I. Abundance of active ingredients in sea-buckthorn oil. *Lipids in Health and Disease*. 2017;16(1):1-11.
- Suryasa, W., Sudipa, I. N., Puspani, I. A. M., & Netra, I. (2019). Towards a Change of Emotion in Translation of Kṛṣṇa Text. *Journal of Advanced Research in Dynamical and Control Systems*, 11(2), 1221-1231.

- Suwija, N., Suarta, M., Suparsa, N., Alit Geria, A.A.G., Suryasa, W. (2019). Balinese speech system towards speaker social behavior. *Humanities & Social Sciences Reviews*, 7(5), 32-40. <https://doi.org/10.18510/hssr.2019.754>
- Gandamay, I. B. M., Antari, N. W. S., & Strisanti, I. A. S. (2022). The level of community compliance in implementing health protocols to prevent the spread of COVID-19. *International Journal of Health & Medical Sciences*, 5(2), 177-182. <https://doi.org/10.21744/ijhms.v5n2.1897>