

SOIL FERTILITY STATUS AND SOYBEAN [*Glycine max* (L) Merr] PERFORMANCE FOLLOWING INTRODUCTION OF INDIGENOUS MYCORRHIZA COMBINED WITH VARIOUS NUTRIENT SOURCES INTO SANDY SOIL

Wahyu Astiko ¹⁾, Ika Rocdjatun Sastrahidayat ²⁾, Syamsuddin Djauhari ²⁾ and Anton Muhibuddin²⁾

¹⁾ Faculty of Agriculture Mataram University

Jl. Majapahit No. 62 Mataram 83124, West Nusa Tenggara

²⁾ Faculty of Agriculture Brawijaya University Jl. Veteran Malang 65145 East Java Indonesia

^{*)}Correspondence author Phone:+62-8123788910 E-mail: astiko_mataram@yahoo.co.id

Received: January 2, 2013/ Accepted: May 29, 2013

ABSTRACT

The experiment tested four treatments, namely inoculation of soil with mycorrhiza, inoculation of soil with mycorrhiza and cattle manure, inoculation of soil with mycorrhiza and rock phosphate, and inoculation of soil with mycorrhiza and inorganic fertilizers. Soil without any inoculation was provided as control. The treatments were arranged in a Completely Randomized Design with four replications. Soil fertility status was based on the concentrations of N, P, K, and organic-C as well as soil pH. Plant performance was determined based on its ability to uptake nutrients (N, P, K, and Ca), its growth and yield. Then, the activity of mycorrhiza was measured based on total population and percentage of root infection. Overall, results of the present study showed that, compared with other treatments, inoculation of mycorrhiza into soil and amended with cattle manure significantly performed higher concentrations of N, P, K, and organic-C. This soil condition caused soybean to absorb significantly higher nutrients, grew well with higher yield compared with plant performance in other treatments. Therefore, results of this study implies that the application of mycorrhiza into soil amended with organic matter is promising to sustain soil productivity under soybean cropping system.

Keywords: *Arbuscular Mycorrhiza*, soil fertility, soybean yield, dryland

INTRODUCTION

Shortage in availability of water, nutrients, and organic matter was a core problems in improving plant production in sandy soil of northern Lombok (Suwardji *et al.*, 2007). Sandy soil with low organic matter content has low

capacity in holding water and nutrients to support optimal soybean performance (Suzuki and Noble, 2007; Bastida *et al.*, 2010). This character of sandy soil is considered as a specific problem in managing sandy soil in northern Lombok, especially to grow soybean. To overcome this particular problem, a strategy in managing soil, in long term, to gain improvement and stability of soil organic matters which then lead to the improvement of soil characteristics especially in holding capacity of water, plant performance and nutrients in the state of ready to be used by plants (Astiko *et al.*, 2013). In addition, soil management by improving role of indigenous arbuscular mycorrhiza (AM) in sandy soil is one of best possible alternative solutions to improve plant productivity (Herrera *et al.*, 1993 and Astiko, 2009). This was due to the role of AM in improving soil quality through the improvement of aggregate and colloid of the soil (Ijdo *et al.*, 2011). This role became better in soil with adequate content of organic matter supplying carbón and micro nutrients required by AM to grow (Öpik *et al.*, 2008; Smith and Read, 2008).

Optimizing role and beneficial characteristic of AM through application of biofertilizer to improve plant productivity in dry land is prospective enough (Sastrahidayat *et al.*, 2001 and Astiko *et al.*, 2005). Application of AM by introducing the organisms into soil is expected to be able to improve plant productivity significantly through role of AM in improving plant capacity to absorb N, P, K, Ca and other micro nutrients. Besides, with its external hypha, AM will improve plant resistance on drought, protect plant root from soil-born pathogen infection, stimulate activity of beneficial microorganisms, and improve soil texture and structure (Gianinazzi and Vosátka, 2004; Feldmann *et al.*, 2009; Ijdo *et al.*, 2011).

Accredited SK No.: 81/DIKTI/Kep/2011

<http://dx.doi.org/10.17503/Agrivita-2013-35-2-p127-137>

Results of many previous studies have proven that nutrient absorption, growth and result of plant inoculated by AM were much higher than control (Azcón-Aguilar and Barea, 1997; Nogueira *et al.*, 2007; Fisher and Jayachandran, 2008). Furthermore, it was reported that the use of indigenous AM had advantages as the microorganism was able to establish and develop well in situ environment as well as its better ability to compete with existing soil microbes compared with introduced AM (Turrini *et al.*, 2008).

Incorporation of AM fungus on soybean in sandy soil of dry-land Northern Lombok was expected to have positive impact on soil characteristic improvement, nutrient absorption, and finally plant growth and yield. This hypothesis was constructed based on results of previous research in Vertisol soil (Astiko *et al.*, 2005) proving that inoculation of AM on soybean improved absorption of P and plant yield was higher than that of plant without AM. The improvement of P absorption was due to AM activity in improving nutrient availability and root proliferation (Smith *et al.*, 2010). This increase of plant yield was suggested due to the ability of AM to increase efficiency in water use, nutrient absorption, and to maintain turgor of plant cells. The external hypha of AM fungus were expected to be able to absorb soil pore water when plant roots enable to do so. In addition, wide distribution of external hypha caused the amount of water taken improved on the soil with low water content (Drew *et al.*, 2003; Smith and Read, 2008). Based on the above mentioned, the aim of this study was to assess soil fertility status and soybean performance following introduction of indigenous mycorrhiza combined with various nutrient sources into sandy soil.

MATERIALS AND METHODS

Experiment Preparation and Maintenance

A series of pot experiments using sandy soil taken Northern Lombok was conducted in a glasshouse of Faculty of Agriculture, Mataram University using 10 kg polybag as experimental unit. Taken from upper layer up to 30 cm depth, soil was passed through 2-mm holes sieve, air dried, then filled into the polybags. The experiment was conducted in two cropping cycles, namely; the first cropping cycle in which soybean was grown and fertilized according to treatments, and the

second cropping cycle in which soybean was grown on soil that had been used for the first cropping cycle without fertilization. The study tested four treatments, namely; soil was inoculated with AM (F1), soil was inoculated with AM and amended with cattle manure (F2), soil was inoculated with AM and amended with rock phosphate (F3), and soil was inoculated with AM and amended with inorganic fertilizers (F4). Soil without AM inoculation and amendment was provided as control (F0). The treatments were completely randomized designed with four replications. AM inoculum was mass produced in corn plants grown in pots. Pots were filled with mixture of sandy soil and sterile cattle manure (ratio 1:1). The soil was watered about field capacity and the plants were maintained in glass house for three months. Then the plants were harvested, dried, blended, and passed through 50-mesh sieve. The final form of the proliferation was powder. The AM inoculums was prepared by mixing spores, powder medium, and root residues. The AM inoculum (20 gram per pot) then was inoculated at sowing time by layering the inoculum at 10 cm depth. Soybean seeds of Kaba var. were sown (2 seeds per pot) at 3 cm depth and 14 days after sowing (das), only one seedling per pot was left to grow further. Nutrient sources applied according to treatments were cattle manure, rock phosphate, Urea, and SP36. Rock phosphate and cattle manure were applied at rate of 1.2 and 2.0 g per plant respectively, and inorganic fertilizers of Urea and SP-36 were applied at rate of 0.1 g and 0.2 g per plant, consecutively. The nutrient sources were buried 5 cm around the plant at depth of 7 cm. Weeding was done manually while watering plants in the afternoon. Plant protection was done by applying organic pesticide 0.5% Azadirachtin (OrgaNeem™) every three days. The plants were harvested 100 das.

Parameters Observation

Parameters dealing with soil fertility status (N, P, K, organic-C and soil pH) were measured before sowing and at 60 and 100 das. The agronomic parameters such as: top and roots dry weight biomass (60 das and 100 das), nutrients uptake (N, P, K, and Ca) at 60 das and component yield of soybean (cops, grain and 100 grain weight). The dry weight of the agronomic parameters were measured after being oven dried at 60 °C for 48 hours. Parameters related to AM activities including fungi population, roots

percentage infections at 60 das were also measured. Plant analysis for N was determined using Kjeldhal method, P using spectrophotometer, C-organic with colorimetric method according to Walkley and Black, K and Ca was recorded using Automatic Absorbance Spectrophotometer (AAS).

Soil, Plant Analysis and Mycorrhiza Observation

Analyses for N, P, and organic-C were done by using Kjeldhal method, spectrophotometer, and colorimetric method according to Walkley and Black, respectively. K and Ca were analyzed by using AAS. Mycorrhiza population was observed using wet sieving technique according to Brundrett *et al.* (1996). The supernatant caught at 38 μ -sieve was added with 60% of sucrose solution and subsequently centrifuged at 3000 rpm for 10 minutes (Daniel and Skipper, 1982). The harvested spores were stored on the Whatman paper with permanent ink marked of 0.5 x 0.5 cm. Counting of mycorrhiza population was done using stereo microscope (extended 40 x). Calculation of roots percentage infections was conducted using modification of clearing and staining method (Kormanik and McGraw, 1982), counted with the *Gridline Intersect* technique (Giovenetti and Mosse, 1980) under stereo-microscope observation.

Statistical analysis

To know if there is any difference among treatments, a mathematic model below was applied:

$$X_{ijk} = \mu + \rho_i + \beta_j + \varepsilon_{ij}$$

Remarks:

- μ = general average
- ρ_i = effect of replication-i
- β_j = effect of treatment-j
- ε_{ij} = error

The model was applied when $X_{ijk} = \mu + \rho_i + \beta_j$ linear and additive to $\sum \rho_i = \sum \beta_j = 0$, and ε_{ijk} was free and normally distributed with average and variance = $(0, \sigma^2)$. Data were analysed by analysis of variance (ANOVA-MStat) and the effect of treatments was determined. When the variance ratio (F) was significant, means for each treatment were separated using a least significant difference test at 5% level.

RESULTS AND DISCUSSION

Soil Chemical Properties

In general, compared with control, all treatments significantly increased soil fertility status as indicated by concentration of total N, available P and K, and organic-C both at 60 das and 100 das (Table 1). The highest increases of soil fertility status were observed in sandy soil inoculated with arbuscular mycorrhiza (AM) and amended with cattle manure (CM). This indicated that AM performed well when combined with CM compared with other combinations. Compared with control, inoculation of AM amended with CM increased N, P, K, and org-C as much as 39%, 105%, 27%, and 85%, respectively on 60 das and 12%, 60%, 10%, and 11%, consecutively on 100 das at the first cropping cycle. Interestingly, these effects were also observed at the second cropping cycle as the total N, P, K, and org-C increased as much as 43%, 120%, 37%, and 36%, respectively on 60 das, and 44%, 53%, 14%, and 36%, respectively on 100 das. Data presented in Table 1 also indicated that at the second cropping cycle soybean performance on treatment of AM plus CM could improve soil fertility status as shown by nutrient concentration of P and N as much as 4 and 48% on 60 das, respectively, and on 100 das the treatment increased N and Org-C as much as 12 and 20%, consecutively.

Table 1. Soil fertility status (N, P, K, organic-C and soil pH) of sandy soil with various treatments after harvesting

Treatments	N (g kg ⁻¹)		P (mg kg ⁻¹)		K (cmol kg ⁻¹)		org-C (g kg ⁻¹)		pH	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
60 DAS										
F ₀	1.15 ^a	0.87 ^a	16.97 ^a	16.54 ^a	0.69 ^a	0.51 ^a	12.1 ^a	24.5 ^a	6.25 ^a	6.13 ^a
F ₁	1.34 ^b	1.15 ^b	23.60 ^b	23.40 ^b	0.75 ^b	0.58 ^b	17.9 ^b	26.5 ^b	6.01 ^b	6.24 ^b
F ₂	1.60 ^c	1.25 ^b	34.83 ^c	36.54 ^c	0.88 ^c	0.70 ^c	22.5 ^c	33.5 ^c	5.95 ^c	6.32 ^c
F ₃	1.44 ^d	1.10 ^b	26.34 ^d	27.01 ^d	0.83 ^d	0.63 ^d	21.0 ^d	29.4 ^d	6.72 ^d	6.28 ^d
F ₄	1.40 ^e	0.93 ^b	20.59 ^d	30.97 ^e	0.78 ^c	0.62 ^{bd}	19.1 ^e	26.3 ^b	6.08 ^e	6.26 ^{bd}
100 DAS										
F ₀	1.31 ^a	1.14 ^a	17.62 ^a	17.33 ^a	0.75 ^a	0.63 ^a	23.8 ^a	23.5 ^a	6.18 ^a	6.54 ^a
F ₁	1.44 ^b	1.36 ^b	20.86 ^b	22.68 ^{bc}	0.77 ^b	0.65 ^b	24.5 ^b	25.3 ^{ab}	6.21 ^b	6.62 ^{ab}
F ₂	1.47 ^c	1.65 ^c	28.25 ^c	26.57 ^d	0.83 ^c	0.72 ^c	26.6 ^c	32.1 ^c	6.24 ^c	6.54 ^{ab}
F ₃	1.38 ^d	1.41 ^b	23.32 ^d	20.62 ^c	0.81 ^d	0.67 ^b	23.9 ^d	28.0 ^d	6.62 ^d	6.75 ^a
F ₄	1.33 ^a	1.43 ^b	24.38 ^e	24.57 ^{bd}	0.82 ^e	0.70 ^c	25.7 ^e	27.5 ^d	6.23 ^c	6.44 ^b
Before exp ¹⁾	1.10	-	13.82	-	0.57	-	12.1	-	6.25	-

Remarks: Means followed by the same letters within the same column are not significantly different ($p=0.05$); 1st and 2nd (first cycles and second cycles); F₀= Control, F₁= AM inoculation, F₂= AM inoculation plus CM, F₃= AM inoculation plus RP, and F₄= AM inoculation plus CF; ¹⁾ pre-treatment data

Results of this study were in accordance with those of done by Jeffries *et al.* (2003) and Gianinazzi *et al.* (2010) reporting that AM inoculation with organic matter amendment can recover soil fertility status on an ecosystem. Furthermore, Douds *et al.* (2006) reported that introduction AM inoculation combined with various nutrient sources into sandy soil could increase soil nutrient content. The same result was also reported by Astiko *et al.* (2013) evaluating contribution of indigenous AM combined with cattle manure to increase corn performance in sandy soil of northern Lombok. Combination of AM and cattle manure resulted in significant improvement on soil fertility status especially N, P, K, and organic-C. This increase was suggested due to activity of enzyme present in external hypha of AM in the rhizosphere able to catalyze and hydrolyze unavailable nutrient complex into available nutrients (Widiastuti *et al.*, 2003).

Previous results reported by Warnock *et al.*, (2007) showed that enrichment of AM could be escalated by the addition of organic matter and the combination has a positive effect on improving soil fertility status and therefore it is beneficial for soybean performance. This signifies that better synergy of the inoculated AM-treatment in combination with CM, as previously confirmed by Kato and Miura (2008). Higher soil extractable-P

found in this study was not only due to indirect contribution of CM in improving soil fertility status but also related to a positive contribution of mycorrhiza in producing phosphatase enzyme, for mineralization of organic-P in soil (Crowley and Rengel, 2000; Joner and Johansen, 2000) and resulted in enhancing insoluble-P in soils (Orcut and Nilsen, 2000).

Nutrient Uptake, Growth and Yield

Nutrient absorption by soybean is presented in Table 2. In general, treatments such as AM inoculation alone or combined with other sources of nutrients significantly increased nutrient uptake by soybean, compared with control. The highest increases were observed at treatment AM inoculation amended with CM. At this treatment, absorption of N, P, K, and Ca increased as much as 214%, 185%, 342%, and 233%, respectively on the first cropping cycle, and 191%, 308%, 224%, and 413%, consecutively on the second cropping series, compared with control. Treatment of AM inoculation followed by CM amendment also caused high absorption of P and Ca as much as 29% and 126%, respectively on the second cropping cycle. This indicated an increase in soybean performance when grown in sandy soil amended with various nutrient sources.

Table 2. Nutrient uptake (N,P, K and Ca) by soybean grown in sandy soil with various treatments.

Treatments	Nutrient uptake (mg.plant ⁻¹)							
	First cycle				Second cycle			
	N	P	K	Ca	N	P	K	Ca
60 DAS								
F ₀	221.59 ^a	22.43 ^a	121.53 ^a	25,60 ^a	224.52 ^a	20.37 ^a	79.21 ^a	37.66 ^a
F ₁	302.83 ^b	29.03 ^b	210.13 ^b	34,16 ^b	301.13 ^b	34.46 ^b	121.63 ^{ab}	59.22 ^b
F ₂	697.16 ^c	64.06 ^c	537.33 ^c	85,30 ^c	653.42 ^c	83.22 ^c	257.43 ^c	93.30 ^c
F ₃	412.73 ^d	39.55 ^d	282.73 ^d	48,20 ^d	429.60 ^d	49.33 ^d	185.20 ^d	12.96 ^d
F ₄	344.10 ^e	34.46 ^e	232.46 ^e	39,96 ^d	311.60 ^b	38.14 ^b	159.44 ^{ad}	96.93 ^d

Remarks: Means followed by the same letters within the same column are not significantly different ($p=0.05$); F₀= Control, F₁= AM inoculation, F₂= AM inoculation plus CM, F₃= AM inoculation plus RP, and F₄= AM inoculation plus CF

The increase of P absorption could possibly cause such a new nutrient balance in plant that induced absorption of other nutrients such as N, K, and Ca. Sufficient availability of K created a condition in which the use of water was efficient as cell turgor was maintained. This condition leads to active metabolism process, K accumulation on the tips, buds, and roots, the accumulation that lead to formation of cortex tissue and cell elongation leading to improvement of soybean performance (Schweiger *et al.*, 2007 and Smith *et al.*, 2010). This result was in accordance with that of Kaschuk *et al.* (2010) stating that the increase of nutrient absorption by host plant lead to plant vitality to supply Carbon to rhizosphere to form AM external hypha. Mathur and Vyas (2000) stated that AM inoculation was also resulted in the increase of accumulation of amino acids, protein, chlorophyll, and sugar contents compared with non-AM plants. N status of shoot tip of plant with mycorrhiza at extreme condition was higher than that of plant without mycorrhiza (Subramanian and Charest, 1999). The same trends were also recorded for N, P, K, and Ca (Liu *et al.*, 2000).

The inoculation of AM followed by CM also increased soybean performance, as observed on plant dry biomass measured by root and shoot dry weights on 60 and 100 das (Table 3). Dry weight of roots and shoots on 60 das increased as much as 164% and 136%, respectively; while on 100 das the increases were as much as 150% and 178%, respectively on the first cropping cycle. On the second cycle, on 60 das the increases were 337% and 718%, while on 100 das the increases were as much as 390% and 1102%, consecutively.

The improvement of biomass was probably due to AM role to influence soil fertility status, especially P. Sufficient availability of P can indirectly induce absorption of other nutrients leading to better plant growth (Carrenho *et al.*, 2001). This positive effect was suggested due to suitability of AM type, plant, and soluble P (Nikolaou *et al.*, 2002; Bhadalung *et al.*, 2005).

This condition was due to good association between plant and AM to perform maximal activity. While AM received carbon from the plant, and the latter got P from the first. It was reported that each particular combination of AM and plant showed such a specific carbon translocation pattern that influenced production of plant biomass (Smith and Read, 2008; Smith *et al.*, 2009). Inefficient symbiosis in use of carbon could decrease plant biomass. AM colonization could result in positive, neutral, or negative impact depending on AM types, plants, and growth environment (Johnson *et al.*, 1997; Hoeksema *et al.*, 2010).

Negative impact of mycorrhiza colonization on initial growth of plant was reported previously (Bethlenfalvay *et al.*, 1982; Koide 1985; Johnson *et al.*, 1997). Such sort of impact may be due to various factors, such as; high availability of P in soil (Mosse *et al.*, 1973), competition for carbon between plant and AM in low light intensity condition (Buwalda and Goh 1982), and difference in biomass allocation pattern between plant with and without mycorrhiza (Smith and Read, 2008).

Table 3. Root and shoot dry weight of soybean grown in sandy soil with various treatments at 60 and 100 das

Treatments	Biomass dry weight (g.plant ⁻¹)			
	First cycle		Second cycle	
	Roots	Shoots	Roots	Shoots
60 DAS				
F ₀	1.88 ^a	11.11 ^a	1.74 ^a	8.50 ^a
F ₁	2.66 ^b	14.51 ^b	2.27 ^b	10.48 ^b
F ₂	4.97 ^c	26.28 ^c	4.38 ^c	18.67 ^c
F ₃	3.62 ^d	20.39 ^d	3.75 ^d	13.51 ^d
F ₄	3.34 ^d	17.10 ^e	2.51 ^b	11.41 ^e
100 DAS				
F ₀	3.41 ^a	22.43 ^a	2.83 ^a	19.94 ^a
F ₁	4.45 ^b	39.63 ^b	2.54 ^a	19.35 ^a
F ₂	8.53 ^c	62.45 ^c	6.27 ^b	32.92 ^b
F ₃	5.22 ^d	44.50 ^d	3.41 ^a	26.40 ^c
F ₄	4.48 ^b	37.85 ^b	2.36 ^a	19.91 ^c

Remarks: Means followed by the same letters within the same column are not significantly different ($p=0.05$); F₀= Control, F₁= AM inoculation, F₂= AM inoculation plus CM, F₃= AM inoculation plus RP, and F₄= AM inoculation plus CF

The use of low solubility of phosphate, like phosphate stone, although with high dosage, in fact, was still effective to support AM and to increase plant performance (Nikolaou *et al.*, 2002). In addition, quite high dependency of plant on AM indicated that in early stage of its growth, plant needs to associate with AM. Table 3 also showed significant effect of AM inoculation combined with CM as indicated by the increase plant performance as observed on root and shoot dry weight. The similar results were also reported previously (Rochdjatun *et al.*, 2011; Astiko *et al.*, 2012).

Inoculation of AM with CM amendment increased plant performance as observed on cobs weight, weight of grain, and weight of 100 grains as much as 180%, 163%, and 139% respectively on the first cropping cycle and as much as 139%, 330%, and 23%, consecutively on the second cycle (Figure 1). These results were in accordance with results of previous studies indicating that fertilizer package with AM inoculation and amended with CM increased soil fertility status, plant performance and yield (Astiko, 2009; Viti *et al.*, 2010). Similar results were also reported on dry paddy (Kabirun, 2002).

The increases of soil fertility status, plant performance and yield on the treatment of AM inoculation amended with CM were caused by the increase of AM activity in absorption of nutrient and water through its external hypha (EH). This was possibly due to EH can reach depletion zone that cannot be reached by plant roots (Zhu *et al.*, 2001). The diameter of EH which is much smaller than that of roots makes the EH possibly to penetrate soil micro pores to take nutrient and water that cannot be reached by roots (Drew *et al.*, 2003).

This ability causes plants with mycorrhiza be able to absorb nutrient, growth and perform better and resistance to drought stress (Smith and Read, 2008). In addition, AM is able to dilute phosphate tied in soil and fertilizer, to improve absorption of N, P, and K, to improve plant tolerance to drought, to improve plant ability to produce growth regulator, to stimulate activity of beneficial microbes, to improve soil structure and aggregation, and to enhance mineral cycling (Cruz, 1990).

Decomposition and mineralization of organic matter were better with the presence of AM which positively affect physic, chemistry, and biology factors of the soil which in turn play key role in improving plant yield (Smith and Read, 2008). All above facts indicated such a suitable functional among AM, host plants, and environment that are able to increase nutrient absorption, plant growth and yield as also reported earlier (Burleigh *et al.*, 2002).

Mycorrhiza Activity

The inoculation of AM followed by CM amendment could increase AM activity as shown by numbers of spores per 100 g of soil and percentage of root infection (Table 4). The number of spores and percentage of infection on the first cropping cycle 60 das on sandy soil inoculated by AM and amended with CM increased as much as 179% and 266%, respectively, while on the second cycle the increases were 24% and 160%, respectively compared with control and were significantly different from other treatments.




Figure 1. Yields of soybean (cobs, grain and 100 grain dry weight) grown in sandy soil with various treatments. A. the first cropping cycle, and B. the second cropping cycle. Bars with the same letters at the same category are not significantly different ($p=0.05$).

Table 4 showed that the number of spores and root infection were significantly high in all treatments when compared with control. Compared with before sowing, the number of spores and root infection 60 das increased significantly. This increases indicated that isolate M_{AA1} were able to compete with indigenous AM present in the rhizosphere of sandy

soil Northern Lombok, particularly in colony forming inside roots. The other point that can be taken from these facts was that the isolate M_{AA1} used was able to produce abundance propagules in the form of spores and such colonized roots that they were able to live on dynamic and competitive habitat (Barrios, 2007; Doud and Johnson, 2007; Gianinazzi *et al.*, 2010).

Result of a research on the role of indigenous mycorrhiza combined with cattle manure in improving yield of maize (*Zea mays* L.) on sandy loam of Northern Lombok showed that inoculation of AM combined with cattle manure resulted in higher number of spores and infected roots both in the first and the second growth season (Astiko *et al.*, 2013). Similar results were also shown by Prasetya and Anderson (2011) on the assessment of the effect of long term tillage on the arbuscular mycorrhiza colonization of vegetable crops grown in andisols. This fact also indicated that isolate M_{AA1} showed high effectiveness. although they were inoculated on unsterile soil. Similar research on soybean inoculated by AM and the application of organic leaf fertilizer "greenstant" also showed similar results (Wangiyana *et al.*, 2007).

Table 4. Biological activity of mycorrhiza (number of spores and percentage of infections) in sandy soil with various treatments.

Treatment s	Spores 100 g soil ⁻¹ and root infection			
	First cycle		Second cycle	
	Spores	Infection	Spores	Infection
60 DAS				
F ₀	1.060 ^a	21 ^a	3.122 ^a	25 ^a
F ₁	2.159 ^b	41 ^b	3.533 ^b	34 ^b
F ₂	2.960 ^c	77 ^c	3.878 ^c	65 ^c
F ₃	2.343 ^d	54 ^d	3.781 ^d	51 ^d
F ₄	2.215 ^b	46 ^e	3.693 ^e	41 ^e
Before exp ¹⁾	371	-	-	-

Remarks: Means followed by the same letters within the same column are not significantly different ($p=0.05$); F₀= Control. F₁= AM inoculation. F₂= AM inoculation plus CM. F₃= AM inoculation plus RP. and F₄= AM inoculation plus CF; ¹⁾ pre-treatment data

CONCLUSION

Inoculation of AM amended with cattle manure improved sandy soil fertility as shown by the increasing soil fertility status as observed on concentrations of N. P. K. and organic-C.

Soybean performance responded positively to the application of AM followed by cattle manure as indicated by plant improvement in nutrient absorption. plant growth and yield. The addition of cattle manure stimulated activity of AM leading to improvement of soil fertility and plant performance.

ACKNOWLEDGEMENTS

This manuscript is a part of the first authors dissertation for the Doctoral Program funded by Post Graduate Scholarships at Agricultural Faculty University of Brawijaya Malang. Thank to the institution of Directorate General of Higher Education (DGHE) as the funders of Ph.D research for 2012 Grant Program by DIPA Number: 023.04.2.415278/2013. December 5. 2012. Letter of Agreement. Impelemtation Research Number: 295.G/SPP-APDD/H18.12/PL/ 2013. May 2. 2013.

REFERENCES

- Astiko, W. , P.W. Wangiyana, I.R. Sastrahidayat and R.S. Tejowulan. 2005. The compatibility of some formulation of VAM fungus P uptake and crop yield grown in various marginal soil in the tropics. ICOM. Rec. 749: 791 (Abstr.)
- Astiko, W. 2009. Fertilizer application package effect on growth and yield of maize on dry land. (in Indonesia). Research Center Mataram University (eds.). Proceeding of the 42th National Seminar Dies Natalis Agriculture Faculty of Mataram University, Mataram. pp 123.
- Astiko, W., I.R. Sastrahidayat, S. Djauhari and A. Muhibuddin. 2012. The role of organic based mycorrhiza to improving soybean yield in the tropical semiarid of Northern Lombok. (in Indonesia) J. Buana Sains. 12(1): 15-20
- Astiko, W., I.R. Sastrahidayat, S. Djauhari and A. Muhibuddin. 2013. The role of indigenous mycorrhiza in combination with cattle manure in improving maize yield (*Zea mays* L.) on sandy loam of Northern Lombok. Eastern of Indonesia. J. of Tropical Soils 18(1): 53-58
- Azcón-Aguilar, C. and J.M. Barea. 1997. Applying mycorrhiza biotechnology to horticulture: significance and potentials. Scientia Horticulturae 68:1-24
- Barrios, E. 2007. Soil biota. ecosystem services and land productivity. Ecol. Econ. 64:269-285
- Bastida, F. ,T. Hernández and C. Garcia. 2010. Soil degradation and rehabilitation: microorganisms and functionality. *in*: Insan H.. I. Franke-Whittle. M. Goberna (editor). Microbes at Work – From Wastes to

- Resources Heidelberg: Springer Verlag. pp. 253-270
- Bethlenfalvy, G., M. Brown and R. Pacovsky. 1982. Parasitic and mutualistic associations between a mycorrhizal fungus and soybean: development of the host plant. *Phytopathol.* 72: 889–893
- Bhadalung, N.N., A. Suwanarit, B. Dell, O. Nopamornbodi, A. Thamchaipenet, and J. Rungchuang. 2005. Effects of long-term NP-fertilization on abundance and diversity of arbuscular mycorrhizal fungi under a maize cropping system. *Plant Soil.* 270: 371–382
- Brundrett, M., N. Bougher, B. Dell, T. Grove and N. Malajczuk. 1996. Working with Mycorrhizas in Forestry and Agriculture. *Aciar Monograph* 32.
- Buwalda, J. and K. Goh. 1982. Host-fungus competition for carbon as a cause of growth depression in vesicular-arbuscular mycorrhizal ryegrass. *Soil Biol Biochem.* 14: 103–106
- Burleigh, S.H., T.R. Cavagnaro and I. Jakobsen. 2002. Functional diversity of arbuscular mycorrhizas extends to the expression of plant genes involved in P nutrition. *J. Exp. Bot.* 53:1593–1601
- Carrenho, R., E.S. Silva, S.F.B. Trufem and V.L.R. Bononi. 2001. Successive cultivation of maize and agricultural practices on root colonization. number of spores and species of AM fungi. *Braz. J. Microbiol.* 32:262–270
- Cruz, R.E. de La. 1990. Final report of the consultant on mycorrhiza program development in The ICU Biotechnology Centre. Bogor Agricultural University . p. 11-30
- Crowley, D.E. and Z. Rengel. 2000. Biology and chemistry of nutrient availability in the rhizosphere. In Z Rengel (eds.) *Mineral nutrition of crops. Fundamental Mechanisms and implications.* The Haworth Press. Inc. NY
- Daniels, B.A. and H.D. Skipper. 1982. Methods for recovery and quantitative estimation of propagules from soil. In N.C. Scenck (Eds.). *Methods and principle of mycorrhiza research.* APS. St. Paul MN. p. 29-36
- Douds, D.D. Jr., G. Nagahashi, P.E. Pfeffer, C. Reider and W.M. Kayser. 2006. On-farm production of AM fungus inoculum in mixtures of compost and vermiculite. *Bioresour Technol.* 97: 809–818
- Douds, D.D. Jr. and N.C. Johnson. 2007. Contributions of arbuscular mycorrhizas to soil biological fertility. Di dalam: Abbott LK. Murphy DV. (editor). *Soil Biological Fertility - A Key to Sustainable Land Use in Agriculture.* New York: Springer Science+ Business Media. p. 129-162.
- Drew, E.A., R.S. Murray and S.E. Smith. 2003. Beyond the rhizosphere: growth and function of arbuscular mycorrhizal external hyphae in sands of varying pore size. *Plant Cell Environ.* 251: 105-114
- Feldmann, F., I. Hutter and C. Schneider. 2009. Best production practice of arbuscular mycorrhizal inoculum. *Soil Biol.* 18: 319-335
- Fisher, J.K. and K. Jayachandran. 2008. Arbuscular mycorrhizal fungi promote growth and phosphorus uptake in *zamia*. a native florida cycad. *Biological Science.* 71(3): 265-272
- Giovannetti, M. and B. Mosse. 1980. An evaluation of techniques to measure vesicular-arbuscular mycorrhiza infection in roots. *New Phytol.* 84: 489-500.
- Gianinazzi, S. and M. Vosátka. 2004. Inoculum of arbuscular mycorrhizal fungi for production systems: science meets business. *Can. J. Bot.* 82: 1264-1271
- Gianinazzi, S., A. Golotte, M.N. Binet, D. van Tuinen, D. Redecker and D. Wipf. 2010. Agroecology: the key role of arbuscular mycorrhizas in ecosystem services. *Mycorrhiza.* 20: 519–530
- Herrera, M.A., C.P. Salamanca and J.M. Barea. 1993. Inoculation of woody legumes with selected arbuscular mycorrhiza fungi and rhizobia to recover desertified Mediterranean ecosystems. *Appl. Environm. Microbiol.* 59: 129-133
- Hoeksema, J.D., V.B Chaudhary, C.A. Gehring, N.C Johnson, J. Karst, R.T Koide. A. Pringle, C.J.D. Zabinski, Bever. J.C. Moore, G.W.T. Wilson, J.N. Klironomos and J. Umbanhowar. 2010. A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecol. Lett.* 13: 394-407
- Ijdo, M., S. Crannenbrouck and S. Declerck. 2011. Methods for large-scale production of AM fungi: past. present. and future. *Mycorrhiza.* 21: 1-16

- Jeffries, P., S. Gianinazzi, S. Perotto, K. Turnau and J.M. Barea. 2003. The contribution of arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil fertility. *Biol. Fertil. Soils* 37: 1-16
- Johnson, N.C., J.H. Graham and F.A. Smith. 1997. Functioning of mycorrhizal associations along the mutualism-parasitism continuum. *New Phytol.* 135: 310-322
- Joner, R.E. and A. Johansen. 2000. Phosphatase activity of external hyphae of two arbuscular mycorrhizal fungi. *Mycol. Res.* 104: 12-26
- Kabirun, S. 2002. Effect of "gogo" rice to arbuscular mycorrhizal and phosphate fertilizers in a entisol soil. *Journal of Soils Science and Environmental.* 3(2): 49-56
- Kato, K. and N. Miura. 2008. Effect of matured compost as a bulking and inoculating agent on the microbial community and maturity of cattle manure compost. *Bioresource Technol.* 99: 3372-3380
- Kaschuk, G., P.A. Leffelaar, K.E. Giller, O. Alberton, M. Hungria and T.W. Kuyper. 2010. Responses of legumes to rhizobia and arbuscular mycorrhizal fungi: A meta-analysis of potential photosynthate limitation of symbioses. *Soil Biol. Biochem.* 42: 125-127
- Kormanik, P.P. and A.C. McGraw. 1982. Quantification of vesicular-arbuscular mycorrhiza in plant roots. In N.C. Scenk (Eds). *Methods and principles of mycorrhizal research.* The American Phytopathological Society. St. Paul. Minnesota. pp. 244
- Koide, R. 1985. The nature of growth depressions in sunflower caused by vesicular-arbuscular mycorrhizal infection. *New Phytol.* 99: 449-462
- Liu, A.C. Hamel, R.I. Hamilton, and D.L. Smith. 2000. Mycorrhizae formation and nutrient uptake of new com (*Zea mays* L.) hybrids with extreme canopy and leaf architecture as influenced by soil N and P levels. *Plant and Soil* 22: 157-166
- Mathur, N. and A. Vyas. 2000. Influence of arbuscular mycorrhizae on biomass production, nutrient uptake and physiological changes in *Ziziphus mauritiana* Lam. Under water stress. *Journal of Arid Environments* 45: 191-195
- Mosse, B. 1973. Plant growth responses to VAM IV. *In* Soil given additional phosphate. *New Phytol.* (72): 127-136
- Nikolaou, N., N. Karagiannidis, S. Koundouras, and I. Fysarakis. 2002. Effects of different P sources in soil on increasing growth and mineral uptake of mycorrhizal *Vitis vinifera* L. (cv Victoria) vines. *J. Int. Sci. Vigne Vin* 36:195-204
- Nogueira, M.A., U. Nehls, R. Hampp, K. Poralla and E.J.B.N. Cardoso. 2007. Mycorrhiza and soil bacteria influence extractable iron and manganese in soil and uptake by soybean. *Plant Soil.* (298): 273-284
- Orcutt, D.M. and E.T. Nilsen. 2000. *The physiology of plants under stress: Soil and biotic factors.* New York. John Wiley and Sons. Inc.
- Öpik, M., Ü. Saks, J. Kennedy and T. Daniell. 2008. Global diversity patterns of arbuscular mycorrhizal fungi—community composition and links with functionality. *In* Varma A. (editor). *Mycorrhiza State of the Art. Genetics and Molecular Biology. Eco-Function. Biotechnology. Eco-Physiology. Structure and Systematics.* Third Edition. Berlin: Springer-Verlag. p. 89-111
- Prasetya, B. and C. Anderson. 2011. Assessment of the effect of long term tillage on the arbuscular mycorrhiza colonization of vegetable crops grown in andisols. *J. Agrivita* 33(1): 85-92
- Rochdjatun, I., S. Djauhari, M. Saleh and A. Muhibuddin. 2011. Control damping off disease caused by *Sclerotium rolfsii* Sacc. using Actinomycetes and VAM fungi on soybean in the dry land based on micro-organism diversity of rhizosphere zone. *J. Agrivita* 33(1): 40-46
- Sastrahidayat, I.R., A.S.M. Subari and M. Bintoro. 2001. Effect of sludge and inoculating arbuscular mycorrhizal fungi on growth and yield of maize. *J. Agrivita* 22(2): 147-155
- Schweiger, P.F., A.D. Robson, N.J. Barrow and L.K. Abbott. 2007. Arbuscular mycorrhizal fungi from three genera induce two-phase plant growth response on a high P-fixing soil. *Plant Soil.* 292: 181-192
- Smith, S.E. and D.J. Read. 2008. *Mycorrhizal symbiosis.* 3rd edn. Elsevier and Academic. New York. London. Burlington. San Diego

- Smith, S.E., H.Y. Li and E.J. Grace. 2009. More than a carbon economy: nutrient trade and ecological sustainability in facultative arbuscular mycorrhizal symbiosis. *New Phytol.* doi: 10.1111/j.1469-8137.2004.01039.x
- Smith, S.E., E. Facelli, S. Pope and F.A. Smith. 2010. Plant performance in stressful environments: interpreting new and established knowledge of the roles of arbuscular mycorrhizas. *Plant Soil* 326: 3-20
- Subramanian, K.S. and C. Charest. 1999. Acquisition of N by external hyphae of an arbuscular mycorrhizal fungus and its impact on physiological responses in maize under drought-stressed and well-watered conditions. *Mycorrhiza* 9: 69-75
- Suardji, G. Suardiari and A.Hippi. 2007. The application of sprinkle irrigation to increase of irrigation efficiency at North Lombok. Indonesia. Paper presented at the Indonesian Soil Science Society Congress IX. Gadjah Mada University. Yogyakarta
- Suzuki, S. and A.D. Noble. 2007. Improvement in water-holding capacity and structural stability of a sandy soil in Northeast Thailand. *Arid Land Research and Management*. 21:37-49
- Turrini, A., A. Luciano, B. Stefano and M. Giovannetti. 2008. In situ collection of endangered arbuscular mycorrhizal fungi in a Mediterranean UNESCO Biosphere Reserve. *Biodivers Conserv.* 17: 643-657
- Viti, C., E. Tetti, F. Dacorosi, E. Lista, E. Rea, M. Tullio, E. Sparvoli and L. Giovannetti. 2010. Compost effect on plant growth-promoting rhizobacteria and mycorrhizal fungi population in maize cultivations. *Compost Science and Utilization* 18(4): 273-281
- Wangiyana, W., M. Sitorus and H. Abdurrachman. 2007. Response of soybean to inoculation with arbuscular mycorrhizal fungi and application of the organic foliar fertilizer "greenstant". *Journal of Agroteksos* 17(3): 157-166 (in Indonesia)
- Warnock, D.D., J. Lehmann, T.W. Kuyper and M.C. Rillig. 2007. Mycorrhizal responses to biochar in soil - concepts and mechanisms. *Plant Soil*. 309: 9-20
- Widiastuti, H., N. Sukarno, L.K. Darusman, D.H. Goenadi, S. Smith and E. Guhardja. 2003. Phosphatase activity and organic acid production in rhizosphere and hyphosphere of mycorrhizal oil palm seedling. *J. of Menara Perkebunan*. 71 (2): 70-81 (in Indonesia).
- Zhu, Y.G., T.R. Cavagnaro, S.E. Smith and S. Dickson. 2001. Backseat driving? Accessing phosphate beyond the rhizosphere-depletion zone. *Trends. Plant Sci.* 6:194-195