

How to Cite:

Pannu, S., Thakur, M., Gulati, P., Kumar, M., Dhingra, D., & Pannu, A. (2022). A recent review on developed analytical methods for detection of curcumin. *International Journal of Health Sciences*, 6(7), 173–194. <https://doi.org/10.53730/ijhs.v6n7.10807>

A recent review on developed analytical methods for detection of curcumin

Shivani Pannu

Assistant Professor

School of pharmacy, Desh Bhagat University, Mandi Gobindgarh, Punjab- 147203

Email: shenupannu70@gmail.com

Mohita Thakur

Assistant Professor

School of pharmacy, Desh Bhagat University, Mandi Gobindgarh, Punjab- 147203

Email: mohitathakur365@gmail.com

Dr. Puja Gulati*

Professor

School of pharmacy, Desh Bhagat University, Mandi Gobindgarh, Punjab- 147203

Professor & Principal, School of Pharmacy, Desh Bhagat University, Mandi, Gobindgarh, Fatehgarh Sahib, Punjab

*Corresponding author email: puja_duggal@yahoo.co.in

Mohit Kumar

Ph.D Research Scholar, Mohit Kumar

Department of Pharmaceutical sciences & Technology, Maharaja Ranjit Singh

Punjab Technical University, Bathinda, Punjab-151001

Email: maddysharma3303@gmail.com

Dishant Dhingra

M. Pharm

Department of Pharmaceutical sciences & Technology, Maharaja Ranjit Singh

Punjab Technical University, Bathinda, Punjab-151001

Email: dhingradishant3@gmail.com

Arzoo Pannu

Ph.D Research Scholar

School of Pharmaceutical sciences, Delhi Institute of Pharmaceutical Sciences and Research, New Delhi-017

Email: arzoopannu@gmail.com

Abstract---Curcumin is a natural yellow pigmented phytoconstituent isolated from the powdered rhizome of *Curcuma longa* linn belongs to family Zinzieberacea. *Curcuma longa* linn is commonly known as


turmeric (Haldi), a well-known Indian spice used as a traditional medicine from ancient era. In literature, there are different analytical methods for detection of curcumin such as HPLC, UPLC and LC-MS/MS. It contains a mixture of different phytoconstituents, therefore it is problematic to separate, identify and isolate pure curcumin from *Curcuma longa* linn. Hence, in the present review article we have tried to summarize the important analytical parameters required to analyze the curcumin compound in *Curcuma longa* linn. as it can help the authors to find out best readings to for identification and isolation of curcumin. This review gives a combined study of various analytical methods used for detection of curcumin with retention time, methods used, column used, wavelength, LOD, LOQ etc using different types of well-established HPLC, LCMS, UPLC methods which can be used for isolation, identification and quantitative estimation of curcumin in *Curcuma longa* linn.

Keywords---Curcumin, HPLC, UPLC, LC-MS, Turmeric

Introduction

Curcumin ((1,7-bis (hydroxyl-3-methoxyphenyl) -1,6-heptadiene-3,5-dione) is a natural yellow pigmented phytoconstituent isolated from the powdered rhizome of *Curcuma longa* linn (Family: Zinzieberacea) (Robinson et al., 2005). *Curcuma longa* linn is commonly known as turmeric (Haldi), which is a well-known Indian spice used as a medicine from ancient era. It has been recorded as precious medicinal spice in Ayurvedic and Unani system of medicine. It is cultivated in tropical and sub-tropical regions around the world but India is the largest producer of turmeric in the world (93.7% of total world production) and is cultivated in 150,000 hectares in India (Phukan et al., 2015). *Curcuma longa* contains different types of curcuminoids, out of which curcumin is reported as the most bio-active compound. These curcuminoids majorily contains 75% curcumin (curcuminI), 20% desmethoxycurcumin (curcuminII), 5% bisdesmethoxycurcumin (curcumin III), essential oils (2-4%), fixed oils (2-3%) (Jayaprakasha et al., 2002) and various volatile oils, including turmerone, atlantone, and zingiberone (Nair et al., 2021). It also contains sugars, proteins and resins.

In 1815, curcumin was isolated from *curcuma longa* for the first time and the characterization was done in 1870. In 1910, the structure of curcumin was elucidated for the first time by Polish scientists. Basically, curcumin is represent to 1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione, which is referred as “curcumin I”, having molecular formula $C_{21}H_{20}O_6$, molecular weight of 368.39 g/mol and melting point 183°C. In addition to “curcumin I”, there are two more compounds known as curcumin, which are curcumin II [demethoxycurcumin, 1-(4-hydroxy-3-methoxyphenyl)-7-(4-hydroxyphenyl)-1,6-heptadiene-3,5-dione] and curcumin III [bisdemethoxycurcumin, 1,7-bis(4-hydroxyphenyl)-1,6-heptadiene-3,5-dione].



(1E,6E)-1,7-bis(4-hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione
Curcumin

The various pharmacological activities of curcumin have been evaluated by many researchers around the world. Curcumin has been found as a significant therapeutic agent due to its great medicinal properties and is currently under clinical trials for treating various conditions i.e. pancreatic cancer, multiple myeloma, colon cancer, psoriasis, oral cancer, mastitis myelodysplastic syndromes, diabetes, alzheimer's disease, periodontal disease, pre-cancerous ulcer, recurrent aphthous stomatitis, and many other conditions etc. Traditionally, it has been used to treat a variety of illness, including hepatic disorders and jaundice, anorexia, rheumatism, and diabetic ulcers. As a polyphenolic compound, it exerts its pharmacological actions via modulating different signalling pathways and biological processes, resulting in antioxidant, anti-inflammatory, antibacterial, immunomodulatory, neuroprotective, anticarcinogenic, vasodilatory, chemoprotective, and antihyperglycemic potentials as shown in Fig. 1.




Fig. 1: Pharmacological activities of curcumin

Curcumin has been characterized as safe by the food and drug administration (FDA) USA, the Natural Health Product Directorate of Canada and the Joint FAO/WHO committee on food additives of the Food and Agriculture Organization/World Health Organization. The average intake of curcumin in Indian diet is approximately 2-2.5 g for 60 Kg individual, which corresponds to a daily intake of approximately 60-100 mg of curcumin (Goel et al., 2008).

The chemical formula of curcumin is 1,7-bis-(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-2,5-dione with molecular formula $C_{21}H_{20}O_6$ and its pK_a value of 8.54. Curcumin predominately resides in its keto-form in acidic and neutral conditions and solid phase. It is reported that the maximum absorption (λ_{max}) of curcumin in methanol occurs at 430 nm (Ansari et al., 2005).


The chemical structure of curcumin makes it less soluble in water at acidic and neutral pH but soluble in methanol, ethanol, dimethyl sulfoxide (DMSO) and acetone. It is a hydrophobic polyphenol and several studies have revealed extremely low water-solubility, low stability, rapid metabolism and poor absorption of these molecules that severely reduces its bioavailability (Mohanty et al., 2012). To overcome the problems, several drug delivery systems have been taken into consideration to provide longer circulation times and increased permeability.

Various HPLC, HPTLC, UPLC, LCMS/MS methods have been reported for the determination of curcumin in various sample matrixes. HPLC analysis of curcumin was first attempted by (Asakawa et al., 1981) using Nucleosil C18 column as stationary phase, a mixture of Acetonitrile: H₂O: Acetic acid (51: 49: 5) as mobile phase and benzyl benzoate as an internal standard. Many HPLC methods have been reported since then. These methods varied depending of different types of matrix, objective of analysis, sensitivity, detector and column used. For a particular HPLC analysis for curcumin, analyst should select chromatographic parameters judiciously based on earlier published methods. So, the objective of this review article is an in-depth analysis of published HPLC articles that will be useful to the readers for selection of various chromatographic parameters for their study.

Method

Reported HPLC Methods

The Chromatography method was performed for extract as per the method was reported in labs.



2.1. HPLC Instruments

Various type of HPLC instrument companies like Shimadzu HPLC instrument-LC-20AD (Japan) equipped with Rheodyne 7725i injection valve with a 20 μ L loop volume and Binary gradient pump, Plus Intelligent LC pump PU-2080 from Jasco (Tokyo, Japan) equipped with a Jasco UV-2075 Intelligent UV/Vis detector and a Rheodyne 7725 injector (Rheodyne, Cotati, CA, USA), A Waters 2695 Alliance HPLC system (Milford, MA, USA), A Agilent 1100 separation module with a photodiode array detector, A Varian® ProStar HPLC system, Agilent quaternary system with 1260 quat pump, injector, variable wavelength detector 1260 VWD VL and auto sampler 1260 ALS, Waters 625LC HPLC system.

2.2. Selection of Column

Various Columns are used in the HPLC system i.e. RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg) (Kadam et al., 2018), Agilent RPC18XDB column(4.6*150mm) (Wulandari et al., 2018), 25cm*4.6mm i.d. stainless steel analytical 5mm particle LiChrosphere-100 RP 18 packing column (Chaudary Hema et al.) , Waters Symmetry Shield (3.9*150 mm) (A.Radha et al.), 5 μ m C₁₈ , AlltactAlltima C₁₈ Column (150*4.6 mm) (Wichitnithadwisut et al.), Thermo Hypersil Gold column (250 mm \times 4.6 mm I.D.: 5 μ m) (Ang Fung Lee et al.).

2.3 Selection of Mobile phase

There are various mobile phases are used in the Hplc system i.e Acetonitrile:Acetic acid (49:51 v/v) (K.Prabaning dyah Niken et al.), Acetonitrile:Acetic acid (50:50 v/v) (Wulandari et al., 2018), Ethanol:Water:Acetonitril (80:10:10 v/v/v) (Thuane Castro Frabeldo Nascimen et al.), Ortho phosphoric acid: acetonitrile (65:35) (Chaudary Hema et al.), Aqueous acetic acid: methanol (15:85v/v) (Cheng et al., 2010) Acetonitril:Tetrahydrofuran:Water (35:20:45 v/v/v) (Han Yangrui et al.), Acetonitrile:water (60:40 v/v) (Hu liandong et al.), Acetonitril:Acetic acid (55:45 v/v) (A.Radha et al.), KH₂PO₄:Methanol (65:35 v/v) (A.Koray Mohamed et al.), Acetonitril:Methanol:Water (40:20:40 v/v/v) (Khalid Syed Haraaon).

2.4 Selection of Detector

These are various Detectors are used for the determination of Curcumin and their other constituents. These detectors are Jasco UV-2075 Intelligent UV/Vis detector, UV spectrophotometer, detector PDA,Detector (Waters 2998) and a fluorescence detector (Waters 2475), a ProStar 330 UV-Vis PDA spectrophotometric detector,Waters 2487 dual channel UV detector,9050 UV visible Detector, detector 1260 VWD VL,Photodiode array detector, Waters 486 tunable absorbance detector,UV vis dual wavelength spectrophotometric detector (SPD-10AVP).

Mobile Phase	Column	Detection	Retention time	LOD	LOQ	Reference
Ammonium acetate in water (pH-3): Acetonitrile (100%) 178	RP18 Water® X-Bridge (250*4.6mm i.d.; 5µm)	424nm	NA	2.23 µg/mL	6.774 µg/mL	(Kadam et al., 2018)
Acetonitrile:Acetic acid (49:51 v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	425 nm	NA	0.056 µg/mL	0.169 µg/mL	(Prabaningdyah et al., 2017)
Acetonitrile:Acetic acid (50:50 v/v)	Algilent RPC18XDB column(4.6*150mm)	425 nm	NA	0.004µg/mL	0.016 µg/mL	(Wulandari et al., 2018)
Water: Acetonitrile	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	254 nm	NA	0.3 µg/ml and 0.03 µg/ml	1 µg/ml and 0.1 µg/ml	(Gugulothu & Patravale, 2012)
Ethanol:Water: Acetonitril (80:10:10 v/v/v)	Algilent RPC18XDB column(4.6*150mm)	250 nm	0.96, 1.26, and 1.36 min	9.65 ng/ml	50 ng/ml	(Castro Frabel do Nascimento et al., 2012)
Ortho phosphoric acid: acetonitrile (65:35)	25cm*4.6mm i.d. stainless steel analytical 5mm particle LiChrosphere-100 RP 18 packing column containing	425 nm	13.09+/-1.01	245.5 55.30 ng/ml	491.1 165.87 ng/ml	Chaudhary et al .
Acetonitril:Water (50:50 v/v)	Stainless steel analytical 5mm particle LiChrosphere-100 RP 18 packing column containing	425 nm	13.6, 12.1 and 10.8 min	11.61 ng/ml	500 ng/ml 5 and 10 ng/ml	(Fonseca-Santos et al., 2017)
Aqueous acetic acid: methanol (15:85v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg) 25cm*4.6mm i.d.	420 nm	8.7 min	40.66, 49.38,29.28 pg.	134.18, 164.44, 97.50 pg	(H. Chen et al., 2009)
Acetonitril:Tetrahydrofuran:Water (35:20:45 v/v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	425 nm	6.5,11.5,19.9 mins	NA	5 & 10 ng/ml	(Han et al., 2011)
Acetonitril:Methanol:Deionized water:Acetic acid (41:23:36:1 v/v/v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	262 nm	NA	0.091 µg/ml	NA	(Heath et al., 2003)
Acetonitrile:water (60:40 v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	428 nm	NA	NA	NA	Hu liandong et.al
Acetonitril:Acetic acid (55:45 v/v)	Waters Symmetry Shield (3.9*150 mm),5µm C ₁₈	425 nm	0.176	0.356,1.80,0.074 ppm	1.078,5 .545,0.224 ppm	(Radha et al., 2016)
KH ₂ PO ₄ :Methanol (65:35 v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	288 nm	55 mins	0.0031,0.0035 µg/ml	0.0104, 0.0118 µg/ml	Koray Mohamed et.al
Acetonitril:	Waters	300 & 428	8.6 & 5.9 min	20 ng/ml	NA	Mazengshuan et.al


Citric buufer (55:45)	SymmetryShield 3.9*150 mm,5µm C ₁₈	nm				
Acetonitril: Water or Chloroform: Methanol	Waters µBondapak™ C ₁₈ (3.9 *300 mm)	265 nm	NA	2.5 ng/ml	NA	Indira PriyadarsiniKavirayani
Chloroform:Me thanol (48:2 v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	425 nm	NA	0.1 µg/spot	NA	(Paramasivam et al., 2008)
Aetonitril:Methanol (20:50 v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	332 nm	2.8 +/- 0.2, 5.1 +/-0.2 mins	NA	0.167,0.271 µg/ml	(Rathore et al., 2008)
Acetonitril:Met hanol:Water (40:20:40 v/v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	370 nm	6.728,6.166,5.66 mins	0.305 µg/ml	2 µg/ml	Khalid Syed Haraaon
Acetonitril:For mic acid (60:40 v/v)	RP18 Water® X-Bridge(250*4.6mm i.d.;5mcg)	NA	5.90,6.64,6.33 mins 13.6,12.1,10.8 mins	NA	1,0.50 ng/ml	Wang Xiu-Mei et.al
Acetonitril:Acetic acid (40:60 v/v)	AlltactAlltima C ₁₈ Column (150*4.6 mm)	425 nm	NA	0.90,0.84,0.08 µg/ml	2.73,2.53,0.23 µg/ml	WichitnithadWisut et.al
Acetonitrile: Acetic acid (40:60)	Thermo Hypersil Gold column (250 mm × 4.6 mm I.D.: 5 µm)	370 nm	32.195, 2.887 and 2.830	0.00488, 0.62500, 0.07813 and 0.03906 µg/mL	0.03906, 2.5000, 0.31250 and 0.07813 µg/mL	(Ang et al., 2014)

Later on, Tonnesen et al. reported a HPLC method which was able to separate three different components of curcuminoids. There are different detectors were used based on UV-Vis and on fluorescence. The fluorescence detector showed sensitivity that was 0-fold higher than the UV-Vis detector in detecting three components of curcuminoids (Curcumin, demethoxycurcumin, and bis – demethoxycurcumin). The five different eluents (combination of EtOH-water, MeOH-water, and butanol) as well as eight stationary phases were evaluated. It was found that the 1,3-diketone group in curcuminoids was adsorbed strongly to the salicylic acid column, hence the analysis was not reproducible.


Mobile Phase	Column	Detection	Retention time	LOD	LOQ	Reference
Toluene:chloroform: methanol (5:4:1 v/v/v) 180	TLC Aluminium Plates Precoated with a silica gel 60F ₂₅₄	NA	NA	50 ng/ml	200 ng/ml	(Ashraf et al., 2012)
N-hexane:ethylacetate: aceticacid:methanol (7:2:0.5:0.5 v/v/v/v)	Precoated TLC silica gel aluminium plates 60 F ₂₅₄ (20*10 cm),250µm thickness	404 nm	30 mins	18.31and 40.50ng/spot	55.50 and 122.74 ng/spot	(Kharat et al., 2017)
Toluene:Ethylacetate :Methanol (8:1:1 v/v/v)	TLC Aluminium Plates Precoated with a silica gel 60F ₂₅₄	293 nm	20 mins	26,31,29 ng/band	77,92,88 ng/band	(Siddiqui, 2015)
Dichloromethane: Methanol(99:1)	Precoated TLC silica gel aluminium plates 60 F ₂₅₄ (20*10 cm),250µm thickness	427 nm	NA	49 ng/spot	148 ng/spot	(Gantait et al., 2011)
Chloroform:Ethanol: Acetic Acid (95:4:5 v/v/v)	Precoated TLC silica gel aluminium plates 60 F ₂₅₄ (20*10 cm),250µm thickness	288 nm	10 mins	NA	NA	Srivastava et.al
Toluene: Ethylacetate: Formic acid (9:6:0.4 v/v/v)	TLC Aluminium Plates Precoated with a silica gel 60F ₂₅₄	200-700 nm	NA	NA	NA	N.TahaMahamoud et.al
Chloroform: Methanol (9.5:0.5)	Precoated Silica gel 60F ₂₅₄ Aluminium Plates(10*20cm)	421nm	NA	8ng/spot	25ng/spot	(Kandasamy & Moscow, 2013)
n-hexane:ethyl acetate:methanol: formic acid (8:2:1:2-3 drops v/v)	Precoated with silica gel 60F ₂₅₄ aluminium plates	421nm	NA	27.3ng	82.7ng	(Kekre & Walode, 2012)
Chloroform:ethyl acetate:formic acid:methanol (7.5:6:0.5:0.5)	Aluminium plates precoated with silicagel 60F ₂₅₄ (20*10cm) thickness 0.2 mm	322nm	NA	29ng	75ng	(Chavan et al., 2015)
Chloroform: methanol (48:2 v/v)	Glass backed silica gel 60GF ₂₅₄ (20*20cm) thickness 300µ	425nm	NA	NA	NA	(Paramasivam et al., 2008)
Chloroform: methanol (9:1)		420nm	NA	17.39ng/spot	52.71ng/spot	(Dixit et al., 2008)
Chloroform:methanol :glacial acetic acid (7:2:0.3)	Precoated with silica gel 60F ₂₅₄	366nm	NA	NA	NA	(Mishra et al., 2017)
Chloroform: ethyl acetate:formic acid (7.5:6:0.2)	Aluminium plates precoated with silica gel 60F ₂₅₄	540nm	NA	0.06ng/spot	0.2 ng/spot	(Vaykole et al., 2014)
Chloroform: methanol (98:2 v/v)	Lichro-sphere aluminium plates Si 60F ₂₅₄ (20*20cm)	366nm	NA	40ng	100ng	(Pathania et al., 2006)
Toluene:ethyl acetate: formic acid (4.5:3:0.2)	Silica gel 60 F ₂₅₄ percolated TLC plate (10*10cm) thickness 0.2mm	366nm	NA	16ng/spot	53.34 ng/spot	(Thakker et al., 2011)

Chloroform: ethyl acetate: formic acid (7.5:6:0.5)	Silica gel 60F254 TLC plates (20*10cm) thickness 0.2mm	254nm	NA	100ng/spot	300ng/spot	(Sonawane et al., 2011)
Toluene: ethyl acetate: methanol	Silica gel 60F254 HPTLC Plates	420nm	NA	25ng	NA	(J. K. Verma & Joshi, 2006)
Chloroform: methanol (9.25:0.75 v/v)	TLC aluminium plates precoated with silica gel 60F254	430nm	NA	8ng/spot	25ng/spot	(Ansari et al., 2005)
Toluene: ethyl acetate: methanol (8:1:1 v/v/v)	Precoated HPTLC Plates (20*10cm)	293nm	NA	26ng/band	77ng/band	(Siddiqui, 2015)

HPTLC method




LC-MS/MS method



Mobile Phase	Column	Detection/ Ion Transition	Retention time	LOD	LOQ	Reference
Acetic acid: Acetonitrile	ZorbaxEclipse C ₁₈ (4.6*150 mm)	425 nm	8.2 \pm 0.2 min and 1.4 \pm 0.1 min	NA	10 ng/ml	DalapathiGu gulothu et.al.pdf.in
Mehanol: Ammonium Acetate (80:20 v/v)	ZorbaxEclipse C ₁₈ (4.6*150 mm)	230 nm	30 mins	0.01, 0.25 μ g/ml	0.4, 1.2 μ g/ml	Memvanga Patrick et.al
Acetonitrile: 1mM Formic Acid (70:30 v/v)	Reverse phase c18 column (150*46mm) Zorbax	367-217 m/z	NA	1ng/ml	5ng/ml	(Yang et al., 2007)
Acetonitrile : water (70:30 v/v)	C18 phenomenex luna, 250mm*4.6mm	372-137 m/z	NA	NA	0.5ng/ml	(Liu et al., 2006)
Methanol:10mM Ammonium Formate(ph 3.0) (1:1)	Waters MS C18 column (2.1mm* 50mm, 3.5 μ m)	367-149 m/z	5.08 mins	NA	2.50ng/ml	(Kunati et al., 2018)
	Phenomenex Luna silica column (50mm*4.6mm* 5 μ m)	369-245 m/z	NA	1 μ g/l	10 μ g/l	(Gören et al., 2009)
0.1% Formic acid in water: Acetonitrile (15:85) (5:95) (25:75)	Agilent zorbax eclipse* DB C 18 column (3.5 μ m, 4.6*50 mm)	369-245-213 m/z	2.5min	0.1ng/ml	1ng/ml	(Li et al., 2018)
0.2% Formic acid : Acetonitrile (50:50)	Gilent poroshell SB C18(2.7 μ m, 4.6*150mm)	369-179 m/z	1.5min	0.5ng/ml	1ng/ml	(Z. Chen et al., 2017)
Acetonitrile: 0.2% formic acid (73:27 v/v)	Halo C18 column (2.7 μ m, 4.6*50mm)	369-177 m/z	1.5min	NA	1ng/ml	(Bhuket et al., 2016)
Acetonitrile: 0.1 formic acid (95:5) (5:95)	C18 column (100mm*2.1mm, 1.7 μ m)	369-177 m/z	4.1min	NA	5ng/ml	(Ma et al., 2015)
Acetonitrile: 0.1% formic acid (50:50) v/v	Sepax BR-C18 (5 μ m, 1.0*100m m)	369-285 m/z	2.2 min	NA	2.5ng/ml	(Ramalingam & Ko, 2014)
50%Acetonitrile: 0.1% formic acid (1:1)	Thermo Beta Basic C8 column (5 μ m, 2.1*50mm)	369-177 m/z	NA	1ng/ml	2ng/ml	(Cao et al., 2014)
0.1% formic acid with water: 1% formic acid with Acetonitrile (50:50) v/v	C8 column (2.1*50mm, 5 μ m)	425nm 545-351m/z	NA	NA	NA	(Zhongfa et al., 2012)

Acetonitrile: 10mM ammonium acetate buffer(ph 3.5) (8:20) v/v	Chromolith C18 (50mm*4.6, 5µm)	367-217 m/z	NA	NA	NA	(Kakkar et al., 2010)
Methanol: 0.05% acetic acid (80:20)	C18 100A column (250mm*4.6mm *5µm)	369.3-285 m/z	1.08min	NA	0.5ng/ml	(Khalil et al., 2013)
100% Methanol: 0.2% formic in water (ph 2.5) (10:90) (90:10)	Synergi™ Fusion-RP C18	NA	1.5min	NA	NA	(Bangphumi et al., 2016)

UPLC method



Mobile Phase	Column	Detection	Retention time	LOD	LOQ	Reference
Aqueous phosphoric acid:Acetonitril (34:66v/v)	Water Acquity® BEH C ₁₈ , h ₅₅ BEH C ₁₈ , h _{ss} T ₃ & BEH Shield RPC ₁₈ BEH RP ₁₈ (2.1&100 nm)	420 nm	1.576,1.43 5,1.30 mins	NA	NA	(Cheng et al., 2010)
5% Aqueous Acetonitril: Acetic Acid &Acetonitril	Water Acquity® BEH C ₁₈ , h ₅₅ BEH C ₁₈ , h _{ss} T ₃ & BEH Shield RPC ₁₈ BEH RP ₁₈ (2.1&100 nm)	426 nm	NA	2.5 ng/ml	NA	H.Marczylo Timothy et.al
2% v/v acetic acid in water: Acetonitrile (55:45) (20:80)	C18(1.7µm, 2.1*50mm)	400nm	1.8min	0.050µg/ml	0.075µg/ml	(Bhuket et al., 2020)
Acetonitrile: water	UPLC BEH C18 (1.7µm, 2.1*50mm)	360nm 435nm	NA	0.008ng/ml 0.003ng/ml	0.02ng/ml 0.01ng/ml	(Cui et al., 2017)
0.05% aqueous phosphoric acid: Acetonitrile (36:66) v/v	BEH Shield RP C18 (2.1*100mm, 1.7µm)	420nm	NA	40.66pg	134.18pg	(Cheng et al., 2010)
Acetonitrile:	C18	NA	NA	NA	NA	(Jude et al.,

Methanol with 10mM ammonium formate in water (70:30)	(2.1*50mm, 1.7µm)					2018)
Acetonitrile : 10mm ammonium formate buffer containing 0.1 % formic acid	BEH HILIC 18 (2.1*100mm, 1.7µm)	NA	NA	NA	NA	(S. Wang et al., 2018)
Formic acid : Acetonitrile (72:28) (10:90)	UPLC® BEH C18 (2.1*50mm, 1.7µm)	369-25 m/z	2.5min	NA	1ng/ml	(Ramadanty et al., 2019)
0.15% Formic acid: Acetonitrile (50:50) v/v	UPLC BEH C18 column 1.7µm, 2.1*100mm)	369-176.95	1.7min	NA	1ng/ml	(Hayun et al., 2018)
0.05% o- phosphoric acid: Acetonitrile (66:34) v/v	HSS-T3 C18 (1.8µm, 2.1*50mm)	425nm	NA	NA	NA	(Saralkar & Dash, 2017)
Acetonitrile: 10mM ammonium acetate buffer	ACQUITY UPLC® BEH C18 (100*2.1mm,	NA	NA	NA	NA	(L. Wang et al., 2019)

(ph-3.5) (85:15) (5:95) (60:40) (75:25) (50:50)	1.7µm)					
10%v/v Acetonitrile in water: Acetonitrile (70:30) (40:60)	ACQUITY UPLC BEH C18 (100*2.1mm, 1.7µm)	NA	NA	NA	NA	(Hocking et al., 2018)
Acetonitrile containing trifluoro acetic acid(0.1% w/v): Water (50:50)	UPLC BEH C18 column (100*2.1mm, 2µm)	425nm	NA	NA	NA	(Xue et al., 2018)
Acetonitrile: 5% Acetonitrile in water with 0.01% acetic acid (75:25) v/v	C18 column (50*2.1mm)	NA	NA	NA	2.0ng/ml	(M. K. Verma et al., 2013)v
Acetonitrile: 10mM ammonium	Waters ACQUITY UPLC BEH	367.0894- 217.1410 m/z	2.18min	0.318ng/ml	1ng/ml	(Ashraf et al., 2015)

formate (70:30)v/v	C18 (100*2.1mm, 1.7µm)					
Acetonitrile: 10mMammoniu m acetate (90:10)v/v	Waters ACQUITY UPLC™ BEH C18 (2.1*100mm, 1.7µm)	367-217.1 m/z	2.18min	NA	NA	(Anwar et al., 2015)
Acetonitrile: 0.1% formic acid with water (80:20)v/v	C18 (50*2.1mm, 3µm)	369-176.9 m/z	NA	NA	1.0ng/ml	(Yu et al., 2019)
0.1% Formic acid : Acetonitriloe (70:30) (25:85)	Waters ACQUITY UPLC™ BEH C18 (2.1*100mm, 1.7µm)	NA	NA	NA	NA	(J. Wang et al., 2015)
Acetonitrile: 2mM ammonium acetate (85:15) v/v	ACQUITY UPLC™ BEH C18 (2.1*100mm, 1.7µm)	367.08-217.03 m/z	1.72min	0.017ng/ml	0.05ng/ml	(Ahmad et al., 2016)

Conclusion

The present review article has summarized the important parameters required to analyse the curcumin compound in *Curcuma longa* linn., since it contain a mixture of different phytoconstituents, it is problematic to separate, identify and isolate pure curcumin from it. HPLC, LCMS, UPLC are reproducible and versatile

chromatographic techniques for the qualitative and quantitative estimation of any kind of secondary metabolites present in plants. Therefore, in this review, we have concluded that there are different types of well-established HPLC, LCMS, UPLC methods which can be used for isolation, identification and quantitative estimation of curcumin in *Curcuma longa* linn.

References

- Ahmad, N., Ahmad, R., & Ahmad, F. J. (2016). Stressed Kinetics and Pharmacokinetics of Curcumin Nanoemulsion Using Validated Ultrahigh-Performance Liquid Chromatography–Synapt Mass Spectrometry (UPLC–MS/MS–ESI–Q–TOF). *Iranian Journal of Science and Technology, Transactions A: Science*, 40(2), 109–123.
- Ang, L. F., Yam, M. F., Fung, Y. T. T., Kiang, P. K., & Darwin, Y. (2014). HPLC method for simultaneous quantitative detection of quercetin and curcuminoids in traditional chinese medicines. *Journal of Pharmacopuncture*, 17(4), 36.
- Ansari, M. J., Ahmad, S., Kohli, K., Ali, J., & Khar, R. K. (2005). Stability-indicating HPTLC determination of curcumin in bulk drug and pharmaceutical formulations. *Journal of Pharmaceutical and Biomedical Analysis*, 39(1–2), 132–138.
- Anwar, M., Ahmad, I., Warsi, M. H., Mohapatra, S., Ahmad, N., Akhter, S., Ali, A., & Ahmad, F. J. (2015). Experimental investigation and oral bioavailability enhancement of nano-sized curcumin by using supercritical anti-solvent process. *European Journal of Pharmaceutics and Biopharmaceutics*, 96, 162–172.
- Asakawa, N., Tsuno, M., Hattori, T., Ueyama, M., Shinoda, A., Miyake, Y., & Kagei, K. (1981). Determination of curcumin content of turmeric by high performance liquid chromatography. *Yakugaku Zasshi*, 101(4), 374–377.
- Ashraf, K., Mujeeb, M., Ahmad, A., Ahmad, N., & Amir, M. (2015). Determination of curcuminoids in *Curcuma longa* Linn. by UPLC/Q-TOF-MS: an application in turmeric cultivation. *Journal of Chromatographic Science*, 53(8), 1346–1352.
- Ashraf, K., Mujeeb, M., Ahmad, A., Amir, M., Mallick, M. N., & Sharma, D. (2012). Validated HPTLC analysis method for quantification of variability in content of curcumin in *Curcuma longa* L (turmeric) collected from different geographical region of India. *Asian Pacific Journal of Tropical Biomedicine*, 2(2), S584–S588.
- Bangphumi, K., Kittiviriyakul, C., Towiwat, P., Rojsitthisak, P., & Khemawoot, P. (2016). Pharmacokinetics of curcumin diethyl disuccinate, a prodrug of curcumin, in wistar rats. *European Journal of Drug Metabolism and Pharmacokinetics*, 41(6), 777–785.
- Bhuket, P. R. N., Niwattisaiwong, N., Limpikirati, P., Khemawoot, P., Towiwat, P., Ongpipattanakul, B., & Rojsitthisak, P. (2016). Simultaneous determination of curcumin diethyl disuccinate and its active metabolite curcumin in rat plasma by LC–MS/MS: Application of esterase inhibitors in the stabilization of an ester-containing prodrug. *Journal of Chromatography B*, 1033, 301–310.
- Bhuket, P. R. N., Wichitnithad, W., Sudtanon, O., & Rojsitthisak, P. (2020). A stability-indicating UPLC method for the determination of curcumin diethyl disuccinate, an ester prodrug of curcumin, in raw materials. *Heliyon*, 6(8), e04561.

- Cao, Y., Xu, R. X., & Liu, Z. (2014). A high-throughput quantification method of curcuminoids and curcumin metabolites in human plasma via high-performance liquid chromatography/tandem mass spectrometry. *Journal of Chromatography B*, 949, 70–78.
- Castro Frabel do Nascimento, T., Meza Casa, D., Facco Dalmolin, L., Cristina de Mattos, A., Maissar Khalil, N., & Mara Mainardes, R. (2012). Development and validation of an HPLC method using fluorescence detection for the quantitative determination of curcumin in PLGA and PLGA-PEG nanoparticles. *Current Pharmaceutical Analysis*, 8(4), 324–333.
- Chavan, A. K., Nirmal, S. A., & Pattan, S. R. (2015). Development and Validation of HPTLC Method to Detect Curcumin and Gallic Acid in Polyherbal Microencapsulated Formulation. *Journal of Liquid Chromatography & Related Technologies*, 38(12), 1213–1217.
- Chen, H., Zhu, H., Zheng, J., Mou, D., Wan, J., Zhang, J., Shi, T., Zhao, Y., Xu, H., & Yang, X. (2009). Iontophoresis-driven penetration of nanovesicles through microneedle-induced skin microchannels for enhancing transdermal delivery of insulin. *Journal of Controlled Release*, 139(1), 63–72.
- Chen, Z., Sun, D., Bi, X., Zeng, X., Luo, W., Cai, D., Zeng, Q., & Xu, A. (2017). Pharmacokinetic based study on “lagged stimulation” of *Curcuma Longae* Rhizoma-Piper nigrum couplet in their main active components’ metabolism using UPLC-MS-MS. *Phytomedicine*, 27, 15–22.
- Cheng, J., Weijun, K., Yun, L., Jiabo, W., Haitao, W., Qingmiao, L., & Xiaohe, X. (2010). Development and validation of UPLC method for quality control of *Curcuma longa* Linn.: Fast simultaneous quantitation of three curcuminoids. *Journal of Pharmaceutical and Biomedical Analysis*, 53(1), 43–49.
- Cui, X., Muhammad, I., Li, R., Jin, H., Guo, Z., Yang, Y., Hamid, S., Li, J., Cheng, P., & Zhang, X. (2017). Development of a UPLC-FLD method for detection of aflatoxin B1 and M1 in animal tissue to study the effect of curcumin on mycotoxin clearance rates. *Frontiers in Pharmacology*, 8, 650.
- Dixit, S., Khanna, S. K., & Das, M. (2008). A Simple 2-Directional High-Performance Thin-Layer Chromatographic Method for the Simultaneous Determination of Curcumin, Metanil Yellow, and Sudan Dyes in Turmeric, Chili, and Curry Powders. *Journal of AOAC International*, 91(6), 1387–1396.
- Fonseca-Santos, B., Gremião, M. P. D., & Chorilli, M. (2017). A simple reversed phase high-performance liquid chromatography (HPLC) method for determination of in situ gelling curcumin-loaded liquid crystals in in vitro performance tests. *Arabian Journal of Chemistry*, 10(7), 1029–1037.
- Gantait, A., Barman, T., & Mukherjee, P. K. (2011). *Validated method for estimation of curcumin in turmeric powder.*
- Goel, A., Jhurani, S., & Aggarwal, B. B. (2008). Multi-targeted therapy by curcumin: how spicy is it? *Molecular Nutrition & Food Research*, 52(9), 1010–1030.
- Gören, A. C., Çıkrıkçı, S., Çergel, M., & Bilsel, G. (2009). Rapid quantitation of curcumin in turmeric via NMR and LC–tandem mass spectrometry. *Food Chemistry*, 113(4), 1239–1242.
- Gugulothu, D. B., & Patravale, V. B. (2012). A new stability-indicating HPLC method for simultaneous determination of curcumin and celecoxib at single wavelength: an application to nanoparticulate formulation. *Pharm Anal Acta*, 3(4), 157.

- Han, Y., Zhu, J., Wang, Y., Wang, X., & Liao, Y. (2011). A simple RP-HPLC method for the simultaneous determination of curcumin and its prodrug, curcumin didecanoate, in rat plasma and the application to pharmacokinetic study. *Biomedical Chromatography*, 25(10), 1144–1149.
- Hayun, H., Rahmawati, R., Harahap, Y., & Sari, S. P. (2018). Development of UPLC–MS/MS method for quantitative analysis of curcumin in human plasma. *Acta Chromatographica*, 30(4), 207–211.
- Heath, D. D., Pruitt, M. A., Brenner, D. E., & Rock, C. L. (2003). Curcumin in plasma and urine: quantitation by high-performance liquid chromatography. *Journal of Chromatography B*, 783(1), 287–295.
- Hocking, A. J., Elliot, D., Hua, J., & Klebe, S. (2018). Administering fixed oral doses of curcumin to rats through voluntary consumption. *Journal of the American Association for Laboratory Animal Science*, 57(5), 508–512.
- Jayaprakasha, G. K., Jagan Mohan Rao, L., & Sakariah, K. K. (2002). Improved HPLC method for the determination of curcumin, demethoxycurcumin, and bisdemethoxycurcumin. *Journal of Agricultural and Food Chemistry*, 50(13), 3668–3672.
- Jude, S., Amalraj, A., Kunnumakkara, A. B., Divya, C., Löffler, B.-M., & Gopi, S. (2018). Development of validated methods and quantification of curcuminoids and curcumin metabolites and their pharmacokinetic study of oral administration of complete natural turmeric formulation (Cureit™) in human plasma via UPLC/ESI-Q-TOF-MS spectrometry. *Molecules*, 23(10), 2415.
- Kadam, P. V., Yadav, K. N., Bhingare, C. L., & Patil, M. J. (2018). Standardization and quantification of curcumin from *Curcuma longa* extract using UV visible spectroscopy and HPLC. *J Pharmacogn Phytochem*, 7(5), 1913–1918.
- Kakkar, V., Singh, S., Singla, D., Sahwney, S., Chauhan, A. S., Singh, G., & Kaur, I. P. (2010). Pharmacokinetic applicability of a validated liquid chromatography tandem mass spectroscopy method for orally administered curcumin loaded solid lipid nanoparticles to rats. *Journal of Chromatography B*, 878(32), 3427–3431.
- Kandasamy, J., & Moscow, S. (2013). Validated method for estimation of curcumin from different varieties of *curcuma longa*. *Int J Pharm Biosci*, 4, 1004–1010.
- Kekre, V. A., & Walode, S. G. (2012). Validated HPTLC method for estimation of curcumin content in dietary supplement formulation. *International Journal of Pharmaceutical Sciences and Research*, 3(10), 3796.
- Khalil, N. M., do Nascimento, T. C. F., Casa, D. M., Dalmolin, L. F., de Mattos, A. C., Hoss, I., Romano, M. A., & Mainardes, R. M. (2013). Pharmacokinetics of curcumin-loaded PLGA and PLGA-PEG blend nanoparticles after oral administration in rats. *Colloids and Surfaces B: Biointerfaces*, 101, 353–360.
- Kharat, S., Namdeo, A., & Mehta, P. (2017). Development and validation of HPTLC method for simultaneous estimation of curcumin and galangin in polyherbal capsule dosage form. *Journal of Taibah University for Science*, 11(5), 775–781.
- Kunati, S. R., Yang, S., William, B. M., & Xu, Y. (2018). An LC–MS/MS method for simultaneous determination of curcumin, curcumin glucuronide and curcumin sulfate in a phase II clinical trial. *Journal of Pharmaceutical and Biomedical Analysis*, 156, 189–198.

- Li, W., Yang, H., Buckley, B., Wang, L., & Kong, A.-N. (2018). A Novel Triple Stage Ion Trap MS method validated for curcumin pharmacokinetics application: A comparison summary of the latest validated curcumin LC/MS methods. *Journal of Pharmaceutical and Biomedical Analysis*, 156, 116–124.
- Liu, A., Lou, H., Zhao, L., & Fan, P. (2006). Validated LC/MS/MS assay for curcumin and tetrahydrocurcumin in rat plasma and application to pharmacokinetic study of phospholipid complex of curcumin. *Journal of Pharmaceutical and Biomedical Analysis*, 40(3), 720–727.
- Ma, W., Wang, J., Guo, Q., & Tu, P. (2015). Simultaneous determination of doxorubicin and curcumin in rat plasma by LC–MS/MS and its application to pharmacokinetic study. *Journal of Pharmaceutical and Biomedical Analysis*, 111, 215–221.
- Mishra, J., Tripathi, A., Misra, K., & Sarada, S. K. S. (2017). HPTLC: A tool for determination of curcumin in mammalian samples. *Int. J. Pharmacog. Phytochem. Res*, 9(4), 498–501.
- Mohanty, C., Das, M., & Sahoo, S. K. (2012). Emerging role of nanocarriers to increase the solubility and bioavailability of curcumin. *Expert Opinion on Drug Delivery*, 9(11), 1347–1364.
- Nair, A., Gopi, S., & Jacob, J. (2021). Bioavailability, pharmacokinetic, pharmacodynamic, and clinical studies of natural products on their antiinflammatory activities. In *Inflammation and Natural Products* (pp. 321–358). Elsevier.
- Paramasivam, M., Aktar, M. W., Poi, R., Banerjee, H., & Bandyopadhyay, A. (2008). Occurrence of curcuminoids in *Curcuma longa*: A quality standardization by HPTLC. ||| *Bangladesh Journal of Pharmacology* |||, 3(2), 55–58.
- Pathania, V., Gupta, A. P., & Singh, B. (2006). Improved HPTLC method for determination of curcuminoids from *Curcuma longa*. *Journal of Liquid Chromatography & Related Technologies*, 29(6), 877–887.
- Phukan, P., Bawari, M., & Sengupta, M. (2015). Promising neuroprotective plants from north-east India. *Int J Pharm Pharm Sci*, 7(3), 28–39.
- Prabaningdyah, N. K., Riyanto, S., Rohman, A., & Siregar, C. (2017). Application of HPLC and response surface methodology for simultaneous determination of curcumin and desmethoxy curcumin in *Curcuma* syrup formulation. *Journal of Applied Pharmaceutical Science*, 7(12), 58–64.
- Radha, A., Ragavendran, P., Thomas, A., & Kumar, D. S. (2016). A cost effective hplc method for the analysis of curcuminoids. *Hygeia JD Med*, 8, 1–15.
- Ramadanty, W. T., Arozal, W., Louisa, M., Soetikno, V., Purbadi, S., & Priyanto, P. (2019). Efficient validated method of UPLC-MS/MS to determine curcumin in rat plasma and ovarium. *Journal of Applied Pharmaceutical Science*, 9(01), 58–65.
- Ramalingam, P., & Ko, Y. T. (2014). A validated LC-MS/MS method for quantitative analysis of curcumin in mouse plasma and brain tissue and its application in pharmacokinetic and brain distribution studies. *Journal of Chromatography B*, 969, 101–108.
- Rathore, R., Jain, J. P., Srivastava, A., Jachak, S. M., & Kumar, N. (2008). Simultaneous determination of hydrazinocurcumin and phenol red in samples from rat intestinal permeability studies: HPLC method development and validation. *Journal of Pharmaceutical and Biomedical Analysis*, 46(2), 374–380.

- Robinson, T. P., Hubbard IV, R. B., Ehlers, T. J., Arbiser, J. L., Goldsmith, D. J., & Bowen, J. P. (2005). Synthesis and biological evaluation of aromatic enones related to curcumin. *Bioorganic & Medicinal Chemistry*, 13(12), 4007–4013.
- Saralkar, P., & Dash, A. K. (2017). Alginate nanoparticles containing curcumin and resveratrol: preparation, characterization, and in vitro evaluation against DU145 prostate cancer cell line. *AAPS Pharmscitech*, 18(7), 2814–2823.
- Siddiqui, N. A. (2015). Evaluation of thermo sensitivity of curcumin and quantification of ferulic acid and vanillin as degradation products by a validated HPTLC method. *Pakistan Journal of Pharmaceutical Sciences*, 28.
- Sonawane, S. D., Nirmal, S. A., Patil, A. N., & Pattan, S. R. (2011). Development and validation of HPTLC method to detect curcumin and gallic acid in polyherbal formulation. *Journal of Liquid Chromatography & Related Technologies*, 34(20), 2664–2673.
- Thakker, V. Y., Shah, V. N., Shah, U. D., & Suthar, M. P. (2011). Simultaneous estimation of gallic acid, curcumin and quercetin by HPTLC method. *Journal of Advanced Pharmacy Education & Research*, 1, 70–80.
- Vaykole, A. M., Nirmal, S. A., Jadhav, R. S., & Pattan, S. R. (2014). Development and validation of HPTLC method to detect curcumin, piperine, and boswellic acid in polyherbal transdermal patch. *Journal of Liquid Chromatography & Related Technologies*, 37(3), 367–378.
- Verma, J. K., & Joshi, A. V. (2006). Rapid HPTLC method for identification and quantification of curcumin, piperine and thymol in an ayurvedic formulation. *JPC-Journal of Planar Chromatography-Modern TLC*, 19, 398–400.
- Verma, M. K., Najar, I. A., Tikoo, M. K., Singh, G., Gupta, D. K., Anand, R., Khajuria, R. K., Sharma, S. C., & Johri, R. K. (2013). Development of a validated UPLC-qTOF-MS Method for the determination of curcuminoids and their pharmacokinetic study in mice. *Daru Journal of Pharmaceutical Sciences*, 21(1), 1–10.
- Wang, J., Ma, W., & Tu, P. (2015). The mechanism of self-assembled mixed micelles in improving curcumin oral absorption: in vitro and in vivo. *Colloids and Surfaces B: Biointerfaces*, 133, 108–119.
- Wang, L., Lu, N., Zhao, L., Qi, C., Zhang, W., Dong, J., & Hou, X. (2019). Characterization of stress degradation products of curcumin and its two derivatives by UPLC–DAD–MS/MS. *Arabian Journal of Chemistry*, 12(8), 3998–4005.
- Wang, S., Lin, Z., Su, K., Zhang, J., Zhang, L., Gao, Z., Wang, Z., Ma, J., & Wang, X. (2018). Effect of curcumin and pifenidone on toxicokinetics of paraquat in rat by UPLC–MS/MS. *Acta Chromatographica*, 30(1), 26–30.
- Wulandari, R., Martono, S., & Rohman, A. (2018). Liquid chromatography and fourier transform infrared spectroscopy for quantitative analysis of individual and total curcuminoid in *Curcuma longa* extract. *Journal of Applied Pharmaceutical Science*, 8(9), 107–113.
- Xue, J., Zhang, Y., Huang, G., Liu, J., Slavin, M., & Yu, L. L. (2018). Zein-caseinate composite nanoparticles for bioactive delivery using curcumin as a probe compound. *Food Hydrocolloids*, 83, 25–35.
- Yang, K.-Y., Lin, L.-C., Tseng, T.-Y., Wang, S.-C., & Tsai, T.-H. (2007). Oral bioavailability of curcumin in rat and the herbal analysis from *Curcuma longa* by LC–MS/MS. *Journal of Chromatography B*, 853(1–2), 183–189.

- Yu, W., Wen, D., Cai, D., Zheng, J., Gan, H., Jiang, F., Liu, X., Lao, B., Yu, W., & Guan, Y. (2019). Simultaneous determination of curcumin, tetrahydrocurcumin, quercetin, and paeoniflorin by UHPLC-MS/MS in rat plasma and its application to a pharmacokinetic study. *Journal of Pharmaceutical and Biomedical Analysis*, 172, 58–66.
- Zhongfa, L., Chiu, M., Wang, J., Chen, W., Yen, W., Fan-Havard, P., Yee, L. D., & Chan, K. K. (2012). Enhancement of curcumin oral absorption and pharmacokinetics of curcuminoids and curcumin metabolites in mice. *Cancer Chemotherapy and Pharmacology*, 69(3), 679–689.
- Suryasa, W., Sudipa, I. N., Puspani, I. A. M., & Netra, I. (2019). Towards a Change of Emotion in Translation of Kṛṣṇa Text. *Journal of Advanced Research in Dynamical and Control Systems*, 11(2), 1221-1231.