



# On an edge partition and root graphs of some classes of line graphs

K. Pravas<sup>a</sup>, A. Vijayakumar<sup>b</sup>

<sup>a</sup>Department of Mathematics, K. K. T. M. Government College, Pullut-680663, India.

<sup>b</sup>Department of Mathematics, Cochin University of Science and Technology, Cochin-682022, India.

pravask@gmail.com, vijay@cusat.ac.in

## Abstract

The Gallai and the anti-Gallai graphs of a graph  $G$  are complementary pairs of spanning subgraphs of the line graph of  $G$ . In this paper we find some structural relations between these graph classes by finding a partition of the edge set of the line graph of a graph  $G$  into the edge sets of the Gallai and anti-Gallai graphs of  $G$ . Based on this, an optimal algorithm to find the root graph of a line graph is obtained. Moreover, root graphs of diameter-maximal, distance-hereditary, Ptolemaic and chordal graphs are also discussed.

**Keywords:** line graph, Gallai, anti-Gallai, root graph

Mathematics Subject Classification : 05C99

DOI:10.5614/ejgta.2017.5.1.12

## 1. Introduction

The line graph  $L(G)$  of a graph  $G$  has as its vertices the edges of  $G$ , and any two vertices are adjacent in  $L(G)$  if the corresponding edges are incident in  $G$ . The Gallai graph  $Gal(G)$  [10, 15] of a graph  $G$  has as its vertices the edges of  $G$ , and any two vertices are adjacent in  $Gal(G)$  if the corresponding edges are incident in  $G$ , but do not span a triangle in  $G$ . The anti-Gallai graph  $antiGal(G)$  [13] of a graph  $G$  has as its vertices the edges of  $G$ , and any two vertices of  $G$  are adjacent in  $antiGal(G)$  if the corresponding edges are incident in  $G$  and lie on a triangle in  $G$ .

In [13] it is shown that the four color theorem can be equivalently stated in terms of anti-Gallai graphs. The problems of determining the clique number and the chromatic number of  $Gal(G)$  are

Received: 8 July 2016, Revised: 14 January 2017, Accepted: 26 February 2017.

NP-Complete[13]. In [3] it is shown that there are infinitely many pairs of non-isomorphic graphs of the same order having isomorphic Gallai graphs and anti-Gallai graphs. In [2] it is shown that the complexity of recognizing anti-Gallai graphs is NP-complete.

A graph  $H$  is forbidden in a graph family  $\mathcal{G}$ , if  $H$  is not an induced subgraph of any  $G \in \mathcal{G}$ . For any finite graph  $H$ , there exist a finite family of forbidden subgraphs for the Gallai graphs and the anti-Gallai graphs to be  $H$ -free [3]. However, both Gallai graphs and anti-Gallai graphs cannot be characterized using forbidden subgraphs [13].

The Gallai and the anti-Gallai graphs are spanning subgraphs of line graphs. In fact, they are complement to each other in  $L(G)$ . Therefore a natural question arises: is it possible to identify the edges of  $Gal(G)$  and  $antiGal(G)$  from  $L(G)$ ? A positive answer to this is given in this paper by introducing an algorithm to partition the edge set of a line graph into the edges of Gallai and anti-Gallai graphs, using the adjacency properties of common neighbors of the edges of a line graph in a hanging [8].

A graph  $G$  is a root graph of the line graph  $H$  if  $L(G) \cong H$ . The root graph of a line graph is unique, except for the triangle and  $K_{1,3}$  [16]. In this paper, using the edge-partition, an algorithm is obtained to find the root graph of a line graph. Also, the root graphs of diameter-maximal, distance-hereditary, Ptolemaic and chordal graphs are obtained.

Let  $H = (V, E)$  be a graph with vertex set  $V = V(H)$  and edge set  $E = E(H)$ . Let  $N(v)$  denote the set of all vertices adjacent to  $v$  and  $N_M(v) = M \cap N(v)$ , where  $M \subseteq V$ . The edge joining  $u$  and  $v$  is denoted by  $uv$ . The common neighbors of  $uv$  is  $N(u) \cap N(v)$  and  $N(uv) = N(u) \cup N(v)$ . The subgraph induced by  $\{v_1, v_2, \dots, v_k\} \subseteq V$  is denoted by  $\langle v_1, v_2, \dots, v_k \rangle$ . A clique is a complete subgraph of a graph. An edge clique cover of  $H$  is a family of cliques  $\mathcal{E} = \{q_1, q_2, \dots, q_k\}$  such that each edge of  $H$  is in at least one of  $E(q_1), E(q_2), \dots, E(q_k)$ .

A path on  $n$  vertices  $P_n$  is the graph with vertex set  $\{v_1, v_2, \dots, v_n\}$  and  $v_i v_{i+1}$  for  $i = 1, 2, \dots, n-1$  are the only edges. The distance between two vertices  $u$  and  $v$ , denoted by  $d(u, v)$ , is the length of a shortest  $u - v$  path in  $H$ . The diameter of  $H$ , denoted by  $d(H)$ , is the maximum length of a shortest path in  $H$ .

The join of two graphs  $G_1$  and  $G_2$ , denoted by  $G_1 \vee G_2$ , is the graph with vertex set  $V(G_1) \cup V(G_2)$  and  $E(G_1 \vee G_2) = E(G_1) \cup E(G_2) \cup \{uv : u \in V(G_1) \text{ and } v \in V(G_2)\}$ .

All graphs mentioned in this paper are simple and connected, unless otherwise specified. Also, all other basic concepts and notations not mentioned in this paper are from [4].

## 2. Adjacency properties of edges of $L(G)$

The hanging [8] of a graph  $H = (V, E)$ , with  $|V| = n$  and  $|E| = m$ , by a vertex  $z$  is the function  $h_z(x)$  that assigns to each vertex  $x$  of  $H$  the value  $d(z, x)$ . The  $i$ -th level of  $H$  in a hanging  $h_z$  is defined as  $L_i = \{x \in H : h_z(x) = i\}$ . A hanging can be obtained using a breadth first search(BFS) [1], which has a time complexity of  $O(m + n)$ .

For a vertex  $v$  in  $L_i$ , a supporter of  $v$  is a vertex in  $L_{i-1}$ , which is adjacent to  $v$ . A vertex in  $L_i$  is an ending vertex if it has no neighbors in  $L_{i+1}$ . An arbitrary supporter of  $v$  is denoted by  $S(v)$ . It is clear that any vertex  $v$  in the level  $L_i$  for  $i \geq 1$  has at least one supporter.

We use the following, well known, forbidden subgraph characterization of a line graph.

**Theorem 2.1.** [6] A graph  $H$  is a line graph if and only if the nine graphs in Fig 1 are forbidden subgraphs for  $H$ .




Figure 1. Forbidden Subgraphs of line graph

**Theorem 2.2.** Consider a hanging of a line graph  $H$  by an arbitrary vertex in  $H$  and let  $uv$  denote the edge joining  $u$  and  $v$  in the same level  $L_i$ . Then, the following statements hold

1. All common neighbors of  $uv$  in  $L_{i-1}$  are adjacent to each other.
2. All common neighbors of  $uv$  in  $L_{i+1}$  are adjacent to each other.
3. If  $uv$  has no common neighbor in  $L_{i-1}$ , then all the common neighbors of  $uv$  in  $L_i$  which are adjacent to all other neighbors of  $uv$  are adjacent to each other.
4. There is at most one common neighbor of  $uv$  in  $L_i$ , which is adjacent to all the neighbors of  $uv$  but not adjacent to the common neighbors of  $uv$  in  $L_{i-1}$  and  $L_i$ .

*Proof.*

1. Let  $x$  and  $x'$  be two (distinct) common neighbors of an edge  $uv$  in  $L_{i-1}$ , then  $i \geq 2$ . Assume that  $x$  and  $x'$  are not adjacent. Now, if  $x$  and  $x'$  have a common neighbor  $w$  in  $L_{i-2}$ , then

$\langle w, x, x', u, v \rangle \cong F_2$  in Fig 1 which contradicts the fact that  $H$  is a line graph. So, let  $w$  and  $w'$  be any two vertices in  $L_{i-2}$  adjacent to  $x$  and  $x'$  respectively. Then  $\langle w, w', x, x', u, v \rangle \cong F_7$  or  $F_4$  according as,  $w$  and  $w'$  are adjacent or not.

2. Let  $w$  and  $x$  be two common neighbors of an edge  $uv$  in  $L_{i+1}$ . Assume that  $x$  and  $w$  are not adjacent. Now, if  $z$  is a supporter of  $u$  in  $L_{i-1}$ , then  $\langle z, u, w, x \rangle \cong K_{1,3}$ , which is a contradiction.
3. Let  $uv$  has no common neighbor in the level  $L_{i-1}$  and hence  $i \geq 2$ . Let  $x$  and  $w$  be two common neighbors of  $uv$  in  $L_i$  which are adjacent to all the neighbors of  $uv$ . Assume that  $x$  and  $w$  are not adjacent. Now  $u$  and  $v$  cannot have a common supporter. So let  $z_1$  and  $z_2$  be two supporters of  $u$  and  $v$  respectively. Since  $z_1$  and  $z_2$  are neighbors of  $uv$ , both  $x$  and  $w$  are adjacent to them. Now, the vertices  $z_1, x, w$  and  $S(z_1)$  induce a  $K_{1,3}$  which is a contradiction.
4. Assume that  $x$  and  $w$  are two nonadjacent common neighbors of  $uv$  in  $L_i$  which are not adjacent to the common neighbors of  $uv$  but adjacent to all the other neighbors of  $uv$  in  $L_{i-1}$  and  $L_i$ . So, it is clear that  $i \geq 2$ . Let  $z$  be a common neighbor of  $uv$  in  $L_{i-1}$ . Now  $u$  must have at least one neighbor in  $L_{i-1}$  other than the common neighbors of  $uv$  in  $L_{i-1}$ , for otherwise, the vertices  $u, x, w$  and  $z$  induce a  $K_{1,3}$  which is a contradiction. Similar is the case for the vertex  $v$ . So let  $z_1$  and  $z_2$  be two neighbors (but not common neighbors) of  $u$  and  $v$  in  $L_{i-1}$  respectively. But, we have,  $\langle S(z_1), z_1, x, w \rangle \cong K_{1,3}$ , which is also a contradiction.

□

**Remark 2.1.** In fact the above theorem is applicable to a larger class of graphs than line graphs as only some of the forbidden sub graphs of line graphs are used in the proof.

### 3. Anti-Gallai triangles in $L(G)$

Let  $uvw$  be a triangle in  $L(G)$  and let  $\bar{u}, \bar{v}$  and  $\bar{w}$  be the edges in  $G$  representing the vertices  $u, v$  and  $w$  respectively in  $L(G)$ . If the edges  $\bar{u}, \bar{v}$  and  $\bar{w}$  induce a triangle in  $G$  then the triangle  $uvw$  in  $L(G)$  is referred to as an anti-Gallai triangle. All the triangles in  $antiGal(G)$  need not be an anti-Gallai triangle and the number of anti-Gallai triangles in  $L(G)$  is equal to the number of triangles in  $G$ . Since each edge of an anti-Gallai graph belongs to some anti-Gallai triangle, the set of all anti-Gallai triangles in  $L(G)$  induces  $antiGal(G)$ .

**Remark 3.1.** When a triangle  $uvw$  in  $L(G)$  is not an anti-Gallai triangle, the edges  $\bar{u}, \bar{v}$  and  $\bar{w}$  in  $G$  have a vertex in common.

**Lemma 3.1.** Consider a line graph  $H \not\cong K_3$ . If a triangle  $uvw$  in  $H$  is an anti-Gallai triangle, then for all  $x \in V(H) \setminus \{u, v, w\}$ , one of the following holds.

- a)  $\langle u, v, w, x \rangle \cong K_4 - e$
- b)  $\langle u, v, w, x \rangle$  is disconnected.

*Proof.* Let  $G$  be the graph such that  $L(G) \cong H$  and assume that the triangle  $uvw$  is an anti-Gallai triangle in  $H$ . Then the edges  $\bar{u}, \bar{v}$  and  $\bar{w}$  in  $G$  induce a triangle in  $G$ . Now corresponding to any vertex  $x$  in  $H$ , there is an edge  $\bar{x}$  in  $G$ . If  $\bar{x}$  is adjacent to the triangle  $\bar{u}\bar{v}\bar{w}$ , then  $\bar{x}$  is adjacent to exactly two of the edges of  $\bar{u}\bar{v}\bar{w}$  and hence  $\langle u, v, w, x \rangle \cong K_4 - e$  in  $H$ . If  $\bar{x}$  is not adjacent to the triangle  $\bar{u}\bar{v}\bar{w}$ , then  $\langle u, v, w, x \rangle$  is disconnected.  $\square$

**Lemma 3.2.** *If a triangle  $uvw$  is not an anti-Gallai triangle in a line graph  $H \cong L(G)$ , then there is at most one common neighbor  $z$  for an edge of  $uvw$  in  $H$  such that  $\langle u, v, w, z \rangle \cong K_4 - e$ .*

*Proof.* Let  $\bar{u}, \bar{v}$  and  $\bar{w}$  be the edges in  $G$ , representing the vertices  $u, v$  and  $w$  respectively in  $H$ . Let  $z$  be such that  $\langle u, v, w, z \rangle \cong K_4 - e$  in  $L(G)$  and let it be a common neighbor of  $uv$ . Then the edge  $\bar{z}$  in  $G$  is adjacent to both the edges  $\bar{u}$  and  $\bar{v}$  and not adjacent to  $\bar{w}$ . Clearly  $\bar{u}, \bar{v}$  and  $\bar{z}$  induce a triangle in  $G$  and hence  $uvw$  is an anti-Gallai triangle in  $L(G)$ . Now assume that  $z'$  is a vertex different from  $z$  such that it is a common neighbor of  $uv$  and  $\langle u, v, w, z' \rangle \cong K_4 - e$ . Then the vertices  $z$  and  $z'$  cannot be adjacent, otherwise  $\langle u, v, z, z' \rangle \cong K_4$  and by Lemma 3.1 it will contradict the fact that  $u, v, z$  is an anti-Gallai triangle. But, we have,  $\langle u, w, z, z' \rangle \cong K_{1,3}$  and hence  $H$  cannot be a line graph by Theorem 2.1.  $\square$

**Theorem 3.1.** *Consider a line graph  $H \not\cong K_3, K_4 - e, C_4 \vee K_1$  and  $C_4 \vee 2K_1$ . A triangle  $uvw$  in  $H$  is an anti-Gallai triangle if and only if  $\langle u, v, w, x \rangle \cong K_4 - e$  or disconnected for all  $x \in V(H) \setminus \{u, v, w\}$ .*

*Proof.* Let  $G$  be the graph such that  $L(G) \cong H$ . The necessary part of the theorem follows from Lemma 3.1.

Conversely, assume that  $uvw$  is a triangle in  $H$  such that  $\langle u, v, w, x \rangle \cong K_4 - e$  or disconnected for all  $x \in V(H)$  and that  $uvw$  is not an anti-Gallai triangle. Then the edges  $\bar{u}, \bar{v}$  and  $\bar{w}$  induce a  $K_{1,3}$  in  $G$ . Note that any vertex which induces a  $K_4 - e$  with the triangle  $uvw$  is adjacent to exactly two vertices among  $u, v$  and  $w$ . Now, since  $H$  is connected and not a  $K_3$ , there is a vertex  $x$  adjacent to the triangle  $uvw$ . Assume that  $x$  is adjacent to  $u$  and  $w$ . Then in  $G$ ,  $\bar{u}, \bar{v}$  and  $\bar{x}$  induce a triangle so that  $uw$  is an anti-Gallai triangle. Since  $H \not\cong K_4 - e$  and also connected, there is a vertex  $y$  adjacent to at least one of the vertices  $u, v, w$  and  $x$ . If there is no vertex adjacent to the triangle  $uvw$ , then it must be adjacent to  $x$  alone, which is a contradiction to the fact that  $uw$  is anti-Gallai triangle. So let  $y$  be adjacent to  $uvw$ . By Lemma 3.2  $y$  cannot be adjacent to  $u$  and  $w$ . So let  $y$  be adjacent to  $v$  and  $w$ . Now we have  $vw$  is also an anti-Gallai triangle. But, since  $H \not\cong C_4 \vee K_1$  and connected, using the same arguments as before, we have a vertex  $z$  adjacent to the triangle  $uvw$  again. The only possibility then is that  $z$  is adjacent to the vertices  $u$  and  $v$ . Now we show that there are no more vertices possible in  $H$ . If not, let  $p$  be a vertex in  $H$  different from  $u, v, w, x, y$  and  $z$ . But, by Lemma 3.2, the vertex  $p$  cannot be adjacent to  $uvw$ . Now if  $p$  is adjacent to  $x$ , it must be adjacent to  $u$  or  $w$  as  $uw$  is an anti-Gallai triangle, which again is not possible. Similarly,  $p$  cannot be adjacent to  $y$  and  $z$ . Hence no such vertex  $p$  can be adjacent to any of the vertices  $u, v, w, x, y$  and  $z$ . So such a vertex does not exist in  $H$ , as  $H$  is a connected graph. Now we have  $H \cong \langle u, v, w, x, y, z \rangle \cong C_4 \vee 2K_1$ , which is a contradiction.  $\square$

We observe that it is possible to suitably re-label the edges in the root graph of  $C_4 \vee K_1$  so that no triangles in  $C_4 \vee K_1$  can be claimed to be an anti-Gallai triangle, see Figure 2. It can be seen




Figure 2. Two possible labellings of  $K_4 - e$  and its line graph  $C_4 \vee K_1$

that  $K_4 - e$  and  $C_4 \vee 2K_1$  also have this property. Theorem 3.1 shows that these three graphs are the only exceptions (the graph  $K_3$  is excluded as it is a trivial case with 3 vertices). Hence, the graphs  $K_4 - e$ ,  $C_4 \vee K_1$  and  $C_4 \vee 2K_1$  are excluded in the following discussions.

**Definition 1.** A triangle in a hanging of a line graph is an  $L\Delta$  ( $M\Delta$ ,  $R\Delta$ ) if it is an anti-Gallai triangle and it is induced by two vertices in one level and one vertex from the lower (same, higher) level of the ordering.

We can see that any anti-Gallai triangle is either an  $L\Delta$ ,  $M\Delta$  or  $R\Delta$  in a hanging of  $L(G)$




Figure 3. A graph and the hanging of its line graph by vertex  $f$ . The dotted lines show an  $L\Delta fgh$ ,  $R\Delta hij$  and an  $M\Delta abc$

**Theorem 3.2.** Let  $uv$  be an edge in any level of a hanging of  $H \cong L(G)$  by an arbitrary vertex in  $H$ , then

1.  $uv$  cannot be an edge of an  $L\Delta$  in any level  $L_i$  for  $i > 1$ .
2.  $uv$  cannot be an edge of an  $M\Delta$  in  $L_1$ .
3. If  $uv$  is an edge in an  $M\Delta$  then  $uv$  cannot be an edge of an  $L\Delta$ .
4. If  $uv$  is an edge in an  $M\Delta$  then  $uv$  cannot be an edge of an  $R\Delta$ .
5. If  $uv$  is an edge in an  $L\Delta$  then  $uv$  cannot be an edge of an  $R\Delta$ .
6.  $uv$  can be an edge of at most one  $L\Delta$  or  $R\Delta$  or  $M\Delta$ .

*Proof.*

1. Let  $uv$  be an edge in an  $L_i$  for  $i > 1$  and let it belong to an  $L\Delta uvx$ , where  $x \in L_{i-1}$ . Let  $w$  be the vertex in  $L_{i-2}$  which is adjacent to  $x$ . Then  $\langle w, x, u, v \rangle$  induces a subgraph which is neither a  $K_4 - e$  nor disconnected, which is a contradiction.
2. Let  $uvx$  be an  $M\Delta$  in  $L_1$  and  $z$  be the vertex, from where the hanging of  $H$  being considered. Then  $d(z) \geq 3$  and  $\langle z, x, u, v \rangle$  induce a  $K_4$  and hence  $uvx$  cannot be an anti-Gallai triangle, which is a contradiction.
3. Let  $uv$  be an edge in  $L\Delta$  then  $uv$  is in  $L_1$  by (1) and hence  $uv$  cannot be an edge of an  $M\Delta$  by (2).

From (3) and Theorem 3.1, it follows that anti-Gallai triangles of a graph cannot share an edge in a line graph. Hence the proof of (4) to (6) follows.

□

Now, Lemma 3.3 follows.

**Lemma 3.3.** *Exactly one triangle of a  $K_4 - e$  in a line graph is an anti-Gallai triangle.*

From Theorems 2.2 and 3.1, we have the following propositions.

**Proposition 3.1.** *The edge  $uv$  is in an  $L\Delta$ , with both its ends in the same level of a hanging of a line graph if and only if it satisfies the following conditions*

1. *Each vertex in  $L_1$  is either adjacent to  $u$  or  $v$  but not to both.*
2. *Each neighbor of  $uv$  in  $L_2$  is a common neighbor of  $uv$ .*

**Proposition 3.2.** *The edge  $uv$  is in an  $M\Delta$  in a hanging of a line graph if and only if it satisfies the following conditions*

1. *The edge  $uv$  has a common neighbor  $x$  in  $L_i$  which is not adjacent to the other common neighbors of  $uv$  in  $L_{i-1}$  and  $L_i$ .*
2. *Either  $u$  or  $v$  is adjacent to each neighbor of  $x$ .*
3. *Each non neighbor of  $x$  is either a common neighbor of  $uv$  or not a neighbor of  $uv$ .*

**Proposition 3.3.** *The edge  $uv$  is in an  $R\Delta$  with both its ends in the  $i^{\text{th}}$  level of a hanging of a line graph if and only if it satisfies the following conditions*

1. *The edge  $uv$  has exactly one common neighbor  $x$  in  $L_{i+1}$ .*
2. *The vertex  $x$  is an ending vertex.*
3. *Either  $u$  or  $v$  is adjacent to each neighbor of  $x$ .*
4. *Each non neighbor of  $x$  in  $L_{i-1} \cup L_i$  is either a common neighbor of  $uv$  or not a neighbor of  $uv$ .*

#### 4. Partitioning the edges of a line graph

We now provide an algorithm to partition the edge set of a line graph into edge sets of its Gallai and anti-Gallai graphs. The following three tests checks whether an edge  $uv \in L_i$  belongs to an  $L\Delta$ ,  $M\Delta$  or  $R\Delta$ .

##### Algorithm 1. $L\Delta$ test

1. If  $i \neq 1$  go to step 7.
2. Find  $N(u)$  and  $N(v)$ .
3. If  $N_{L_i}(u) \cup N_{L_i}(v) \neq L_i$  then go to step 7.
4. If  $N_{L_i}(u) \cap N_{L_i}(v) \neq \emptyset$  then go to step 7.
5. If  $N_{L_{i+1}}(u) \neq N_{L_{i+1}}(v)$  then go to step 7.
6. Triangle  $uvz$  is an  $L\Delta$ .
7. The edge  $uv$  is not in  $L\Delta$ .

##### Algorithm 2. $M\Delta$ test

1. If  $i = 1$  go to step 9.
2. Find the set  $C$  of common neighbors  $w_j$  of  $uv$  in  $L_i$ . If  $C = \emptyset$ , go to step 9.
3. Find the set  $B$  of common neighbors  $x_j$  of  $uv$  in  $L_{i-1}$  and  $L_{i+1}$ .
4. For each  $x_j \in B$ , delete the members of the set  $N_C(x_j)$  from  $C$ . If  $C = \emptyset$  go to step 9.
5. For each  $w_j$ , if  $|N_C[w_j]| > 1$ , delete the members of the set  $N_C[w_j]$ . If  $|C| \neq 1$  go to step 9.
6. Find the set  $N(uv)$  in  $H$ .
7. If  $|N_C(y_j)| = 1$ , for each  $y_j \in N(uv) \setminus (B \cup C)$ , go to step 8. Else go to step 9.
8. Triangle  $uvx$  is an  $M\Delta$ .
9. The edge  $uv$  is not in  $M\Delta$ .

##### Algorithm 3. $R\Delta$ test

1. Find the set  $C_R$  of common neighbors of  $uv$  in  $L_{i+1}$ .
2. If  $|C_R| \neq 1$  go to step 7. Else choose the common neighbor of  $uv$  in  $L_{i+1}$  as  $x$ .
3. If the vertex  $x$  is not an ending vertex, go to step 7.

4. Either  $u$  or  $v$  is adjacent to each neighbor of  $x$ . Else go to step 7.
5. Each non neighbor of  $x$  is either a common neighbor of  $uv$  or not a neighbor of  $uv$ . Else go to step 7.
6. Triangle  $uvx$  is an  $R\Delta$ .
7. The edge  $uv$  is not in  $R\Delta$ .

Given a line graph  $H \cong L(G)$ , obtain a hanging  $h_z$  by an arbitrary vertex  $z$ . Consider all the edges starting from a vertex  $u$  in  $L_1$ . For each edge of the form  $uv$  for some  $v \in L_1$ , apply tests 1, 2 and 3 one by one. Choose another edge whenever an anti-Gallai triangle is found or when all the tests fail. When all the edges in a level are considered, go to the next level and repeat the procedure. This algorithm ends when all the edges in the last level of the hanging are considered and uses a time complexity of  $O(m)$ .

We now observe that in a line graph  $L(G)$ , any edge that is in the edge set of  $antiGal(G)$  belongs to some anti-Gallai triangle. Hence the set of all the edges of the anti-Gallai triangles gives the edge set of  $antiGal(G)$  and the remaining edges of the  $L(G)$  corresponds to the edge set of  $Gal(G)$ .

## 5. An algorithm to find the root graph of a line graph

An optimal algorithm to recognize a line graph and out put its root graph can be seen in [14], the time complexity of which is  $O(n) + m$ . Using the above edge partition, an algorithm, which uses a time complexity of  $O(m) + O(n)$ , is provided to find the root graph of a line graph  $H$ . The same algorithm can be used as a recognition algorithm for line graphs. For this, applying the above three tests for the edges in an arbitrary graph, we call a triangle type I if it belongs to the category of anti-Gallai triangles and type II otherwise.

### Algorithm 4. Root graph of a line graph

Consider a connected graph  $H = (V, E)$  with  $|V| = n$ ,  $|E| = m$  and its hanging  $h_z$ , by an arbitrary vertex  $z$ .

Let  $M = \{z, u\}$ , where  $u$  is a neighbor of  $z$ . Let  $G$  be a path on three vertices with  $V(G) = \{\{z\}, \{z, u\}, \{u\}\}$  and  $E(G) = \{(\{z\}, \{z, u\}), (\{z, u\}, \{u\})\}$ . Here the labels of vertices of  $G$  are represented as sets which can be re-labeled, in the steps of the following algorithm, using set operations.

1. Choose a vertex  $v$  from  $V(H) \setminus M$  with  $N_M(v) \neq \emptyset$ .
2. If  $v$  induces a clique in  $N_M(v)$  and does not induce a type I triangle go to step 3. Else go to step 4.
3. Make  $V(G) = V(G) \cup \{v\}$ , and join  $\{v\}$  with a vertex  $C \in V(G)$ , where  $C = N_M(v)$ , and make  $M = M \cup \{v\}$  and  $C = C \cup \{v\}$ . If no such vertex  $C$  exists, go to step 4.

4. Find two vertices  $A$  and  $B$  in  $V(G)$  such that  $A \cup B = N_M(v)$  and make  $M = M \cup \{v\}$ ,  $A = A \cup \{v\}$  and  $B = B \cup \{v\}$ . Go to step 1.

The algorithm ends whenever  $M = V(H)$  or there does not exist  $C$  or  $A$  and  $B$  as required. Here the graph  $G$  represents the root graph of the line graph  $H$  and in the latter case it can be concluded that the graph  $H$  is not a line graph of any graph.

The correctness of the algorithm can be verified with the help of the following theorem due to Krausz [12].

**Theorem 5.1.** *A graph  $H$  is a line graph if and only if it has an edge clique cover  $\mathcal{E}$  such that both the following conditions hold:*

1. *Every vertex of  $H$  is in exactly two members of  $\mathcal{E}$ .*
2. *Every edge of  $H$  is in exactly one member of  $\mathcal{E}$ .*

Since the vertex labels of  $G$  are represented as sets, a vertex in  $\langle M \rangle$  is an element of some vertex label(set), of  $G$ . Here the elements of each vertex label in  $V(G)$  induce a clique in  $\langle M \rangle$  of  $H$ , since  $x, y$  are in a vertex label of  $G$  if and only if  $x$  and  $y$  are adjacent in  $\langle M \rangle$  of  $H$ . Now from the construction of  $G$ , each vertex of  $\langle M \rangle$  is an element of exactly two vertex labels of  $G$  and also any adjacent vertices in  $\langle M \rangle$  belong to a vertex label of  $G$ . Now  $V(G)$  gives an edge clique cover of  $\langle M \rangle$  which satisfies the two conditions given in Krausz's theorem. Hence the algorithm obtains a graph  $G$  with  $L(G) \cong H$  if and only if  $M = V(H)$ .

We now provide the difference between our algorithm and the algorithm in [14].

Given a graph  $H$ , the algorithm in [14] assumes that  $H$  is a line graph and defines a graph  $G$  such that  $H$  is necessarily the line graph of  $G$ . A comparison of  $L(G)$  and  $H$  is then made to check whether the given graph is actually a line graph. The algorithm starts with two adjacent basic nodes, labeled 1-2 and 2-3, and labels the vertices in  $H$ , on the go, depending on their adjacency. The algorithm proceeds to determine all connections in  $G$  corresponding to a clique, containing the basic nodes in  $H$ , simultaneously finding an anti-Gallai triangle  $\{1-2, 2-3, 1-3\}$ , if it exists. In each step, the cliques sharing the vertices, which are already worked out, are considered and the algorithm finally outputs a labeled graph  $G$ .

In our algorithm, the types of triangles are found using the first three algorithms, the time complexity of which is calculated as follows. We can see that a hanging of the graph  $H$  can be obtained in  $O(m + n)$  steps. In each of the algorithms 1, 2 and 3 only a subset of  $E(H)$  are considered (as edges between the levels are not included) and the algorithm 4, which assumes that algorithms 1, 2 and 3 are already done, finishes in  $O(n)$  steps. Hence using these algorithms the root graph of a line graph can be obtained in  $O(m) + O(n)$  steps. It can be noted, as a consequence of Theorem 3.1, that irrespective of the starting set  $M$  of nodes, any pre-labeled line graph  $H$  with more than four vertices gives a uniquely labeled root graph  $G$ .

## 6. Root graphs of diameter-maximal line graphs

A graph  $G$  is diameter-maximal [7], if for any edge  $e \in E(\overline{G})$ ,  $d(G + e) < d(G)$ .

**Theorem 6.1.** [7] A connected graph  $G$  is diameter-maximal if and only if

1.  $G$  has a unique pair of vertices  $u$  and  $v$  such that  $d(u, v) = d(G)$ .
2. The set of nodes at distance  $k$  from  $u$  induce a complete sub graph.
3. Every node at distance  $k$  from  $u$  is adjacent to every node at distance  $k + 1$  from  $u$ .

**Lemma 6.1.** Let  $G$  be a diameter-maximal line graph and  $u, v$  be two vertices of  $G$  with  $d(u, v) = d(G)$ . Let  $L^* = (|L_0|, |L_1|, \dots, |L_d|)$  be the sequence generated from the hanging  $h_u$ . Then,  $|L_i| \leq 2$  for  $i = 0, 1, \dots, d$ .

*Proof.* Clearly  $|L_0| = |L_d| = 1$  in  $L^*$ . If possible, let  $u, v$  and  $w$  be three vertices in  $L_i$  for some  $i$  for  $0 < i < d$ . By Theorem 6.1,  $\langle u, v, w \rangle \cong K_3$  and there exist vertices  $x$  in  $L_{i-1}$  and  $y$  in  $L_{i+1}$  such that  $u, v$  and  $w$  are adjacent to both  $x$  and  $y$ . But, then,  $\langle x, u, v, w, y \rangle \cong F_3$  which is a contradiction.  $\square$

A sequence  $S$  is forbidden in  $L^*$  if the consecutive terms of  $S$  do not appear consecutively in  $L^*$ .

**Theorem 6.2.** For every  $d \geq 3$ , there exists three diameter-maximal line graphs with diameter  $d$ .

*Proof.* First, we show that the sequence  $(a_1, a_2, 2, a_3, a_4)$ , where  $a_i \in \{1, 2\}$ , is forbidden in  $L^*$ . For, assuming the contrary, let  $|L_i| = 2$  for some  $i$ ,  $2 \leq i \leq d - 2$ , and  $L_i = \{v_1, v_2\}$ . Let  $v_3, v_4, v_5$  and  $v_6$  be arbitrary vertices in  $L_j$ , for  $j = i - 2, i - 1, i + 1$  and  $i + 2$  respectively. But  $\langle v_1, \dots, v_6 \rangle \cong F_4$  which is a contradiction.

Applying the same argument, we see that the sequences  $(a_1, a_2, 2, 2)$ ,  $(2, 2, a_1, a_2)$  and  $(2, 2, 2)$  are also forbidden in  $L^*$ , so that the integer 2 appears at most twice in  $L^*$  and hence either (i)  $|L_1| = |L_{d-1}| = 2$ , (ii)  $|L_1| = 2$  or (iii) all the entries of  $L^*$  are 1. Note that the case when  $L^*$  has  $|L_{d-1}| = 2$  is not considered, as it is similar to (ii). Hence there are only three possible sequences of  $L^*$  when  $d \geq 3$ . As the three sequences are different and the pair  $(u, v)$  in Theorem 6.1 is unique, there exist exactly three diameter-maximal line graphs.  $\square$

**Corollary 6.1.** The root graphs of diameter-maximal line graphs with diameter  $d$  are of the form  $G$  in Table 1.




Diameter of $L(G)$	$d = 1$	$d = 2$	$d \geq 3$
$G$			

Table 1. Graph  $G$ , for Corollary 6.1

## 7. Root graphs of DHL graphs

A graph  $G$  is distance-hereditary if for any connected induced subgraph  $H$ ,  $d_H(u, v) = d_G(u, v)$ , for any  $u, v \in V(H)$ . A detailed study can be seen in [5]. A graph  $G$  is chordal if every cycle of length at least four in  $G$  has an edge(chord) joining two non-adjacent vertices of the cycle [4]. A graph is Ptolemaic if it is both distance-hereditary and chordal [11].

In this section, the family of root graphs of distance-hereditary line (DHL) graphs is obtained. The root graphs of chordal and Ptolemaic graphs are also discussed.

**Theorem 7.1.** [5] *Let  $G$  be a connected graph. Then  $G$  is distance-hereditary if and only if the graphs of Fig 4 and the cycles  $C_n$  with  $n \geq 5$  are forbidden subgraphs of  $G$ .*




Figure 4. The graphs for Theorem 7.1: house, domino and gem graphs

**Theorem 7.2.** [11] *Let  $G$  be a graph. The following conditions are equivalent*

1.  $G$  is a Ptolemaic graph
2.  $G$  is distance-hereditary and chordal
3.  $G$  is chordal and does not contain an induced gem

A vertex  $v$  is simplicial if  $N(v)$  is a clique. The ordering  $\{v_1, \dots, v_n\}$  of the vertices of  $H$  is a perfect elimination ordering if, for all  $i \in \{1, \dots, n\}$ , the vertex  $v_i$  is simplicial in  $H_i = \langle v_i, \dots, v_n \rangle$ .

**Theorem 7.3.** [9] *Let  $G$  be a graph. The following statements are equivalent:*

1.  $G$  is a chordal graph.
2.  $G$  has a perfect elimination ordering. Moreover, any simplicial vertex can start a perfect elimination ordering.

**Theorem 7.4.** *In a DHL graph if a vertex is adjacent to at least one vertex in a  $C_4$  then it must be adjacent to all the vertices of that  $C_4$  and to no other vertices in the graph.*

*Proof.* Let  $H$  be a DHL graph which contains a  $C_4$  and let a vertex  $u$  be adjacent to at least one vertex of the  $C_4$ . If  $u$  is adjacent to exactly one vertex of  $C_4$  then a  $K_{1,3}$  is formed in  $H$ , which is a contradiction. Let  $u$  be adjacent to exactly two vertices of  $C_4$ . Then either a house, when  $u$  is adjacent to two adjacent vertices of  $C_4$ , or a  $K_{1,3}$ , when  $u$  adjacent to two non-adjacent vertices of

$C_4$  is formed, which is also a contradiction. Since an  $F_2$  is obtained when  $u$  is adjacent to three vertices of a  $C_4$ ,  $u$  must be adjacent to all the four vertices of the  $C_4$ .

Next we show that two adjacent vertices can not be made adjacent to a  $C_4$  in  $H$ . For, otherwise each of the two vertices must be adjacent to all the vertices of  $C_4$  and hence induces  $C_4 \vee K_2$ . But a copy of  $F_3$  is induced in  $C_4 \vee K_2$ , which is a contradiction. If only one vertex of two adjacent vertices is adjacent to  $C_4$ , a  $K_{1,3}$  is induced in  $H$  which is also a contradiction.  $\square$

**Corollary 7.1.** *A DHL graph contains at most one  $C_4$ .*

**Corollary 7.2.** *The root graphs of DHL graphs which contain a  $C_4$  are  $K_4$ ,  $K_4 - e$  and  $C_4$ .*

*Proof.* The proof is complete as we see from Corollary 7.1 that the only DHL graphs which contain a  $C_4$  are  $C_4 \vee 2K_1$ ,  $C_4 \vee K_1$  and itself.  $\square$

As there are only three DHL graphs containing a  $C_4$ , we restrict our discussion in the following sections to DHL graphs not containing  $C_4$ 's.

If  $H$  is a DHL graph containing no anti-Gallai triangle then its root graph contains no triangles. Also, a DHL graph is  $C_n$ -free,  $n \geq 5$ . Now, together with Corollary 7.2, we have the following result.

**Theorem 7.5.** *Let  $H \not\cong C_4$  be a DHL graph not containing an anti-Gallai triangle, then  $H$  is a line graph of a tree.*

**Lemma 7.1.** *An anti-Gallai triangle in a DHL graph has a vertex of degree two.*

*Proof.* Let  $uvx$  be an anti-Gallai triangle in a DHL graph  $H \not\cong K_3$ . Then  $uvx$  is in some  $K_4 - e$  in  $H$ . Let  $uvy$  be a triangle such that  $u, x, y, w \cong K_4 - e$ . We now show that degree of the vertex  $x$  is two. Consider  $h_x$ , we just need to show that  $L_1$  contains no vertices other than  $u$  and  $v$ . For, let  $w$  be a vertex in  $L_1$ . Then  $wx$  is an edge and, by Theorem 3.1, either  $u$  or  $v$  is adjacent to  $w$ . Then  $y$  cannot be adjacent to  $w$  as  $N(w) \cap \{u, v, x, y\}$  together with  $w$  induce  $C_4 \vee K_1$ . But,  $\langle u, v, w, x, y \rangle$  is a gem, a contradiction.  $\square$

By lemma 7.1, it now follows that each triangle in the root graph of a DHL graph is attached to the graph by sharing at the most one vertex. Let  $\mathcal{T}$  be the family of trees. Let  $\mathcal{T}_\Delta$  be the family of graphs obtained by attaching some triangles to some vertices in a tree  $T$ , for each  $T \in \mathcal{T}$ .

**Theorem 7.6.** *A graph  $G$  is a root graph of a  $C_4$ -free DHL graph if and only if  $G \in \mathcal{T}_\Delta$ .*

*Proof.* The proof is by induction on the number of edges in a  $T \in \mathcal{T}_\Delta$ . It can be verified that the root graphs of distance-hereditary graphs of size  $\leq 3$  are in  $\mathcal{T}_\Delta$  and hence the theorem is true for all  $m \leq 3$ .

Let  $T \in \mathcal{T}_\Delta$  has  $m$  edges and  $T$  is a root graph of a DHL graph. Let  $T'$  be a graph in  $\mathcal{T}_\Delta$  with  $E(T') = E(T) \cup \{e\}$ . Since  $T'$  must be connected, there can be two cases: either (i) the edge  $e$  is added as a pendent edge to  $T$  or (ii) the edge  $e$  is formed by joining two vertices in  $T$ .

Let  $l_e$  be the vertex in  $L(T')$  corresponding to the edge  $e$  in  $T'$ . In case(i), since  $e$  is a pendant edge in  $T'$ ,  $l_e$  is simplicial in  $L(T')$ . We can now show that  $L(T')$  is gem-free. If possible let a gem

is there in  $L(T')$ . Since  $L(T)$  is distance-hereditary and  $C_4$ -free, it is chordal. By Theorem 7.2  $L(T)$  is gem-free,  $l_e$  must be a vertex in the induced gem. But,  $N(l_e)$  is complete so that  $l_e$  is one of the degree two vertices in the gem. Now  $l_e$  is in a  $K_4 - e$ . By Lemma 7.1, one of the two triangles in the  $K_4 - e$  must be an anti-Gallai triangle. But the triangle containing  $l_e$  cannot be so, as  $e$  is a pendant edge in  $T'$ . But the other triangle has no vertex of degree 2 in the induced gem. This is a contradiction, by Lemma 7.1, to the assumption that  $L(T')$  contains a gem.

In case(ii), as  $T$  is connected, adding an edge  $e$  joining two vertices of  $T$  makes a cycle in  $T'$ . But  $T \in \mathcal{T}_\Delta$  is  $C_n$ -free,  $n \geq 4$ , and contains no  $K_4 - e$ . Hence  $e$  joins two pendant vertices of  $T$ , forming a triangle and has end vertices of degree two. Therefore in  $L(T')$ , the corresponding vertex  $l_e$  is in an anti-Gallai triangle and has degree two. It now follows that  $l_e$  is simplicial. If  $L(T')$  contains a gem,  $l_e$  must be one of the degree two vertices in the induced gem. But in this case the anti-Gallai triangle containing  $l_e$  do not satisfy Theorem 3.1 with the other vertex of degree two in the induced gem, which is again a contradiction.

In both the cases we have a one-vertex extension  $L(T')$  of a gem-free chordal graph  $L(T)$  and hence  $L(T')$  is a DHL graph.  $\square$

**Corollary 7.3.** *A graph  $L(G)$  is Ptolemaic if and only if  $G \in \mathcal{T}_\Delta$*

**Corollary 7.4.** *Let  $\mathcal{T}_\Delta^c$  be the family of graphs obtained by attaching some triangles to some vertices in a tree  $T$  and identifying each edge of  $T$  by an edge of at most one triangle, for each  $T \in \mathcal{T}$ . Then  $L(G)$  is a chordal graph if and only if  $G \in \mathcal{T}_\Delta^c$*

**Acknowledgment:** The authors thank the referee for some suggestions towards the improvement of the paper.

## References

- [1] A.V. Aho, J.E. Hopcroft and J.D. Ullman, *The design and analysis of computer algorithms*, Addison Wesley, (1976).
- [2] P. Anand, et al., On the hardness of recognizing triangular line graphs, *Discrete Math.* **312** (2012), 2627–2638.
- [3] L.S. Aparna, S.B. Rao and A. Vijayakumar, Gallai and anti-Gallai graphs of a graph, *Math. Bohem.* **132** (2007), 43–54.
- [4] R. Balakrishnan and K. Ranganathan, *A Textbook of Graph Theory*, second edition, Heidelberg: Springer (2012).
- [5] H.J. Bandelt and H.M. Mulder, Distance-hereditary graphs, *J. Combin. Theory Ser. B* **41** (1986), 182–208.
- [6] L.W. Beineke, On derived graphs and digraphs, *Beitrage zur Graphentheorie*, Leipzig (1968), 17–23.
- [7] F. Buckley and F. Harary, *Distance in Graphs*, Addison-Wesley, Redwood City, CA (1990).

- [8] A. D'Atri and M. Moscarini, Distance hereditary graphs, Steiner trees, and connected domination, *SIAM J. Comput.* **17** (1988), 521–538.
- [9] D.R. Fulkerson and O.A. Gross, Incidence matrices and interval graphs, *Pacific J. Math.* **15** (1965), 835–855.
- [10] T. Gallai, Transitiv Orientierbare Graphen, *Acta math. Acad. Sci. Hung.* **18** (1967), 25–66.
- [11] E. Howorka, A characterization of distance-hereditary graphs, *Quart. J. Math. Oxford. Ser.* **28** (1977), 417–420.
- [12] J. Krausz, Démonstration nouvelle d'une théorème de Whitney sur les réseaux, *Mat. Fiz. Lapok* **50** (1943), 75–85.
- [13] V.B. Le, Gallai graphs and anti-Gallai graphs, *Discrete Math.* **159** (1996), 179–189.
- [14] P.G.H. Lehot, An optimal algorithm to detect a line graph and output its root graph, *J. ACM* **21** (1974), 569–575.
- [15] L. Sun, Two classes of perfect graphs, *J. Combin. Theory Ser. B* **53** (1991), 273–292.
- [16] H. Whitney, Congruent graphs and the connectivity of graphs, *Amer. J. Math.* **54** (1932), 150–168.