


Berita Biologi

Jurnal Ilmu-ilmu Hayati



BERITA BIOLOGI

Vol. 16 No. 1 April 2017

Terakreditasi Berdasarkan Keputusan Kepala Lembaga Ilmu Pengetahuan Indonesia
No. 636/AU3/P2MI-LIPI/07/2015

Tim Redaksi (*Editorial Team*)

Andria Agusta (Pemimpin Redaksi, *Editor in Chief*)
Kusumadewi Sri Yulita (Redaksi Pelaksana, *Managing Editor*)
Gono Semiadi
Atit Kanti
Siti Sundari
Evi Triana
Kartika Dewi
Dwi Setyo Rini

Desain dan Layout (*Design and Layout*)

Muhamad Ruslan, Fahmi

Kesekretariatan (*Secretary*)

Nira Ariasari, Enok, Budiarjo

Alamat (*Address*)

Pusat Penelitian Biologi-LIPI
Kompleks Cibinong Science Center (CSC-LIPI)
Jalan Raya Jakarta-Bogor KM 46,
Cibinong 16911, Bogor-Indonesia
Telepon (021) 8765066 - 8765067
Faksimili (021) 8765059
Email: berita.biologi@mail.lipi.go.id
jurnalberitabiologi@yahoo.co.id
jurnalberitabiologi@gmail.com

Keterangan foto cover depan (*Notes of cover picture*): Bentuk koloni isolat bakteri Bt, BLSP-4, dan BLSP-3: (A) pada media pertumbuhan NA dan (B) pada pengamatan secara mikroskopis dengan perbesaran 100x (*Bacterial colony shapes of Bt, BLSP-4 and BLSP-3, respectively: (A) bacterial colony in growth medium NA (B) bacterial colony on 100 x microscopic magnification*), sesuai dengan halaman 15.



ISSN 0126-1754

636/AU3/P2MI-LIPI/07/2015

Volume 16 Nomor 1, April 2017

Berita Biologi

Jurnal Ilmu-ilmu Hayati

Berita Biologi	Vol. 16	No. 1	Hlm. 1 - 110	Bogor, April 2017	ISSN 0126-1754
----------------	---------	-------	--------------	-------------------	----------------

Pusat Penelitian Biologi - LIPI

Ucapan terima kasih kepada
Mitra Bebestari nomor ini
16(1) – April 2017

Dr. Heddy Julistiono
Ir. Suciati M.Si.
Dr. Nuril Hidayati
Drs. Haryono M.Si
Drs. Awit Suwito, M.Si
Dr. Rizkita Rachmi Esyanti
Prof. Dr. Amarila Malik, MSi., Apt.
Ir. I Gusti Bagus Adwita Arsa, MP.
Dr. Shanti Ratnakomala, M.Si
Dr. Fenny M. Dwivany
Dr. Ir. Barep Sutiyono, M.S.
Dr. I Made Sudiana, M.Sc.
Dr. Tri Muji Ermayanti
Dr. Ika Roostika Tambunan, SP. MSi.
Ucu Yanu Arbi M.Si.
Vani Nur Oktaviany Subagyo SP., Msi

CLONING, EXPRESSION, AND PARTIAL PURIFICATION OF PLANTARICIN W LOCUS PRODUCED BY *Lactobacillus plantarum* S34 [Kloning, Ekspresi, dan Purifikasi Parsial Lokus Plantarisin W Diproduksi oleh *Lactobacillus plantarum* S34]

Rifqiyah Nur Umami^{1✉}, Apon Zaenal Mustopa¹, Linda Sukmarini¹, Hasim Danuri²,
Andini Setyanti Putri², and Krisna Dwi Aria Wibowo³

^{✉1}Research Center for Biotechnology, Indonesian Institute of Sciences (LIPI), Cibinong, Bogor 16911

²Department of Biochemistry, Bogor Agricultural University (IPB), Darmaga, Bogor 16680

³Faculty of Biology, Gadjah Mada University (UGM), Yogyakarta 55281

email: rifqiyahnurumami@gmail.com

ABSTRAK

Lactobacillus plantarum S34 dilaporkan mempunyai aktivitas antibakteri yang terkait dengan produksi bakteriosin. Bagian dari gen yang menyandikan salah satu lokus bakteriosin yang diproduksi oleh *L. plantarum* S34, disebut dengan plantarisin W (*plnW*), diamplifikasi dari plasmid dan dikloning menggunakan sistem vektor pGEM[®]-T Easy ke dalam *Escherichia coli* DH5 α . Sekuens nukleotida *plnW* ($\pm$ 405 pb) diidentifikasi sebagai protein integral membran. Lebih lanjut, *plnW* diekspresikan secara heterologus sebagai fusi protein dengan His(6)-tag tioredoksin menggunakan vektor ekspresi pET-32a(+) ke dalam *E. coli* BL21 (DE3) pLysS. Protein fusi rekombinan *plnW* terdapat dalam sitoplasma sel, tetapi selain fraksi terlarut terdapat juga fraksi tidak terlarut berupa badan inklusi. Purifikasi parsial dilakukan menggunakan kromatografi afinitas ligan Co²⁺ untuk fraksi terlarut dan metode elektroelusi gel poliakrilamid untuk fraksi tidak terlarut. Massa molekul berukuran kurang lebih 33 kDa terdeteksi berdasarkan pemisahan SDS-PAGE dan dikonfirmasi dengan Western blot sebagai protein fusi rekombinan *plnW*. Protein yang sudah terpurifikasi bermanfaat untuk mengetahui kaitan antara struktur dan fungsi bakteriosin.

Kata kunci: bakteriosin, badan inklusi, *Lactobacillus plantarum* S34, plantarisin W.

ABSTRACT

Lactobacillus plantarum S34 was reported to have antibacterial activities which associated to bacteriocins production. Part of the gene encoding one of the bacteriocin loci that produced by *L. plantarum* S34, termed plantaricin W (*plnW*), was amplified from plasmid and cloned into *Escherichia coli* DH5 α using pGEM[®]-T Easy vector system. *PlnW* nucleotide sequence ($\pm$ 405 bp) was identified as a putative integral membrane protein. Moreover, *plnW* was heterologously expressed as a fusion protein with His(6)-tagged thioredoxin using pET-32a(+) expression vector into *E. coli* BL21 (DE3) pLysS. Recombinant *plnW* fusion protein was accumulated in the cell cytoplasm, however, along with soluble fractions, insoluble aggregates identified as inclusion bodies were also exhibited. Partial purifications were conducted for the soluble and insoluble fractions using ligand Co²⁺ affinity chromatography and polyacrylamide gel electroelution method, respectively. Molecular mass of approximately 33 kDa was detected based on SDS-PAGE separation and confirmed by Western blot as a recombinant *plnW* fusion protein. The purified protein will be useful for deciphering the bacteriocin structural and functional relationships.

Key words: bacteriocin, inclusion body, *Lactobacillus plantarum* S34, plantaricin W.

INTRODUCTION

Lactic acid bacteria (LAB), group of bacteria acquired generally recognized as safe (GRAS) status, have been known for their valuable role as culture starters and natural preservatives in fermented products since they can inhibit the growth of spoilage microorganisms. One of the antimicrobial agents produced by LAB is bacteriocin, the ribosomally synthesized and extracellularly released peptide or protein which has either narrow or broad spectrum antibacterial effects (Gálvez *et al.*, 2007; Karpiński and Szkaradkiewicz, 2013; Yang *et al.*, 2014). Studies on bacteriocins have been focused on identification, characterization of gene clusters, expression, production, mode of action, biosynthesis, bioactivities, and elucidation of their functional structures (Field *et al.*, 2007; Todorov, 2009).

Lactobacillus plantarum, an adaptable LAB species that can be found in different kinds of carbohydrate rich niches, is able to produce a number of bacteriocins that varies within strains. *L. plantarum* C11 has been reported as one of the earliest bacteriocin producer strains which already genetically characterized in terms of biosynthesis and gene regulator determinants (Diep *et al.*, 1996). In addition, other *L. plantarum* strains including, NC8 (Maldonado *et al.*, 2003), WCFS1 (Kleerebezem *et al.*, 2003), J51 (Navarro *et al.*, 2008), J23 (Rojo-Bezares *et al.*, 2008), V90 (Diep *et al.*, 2009), JDM1 (Zhang *et al.*, 2009), and ST-III (Wang *et al.*, 2011), have been found to retain similar bacteriocin loci in their genome. These loci, termed plantaricins (*pln*), can be either plasmid or chromosomally encoded. Recently, plantaricins have

been received great attention due to their diversities and potential development of new antimicrobial additives (Diep *et al.*, 2009; Sáenz *et al.*, 2009).

There are about 22–25 genes within *pln* gene clusters of *L. plantarum* which being arranged into 5–6 operons under the regulation of multiple promoters and responsible for the biosynthesis of various bacteriocins. Two-peptide bacteriocins derived from plantaricin loci have been reported such as plantaricin A (encoded by two peptides α and β), plantaricin S (encoded by *plsA* and *plsB*), plantaricin W (encoded by *phwa* and *phwB*), plantaricin NC8 (encoded by *plNC8 α* and *plNC8 β*), and plantaricin J51 (encoded by *orf3* and *orf4*) (Issen-Meyer *et al.*, 1993; Stephens *et al.*, 1998; Holo *et al.*, 2001; Diep *et al.*, 2009).

There are some important operons including *plnABCD*, *plnMNOP* and *plnGHSTUVWXY*. *PlnTUVWXY* codes putative integral membrane proteins acquiring protease CAAX motif which belong to Abi family and associated to bacteriocin self-immunity. These putative transmembrane segments are essential for the integration of immunity proteins into the membrane of the bacteriocin producer. Despite the extensive studies of the operon, the exact role of *plnTUVWXY* remains to be elucidated (Diep *et al.*, 2009; Sáenz *et al.*, 2009; Tai *et al.*, 2015).

Lactobacillus plantarum S34 is one of LAB isolates originated from traditional fermented beef product named *bekasam*, commonly found at Way Kanan, Lampung, Indonesia. Recently, 16S rRNA sequencing analysis shows that *L. plantarum* S34 has 99% nucleotide homology with *L. plantarum* WCFS1 [GenBank Accession No. AL935263]. *L. plantarum* S34 is selected for further molecular genetic study since it shows potential antibacterial activities against representative pathogens such as *Escherichia coli*, *Salmonella typhi*, and *Bacillus subtilis*, which associated to bacteriocins production (Mustopa and Fatimah, 2014). This study aims to gain high yield protein from one of plantaricin loci of *L. plantarum* S34 designated plantaricin W (*plnW*), as a production model of bacteriocin integral membrane protein through heterologous expression in *Escherichia coli*.

MATERIALS AND METHODS

Plasmid DNA Isolation

Plasmid DNA of *L. plantarum* S34 were isolated by alkaline lysis method based on Sambrook *et al.*, 1989; and O'Sullivan and Klaenhammer, 1993, with several modifications. Pellets were collected from *L. plantarum* S34 overnight cultures (MRS medium) and resuspended in alkaline lysis I solution (25 mM Tris-HCl pH 8.0, 50 mM glucose, and 10 mM EDTA) containing 10 mg mL⁻¹ lysozyme and incubated at 37°C for 2 hours. Freshly prepared alkaline lysis II solution (0.2 N NaOH and 1% SDS) were mixed and incubated at RT for 5 minutes. Next, ice-cold alkaline lysis III solution (5 M potassium acetate, glacial acetic acid, and H₂O) were mixed and incubated on ice for 10 minutes followed by centrifugation at 13,000 rpm at 4°C for 10 minutes. The upper phase were mixed (1:1) with PCI (phenol:chloroform:isoamyl alcohol = 25:24:1) followed by centrifugation at 13,000 rpm at 4°C for 10 minutes. The upper phase were precipitated with isopropanol (1:1) by incubation at -20°C for 2 hours and followed by centrifugation at 13,000 rpm at 4°C for 10 minutes. The pellets were rinsed with 70% ethanol, air-dried and resuspended in H₂O containing 0.1 mg mL⁻¹ RNase. Plasmid DNA isolation from *E. coli* DH5 α , *E. coli* TOP10 and *E. coli* BL21 (DE3) *plysS* were performed using QIAGEN plasmid MINI and MIDI kits (Qiagen, Germany), according to manufacture's instructions.

Cloning

PlnW gene was amplified from native plasmid of *L. plantarum* S34 by PCR using Platinum *taq* DNA polymerase (Invitrogen, USA) and primers previously described by Sáenz *et al.*, 2009 (forward primer *plnW*MY42-F: 5'-GATCAGCCACGATAC-CAAC-3'; reverse primer *plnW*MY42-R: 5'-CTAAAGAAAAAGCCCCCTGAAAC-3'). The PCR cycles were performed as follow: initial denaturation at 94°C for 3 minutes, 35 cycles of amplification consisting: denaturation at 94°C for 1 minute, annealing at 58.5°C for 1 minute, and extension at 72°C for 90 seconds. The final extension was performed at 72°C for 6 minutes. PCR product ($\pm$ 750 bp) was confirmed by 1.0% agarose gel electrophoresis. PCR purifications were done using

Qiaquick Gel Extraction kit (Qiagen, Germany) according to manufacture's instructions. Cloning of *plnW* gene into *E. coli* DH5 α (Novagen, USA) was performed using pGEM[®]-T Easy vector system (Promega, USA) according to manufacture's instructions. The molecular size of DNA target ($\pm$ 750 bp) was confirmed by restriction enzymes (*Nco* I and *Sal* I). Sequence confirmation was performed using T7 promoter and SP6 terminator primers and analyzed using BioEdit software.

Based on the obtained sequence, new primer pairs were designed containing *Bam*H I and *Hind* III restriction sites (forward primer plnWS34BHinF: 5'-GACTGGATCCATGTTACAGAAGAATTTACG-3'; reverse primer plnWS34BHinR: 5'-GATCAAGCTTTCAAATGACGGCATCGAGTG-3'). Plasmid derived from pGEM-plnW were used as template for the PCR cycles as mentioned above with annealing temperature at 55°C for 1 minute. PCR product ($\pm$ 405 bp) was confirmed and purified.

The target gene was ligated into pET32a (+) (Novagen, USA) expression vector, in-frame with His(6)-tagged thioredoxin using restriction enzymes and T4 ligase (Takara Bio Inc, Japan). The assembled vector was firstly sub-cloned into *E. coli* TOP 10 (Novagen, USA) and then transformed into *E. coli* BL21 (DE3) pLysS (Novagen, USA) as expression host using standard method described by Sambrook *et al.*, 1989. LB agar medium (1.5% w/v) containing appropriate antibiotics (100 μ g mL⁻¹ ampicillin for *E. coli* TOP 10; 100 μ g mL⁻¹ ampicillin and 25 μ g mL⁻¹ chloramphenicol for *E. coli* BL21 (DE3) pLysS) were used as selection markers. The bacterial transformants were selected and confirmed by PCR and nucleotide sequencing. Furthermore, amino acid alignments were compared with the reference strain sequences (AL935263.2, DQ323671.2, CP002222.1, FJ809773.1, CP001617.1, X94434.2, and DQ340868.2) using Basic Local Alignment Search Tool (BLAST) program (<http://www.ncbi.nlm.nih.gov/BLAST>).

Expression

The expression of *plnW* protein recombinant was performed according to Putri *et al.*, 2014, by inoculating 10% (v/v) of an overnight culture of *E. coli* BL21(DE3) pLysS harboring assembled plasmid

into 200 mL of LB broth containing 100 μ g mL⁻¹ ampicillin at 37°C with agitation at 150 *rpm*. As inducer, 0.5 mM of isopropyl- β -D-thiogalactopyranoside (IPTG - Thermo Scientific, USA) were added when OD₆₀₀ $\approx$ 0.6, followed by further incubation at 22°C with agitation at 150 *rpm* for 5 hours. The cells were harvested and after three times of freezing and thawing, pellets were resuspended in lysis buffer (10 mM Tris-HCl pH 8.5, 100 mM NaCl, and 0.25% (v/v) Tween-20) and subjected into on-ice sonication (5 cycles of 15 seconds pulse on and 60 seconds pulse off) for cell disruption. The cell lysate were centrifuged at 13,000 *rpm* at 4°C for 30 minutes. Both of the obtained supernatants (soluble fractions) and pellets (insoluble fractions) were subjected for further purification methods.

Partial Purification of Soluble Fraction

The soluble fractions were partially purified using immobilized metal affinity chromatography (IMAC) resin charged with cobalt (Co²⁺) (TALON, Clontech, Takara Bio) as previously described by Utama *et al.*, 2000. Briefly, soluble fractions were applied on lysis buffer-calibrated resin by gently binding it for 3 hours at 4°C. The resin-bound protein was collected by brief centrifugation followed by two times washing with lysis buffer containing 10 mM imidazole. The recombinant protein was purified and eluted with lysis buffer containing 400 mM imidazole. The purified protein was denaturated and analyzed using 12% SDS PAGE (sodium dodecyl sulfate polyacrylamide gel electrophoresis) as described by Laemmli, 1970, stained with staining solution (0.5% (w/v) Coomassie Brilliant Blue G (CBB-G, Sigma, USA), 50% (v/v) methanol, and 10% acetic acid), and destained with destaining solution (50% (v/v) methanol and 10% acetic acid). Precision Plus Protein Standards (Bio Rad, USA) was used as molecular weight marker.

Partial Purification of Insoluble Fraction

The remaining insoluble fractions were washed three times with washing buffer (50 mM Tris-HCl pH 8.0, 100 mM NaCl, 2 M Urea, and 1% Triton X-100) according to Liu *et al.*, 2011, before solubilized

with solubilizing buffer (100 mM Tris-HCl pH 12.5 and 2 M Urea) by an overnight constant mixing at 4°C, according to Patra *et al.*, 2000. The denaturing agents were removed from solubilized protein by step-wise dialysis system, in which Urea concentration was gradually reduced, according to Tsumoto *et al.*, 1998, with several modifications. Firstly, the solubilized protein was dialyzed against denaturing removal buffer 1a (50 mM Tris-HCl pH 8.0, 0.2 mM EDTA, and 1 M Urea) followed by denaturing removal buffer 1b (50 mM Tris-HCl pH 8.0, 0.2 mM EDTA, and 0.5 M Urea). Secondly, protein refolding was performed by dialysis against refolding buffer 2a (50 mM Tris-HCl pH 8.0, 0.2 mM EDTA, 0.4 M Urea, and 0.25 mM 2-mercaptoethanol) followed by stabilizer buffer 2b (50 mM Tris-HCl pH 8.0, 0.2 mM EDTA, 0.4 M Urea, 0.25 mM 2-mercaptoethanol, and 1% (v/v) glycerol). Each dialysis was performed by gentle agitation for 24 hours at 4°C using cellulose membrane dialysis tubing (12,400 MWCO, SIGMA, Germany) pre-heated with 1 mM EDTA and H₂O for 10 minutes, subsequently.

The refolded protein was pooled and directly subjected into preparative SDS-PAGE for further partial purification using electroelution method according to Mohammadian *et al.*, 2010 with some modifications. Briefly, after being stained and destained with the same method above, the target band was precisely excised and sliced into small pieces using clean scalpel. Gel fragments were carefully put into membrane dialysis tubing containing electroelution buffer (25 mM Tris pH 8.3, 0.192 M glycine, and 0.1% SDS). The membrane tube was then placed into electrophoresis apparatus filled with electroelution buffer and the electroelution was performed at 60 V at 4°C until CBB-G was being removed and the gel become colorless.

Protein precipitation was done by adding four times volume of cold acetone to the electroeluted protein and incubated on ice for 30 minutes. The precipitated protein was separated by centrifugation at 10,000 *rpm* at 4°C for 10 minutes, allowed to dry and dissolved in 20 µL of 8 M Urea in dilution buffer (50 mM Tris pH 7.9, 20% glycerol, 0.1 mM EDTA, 1 mM DTT, 0.15 M NaCl, and 0.1% SDS), followed

by incubation at room temperature for 20 minutes. The solution was diluted 10 times using dilution buffer and permitted to renaturation for 1 hour at room temperature. The purified protein was then analyzed by 12% SDS-PAGE for molecular weight confirmation.

Western Blot Analysis

Western blot analysis was performed using Anti-His(C-term)-HRP Antibody (Invitrogen, USA), according to the manufacture's instructions. Briefly, the obtained solubilized lysate were separated using 12% SDS-PAGE without staining and then electrophoretically transferred (30 V, 40 mA; overnight incubation at 4°C) to nitrocellulose membrane (blotting) in transfer buffer (25 mM Tris pH 8.3, 192 mM glycine, and 20% (v/v) methanol). The membrane was blocked by gentle agitation in 10 mL blocking buffer (PBS/T: 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄, and 0.05% (v/v) Tween-20; and 5% (w/v) nonfat, dry milk) for 1 hour at room temperature. The membrane was then washed two times with PBS/T for 5 minutes each, and transferred to 10 mL blocking buffer containing 3 µL Anti-His(C-term)-HRP Antibody followed by overnight incubation with gentle agitation at 4°C. After the membrane was washed two times with PBS/T for 5 minutes each, the target band was detected by Novex HRP Chromogenic Substrate (Invitrogen, USA) consist of TMB (3,3',5,5'-tetramethylbenzidine) which forms a blue precipitate upon reaction with HRP (horseradish peroxidase).

RESULT

Cloning and Expression

The PCR target product (± 750 bp) was successfully cloned into *E. coli* DH5α using pGEM®-T Easy vector system and confirmed by restriction enzymes (*Nco* I and *Sal* I) (Figure 1). Sequence analysis showed that there was additional region (partial *plnV* locus) within the amplified product (data not shown) possibly due to nonspecific amplification in PCR. New primer pairs with additional restriction sites (*Bam*H I and *Hind* III) were designed in order to amplify specific region of *plnW* locus. The desired PCR product (± 405 bp) harboring *plnW* region was successfully amplified




Figure 1. DNA target (± 750 bp) was confirmed using restriction enzymes (*Nco* I and *Sal* I) and visualized in 1.0% gel electrophoresis. (DNA target (± 750 bp) dikonfirmasi menggunakan enzim restriksi (*Nco* I dan *Sal* I) dan divisualisasi menggunakan gel elektroforesis 1%).




Figure 2. Sequence analysis of recombinant *plnW* gene in fusion with His(6)-tagged thioredoxin.
(Analisis sekuens fusi rekombinan gen *plnW* dengan His(6)-tagged thioredoxin)

and transformed into *E. coli* TOP 10 and *E. coli* BL21 (DE3) pLysS using pET32a(+) as expression vector and confirmed by nucleotide sequencing (Figure 2).

The amino acid multiple alignments showed that the obtained *plnW* amino acid sequence had similar identity compared to the previously published *plnW* amino acid sequences (strains WCFS1, J23, STIII, V90, JDM1, C11, and J51; <http://www.ncbi.nlm.nih.gov>) (Figure 3). However, only 135 out of 228 amino acid from the whole part of *plnW* locus could be confirmed in this work. The PCR amplification using primer pairs targeted the

whole part of *plnW* was unsuccessful (data not shown). Nonetheless, partial *plnW* locus could also be seen on the other *L. plantarum* strains such as C11 (219 amino acid) and J51 (150 amino acid).

Partial Purification of Soluble and Insoluble Fractions

The recombinant fusion protein derived from soluble fractions which partially purified using IMAC resin was visualized by 12% SDS-PAGE stained by CBB-G (Figure 4A). The fusion protein derived from insoluble fractions was partially purified and confirmed by CBB-G (Figure 4B). The

partially purified fusion protein derived from insoluble fractions exhibited higher purity and

concentration compared to the one derived from soluble fractions.



Figure 3. Multiple alignments of *plnW* amino acid isolated in this study compared to the previously published *plnW* locus derived from several reference strains of *L. plantarum*. (*Multiple Alignment asam amino plnW yang diisolasi pada penelitian ini dengan locus plnW dari beberapa strain L. plantarum yang sudah dipublikasi sebelumnya*).




Figure 4. SDS-PAGE analysis of recombinant *plnW* fusion protein; (A). Partial purification of soluble fractions using IMAC: M. Marker; 1. Cell lysate; 2. Protein target (± 33 kDa) eluted with 400 mM imidazole; (B). Partial purification of insoluble fractions using electroelution: M. Marker; 1. Solubilized protein; 2. Dialyzed protein; 3. Protein target (± 33 kDa) partially purified with electroelution, precipitated, and diluted with dilution buffer. (*Analisis SDS-PAGE fusi rekombinan protein plnW: (A). Purifikasi parsial fraksi terlarut menggunakan IMAC: M. Marker; 1. Cell lysate; 2. Protein target (± 33 kDa) dielusi dengan 400 mM imidazole; (B). Purifikasi parsial fraksi tidak terlarut menggunakan elektroelusi: M. Marker; 1. Protein tersolubilisasi; 2. Protein terdialisis; 3. Protein target (± 33 kDa) yang dipurifikasi parsial dengan elektroelusi, dipresipitasi dan didilusi dengan bufer dilusi*).




Figure 5. The target protein (± 33 kDa), which indicated by white arrow, was successfully visualized by Novex HRP Chromogenic Substrate (Invitrogen, USA). [Protein target (± 33 kDa), yang ditunjukkan dengan tanda panah putih, berhasil divisualisasi dengan Novex HRP Chromogenic Substrate (Invitrogen, USA)]

Western Blot Analysis

Anti-His(C-term)-HRP Antibody allowed the detection of the recombinant fusion protein (His (6) tag-*plnW*) containing a C-terminal polyhistidine tag with a free carboxyl group within the solubilized protein. The result showed that the desired fusion protein (± 33 kDa) was successfully detected (Figure 5).

DISCUSSION

Bacteriocins production is one of the alternative solutions toward the increasing number of antibiotic-resistant pathogens and the side effect of chemical preservatives. Nevertheless, bacteriocins quorum sensing-dependent production in wild environments during late lag growth phase is often result in low concentration (Diep *et al.*, 2009). One approach that can be done to address this issue is using recombinant technology to produce the target bacteriocin abundantly and more stable. The excessive production of recombinant bacteriocin also beneficial for further elucidation of genetic structural and functional relationships, thus, will lead to better understanding of their mechanism of action at molecular level (Meyer *et al.*, 2010).

However, overproduction of bacteriocins, including plantaricins, may also cause lethal effect to the host cell due to their toxicity. Thus, currently there is no ideal system for the expression of native plantaricin. It is note worthy that the expression host plays important role on the quality and quantity of

the produced recombinant plantaricin. *E. coli* is commonly used as expression host since it is easy to handle. Mustopa *et al.*, 2016, reported that plantaricin E and F derived from *L. plantarum* S34 have been successfully cloned and expressed into *E. coli*.

Therefore, this study was heterologous production system of recombinant plantaricin W, one of the bacteriocin loci encoded by plasmid-borne genes derived from *L. plantarum* S34, in *E. coli*. Plantaricin W (*plnW*) has been identified as a putative integral membrane protein located downstream of the structural genes. This locus generally involved in a wide variety of cellular functions, one of them is self-immunity system, which protect the producer bacteria from being killed by its own bacteriocin.

This study suggests a serial method of isolation, cloning, expression, and partial purification of recombinant *plnW*. Although a truncated region has been identified, the sequence analysis shows a high identity compare to the reference strains aligned (Figure 3). These genomic discrepancies may be due to the alteration of gene expression within *L. plantarum* strains, which derived from distinctive niches, as the result of environmental adaptation.

It is important to consider some drawbacks such as potential toxicity of *plnW* to the host, lack of secretion system for efficient release of *plnW*, and limited ability to facilitate extensive disulfide bond formation due to overproducing of *plnW* during

heterologous expression in *E. coli* BL21 (DE3) pLysS. Therefore, the fusion system (His(6)-tagged thioredoxin within pET-32a(+) expression vector) is used to prevent inclusion body formation and to generate standard purification method. His-tags enable purification method based on metal ions affinity which allows single step purification from crude extract cells, while thioredoxin is a hydrophilic tag which can increase the protein solubility.

The heterologous expression method of *plnW* in *E. coli* was optimized as previously reported by Putri *et al.*, 2014. In order to increase the solubility, *plnW* was expressed at lower temperature and induced by low concentration of IPTG. Our result showed that the purification using affinity chromatography was insufficient to obtain high yield of recombinant *plnW*. The fusion protein could not effectively bound to the resin, and most of it was remained in cell lysate, thus resulted in low purity.

Unfortunately, the formation of inclusion body still become bottleneck during the expression of recombinant *plnW*. We used several strategies to recover the active *plnW*, including solubilization of inclusion body, refolding of solubilized protein and partial purification of refolded protein. Washing step using low concentration of Triton X-100 helped in removing several impurities. Mild solubilization using low concentration of Urea at alkaline condition could solubilize the inclusion body aggregates while preserving the native-like protein structure. We also described step-wise dialysis refolding method, which controls the folding pathway by gradually adjusting the concentration of the denaturant to induce sequential folding.

Partial separation of denatured protein was best achieved by preparative SDS-PAGE followed by electroelution of the protein which results in high-throughput recovery of the proteins from inclusion body. However, additional purification step such as thioredoxin removal using enterokinase enzyme is still needed.

CONCLUSION

This study suggests heterologous expression and partial purification system of plantaricin W derived from *L. plantarum* S34 using *E. coli* as expression host, which allows convenient and

efficient way of recovering properly folded recombinant protein from inclusion body aggregates. The characterization of this locus will be useful for the exploration of *L. plantarum* S34 antibacterial properties. More information on the genetic structures of plantaricin loci of indigenous lactic acid bacteria from Indonesia is needed. Further research on some important gene including for the proposing of new species or genus is mandatory.

ACKNOWLEDGMENTS

This study was financially supported by Kompetitif LIPI program 2012-2014. We thank to Muhamad Ridwan, Dwianty Putri Meitasari, Ike Rahmawati, and Kusdianawati for technical assistance.

REFERENCES

- Diep DB, LS Håvarstein and IF Nes. 1996. Characterization of The Locus Responsible for The Bacteriocin Production in *Lactobacillus plantarum* C11. *Journal of Bacteriology* **178**(15), 4472-4483.
- Diep DB, D Straume, M Kjos, C Torres and IF Nes. 2009. An Overview of The Mosaic Bacteriocin *pln* Loci from *Lactobacillus plantarum*. *Peptides* **30**, 1562-1574.
- Field D, PD Cotter, C Hill and RP Ross. 2007. Bacteriocin Biosynthesis, Structure, and Function. In: *Bacteriocins Current Research and Applications*, 5-40. MA Riley and O Gillor (Eds), Horizon Bioscience. Norfolk, UK.
- Gálvez A, H Abriouel, RL López and N Ben Omar. 2007. Bacteriocin-Based Strategies for Food Biopreservation. *International Journal of Food Microbiology* **120**, 51-70.
- Holo H, Z Jeknic, M Daeschel, S Stevanovic and IF Nes. 2001. Plantaricin W from *Lactobacillus plantarum* Belongs to A New Family of Two-Peptide Lantibiotics. *Microbiology* **147**, 643-651.
- Issen-Meyer J, AG Larsen, K Sletten, M Daeschel and IF Nes. 1993. Purification and Characterization of Plantaricin A, a *Lactobacillus plantarum* Bacteriocin Whose Activity Depends on The Action of Two Peptides. *Journal of General Microbiology* **139**, 1973-1978.
- Karpiński TM and AK Szkaradkiewicz. 2013. Characteristic of Bacteriocines and Their Application. *Polish Journal of Microbiology* **62**(3), 223-235.
- Kleerebezem M, J Boekhorst, R Van Kranenburg, D Molenaar, OP Kuipers, R Leer, R Tarchini, SA Peters, HM Sandbrink, MWEJ Fiers, W Stiekema, RMK Lankhorst, PA Bron, SM Hoffer, MNN Groot, R Kerkhoven, M De Vries, B Ursing, WM De Vos and RJ Siezen. 2003. Complete Genome Sequence of *Lactobacillus plantarum* WCFS1. *Proceeding of The National Academy of Sciences USA* **100**, 1990-1995.
- Laemmli UK. 1970. Cleavage of Structural Proteins During The Assembly of The Head of Bacteriophage T4. *Nature* **227**(5259), 680-685.
- Liu S, Y Han and Z Zhou. 2011. Fusion Expression of *pedA* Gene to Obtain Biologically Active Pediocin PA-1 in *Escherichia coli*. *Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology)* **12**(1), 65-71.
- Maldonado A, JL Ruiz-Barba and R Jiménez-Díaz. 2003. Purification and Genetic Characterization of Plantaricin NC8, a Novel Coculture-Inducible Two-Peptide Bacteriocin from *Lactobacillus plantarum* NC8. *Applied*

- and *Environmental Microbiology* **69**(1), 383-389.
- Meyer JN, C Oppedgaard, P Rogne, HS Haugen and PE Kristiansen. 2010. Structure and Mode-of-Action of the Two-Peptide (Class-IIb) Bacteriocins. *Probiotics and Antimicrobials Proteins* **2**, 52-60.
- Mohammadian T, M Doosti, M Paknejad, F Siavoshi and S Massarrat. 2010. Preparative SDS-PAGE Electroelution for Rapid Purification of Alkyl Hydroperoxide Reductase from *Helicobacter pylori*. *Iranian Journal of Public Health* **39**(1), 85-91.
- Mustopa AZ and Fatimah. 2014. Diversity of Lactic Acid Bacteria Isolated from Indonesian Traditional Fermented Foods. *Microbiology Indonesia* **8**(2), 48-57.
- Mustopa AZ, Kustianawati, Fatimah, RN Umami, BR Budiarto and H Danuri. 2016. Cloning and Expression of Plantaricin E and F Genes of *Lactobacillus plantarum* S34 Isolated From Indonesia Traditional-Fermented Meat (*Bekasam*). *International Food Research Journal* **23**(2), 762-769.
- Navarro L, B Rojo-Bezares, Y Sáenz, L Díez, M Zarazaga, F Ruiz-Larrea and C Torres. 2008. Comparative Study of The *pln* Locus of The Quorum-Sensing Regulated Bacteriocin-Producing *L. plantarum* J51 Strain. *International Journal of Food Microbiology* **128**, 390-394.
- O'Sullivan DJ and TR Klaenhammer. 1993. Rapid Mini-Prep Isolation of High-Quality Plasmid DNA from *Lactococcus* and *Lactobacillus* spp. *Applied and Environmental Microbiology* **59**(8), 2730-2733.
- Patra AK, R Mukhopadhyay, R Mukhija, A Krishnan, LC Garg and AK Panda. 2000. Optimization of Inclusion Body Solubilization and Renaturation of Recombinant Human Growth Hormone from *Escherichia coli*. *Protein Expression and Purification* **18**(2), 182-192.
- Putri AS, RN Umami, AZ Mustopa and H Danuri. 2014. Heterologous Expression of Recombinant Plantaricin WS34 in *Escherichia coli*. *Proceeding The ASEAN Conference on Science and Technology 2014 (ASEAN CoSaT 2014)*, IPB International Convention Center, Bogor, West Java, Indonesia 18th -19th August 2014. Estiko Rijanto (Ed.), 65-72. LIPI Press, Jakarta.
- Rojo-Bezares B, Y Sáenz, L Navarro, R Jiménez-Díaz, M Zarazaga, F Ruiz-Larrea and C Torres. 2008. Characterization of A New Organization of The Plantaricin Locus In The Inducible Bacteriocin-Producing *Lactobacillus plantarum* J23 of Grape Must Origin. *Archives of Microbiology* **189**, 491-499.
- Sáenz Y, B Rojo-Bezares, L Navarro, L Díez, S Somalo, M Zarazaga, F Ruiz-Larrea and C Torres. 2009. Genetic Diversity of The *pln* Locus Among Oenological *Lactobacillus plantarum* Strains. *International Journal of Food Microbiology* **134**, 176-183.
- Sambrook J, EF Fritsch and T Maniatis. 1989. Molecular Cloning A Laboratory Manual. 2nd edn. Cold Spring Harbor Laboratory Press, New York, USA.
- Stephens SK, B Floriano, DP Cathcart, SA Bayley, VF Witt, R Jiménez-Díaz, PJ Warner and JL Ruiz-Barba. 1998. Molecular Analysis of The Locus Responsible For Production of Plantaricin S, A Two-Peptide Bacteriocin Produced by *Lactobacillus plantarum* LPCO10. *Applied and Environmental Microbiology* **64**(5), 1871-1877.
- Tai HF, HL Foo, RA Rahim, TC Loh, MP Abdullah and K Yoshinobu. 2015. Molecular Characterisation of New Organisation of *plnEF* and *plw* Loci of Bacteriocin Genes Harbour Concomitantly in *Lactobacillus plantarum* I24UL4. *Microbial Cell Factories* **14**(89), 1-13.
- Todorov SD. 2009. Bacteriocins From *Lactobacillus plantarum* – Production, Genetic Organization and Mode of Action. *Brazilian Journal of Microbiology* **40**, 209-221.
- Tsumoto K, K Shinoki, H Kondo, M Uchikawa, T Juji and I Kumagai. 1998. Highly Efficient Recovery of Functional Single-Chain Fv Fragments From Inclusion Bodies Overexpressed in *Escherichia coli* by Controlled Introduction of Oxidizing Reagent - Application to A Human Single-Chain Fv Fragment. *Journal of Immunological Methods* **219**, 119-129.
- Utama A, H Shimizu, S Morikawa, F Hasebe, K Morita, A Igarashi, M Hatsu, K Takamizawa and T Miyamura. 2000. Identification and Characterization of The RNA Helicase Activity of Japanese Encephalitis Virus NS3 Protein. *FEBS Letters* **465**(1), 74-78.
- Wang Y, C Chen, L Ai, F Zhou, Z Zhou, L Wang, H Zhang, W Chen and B Guo. 2011. Complete Genome Sequence of The Probiotic *Lactobacillus plantarum* ST -III. *Journal of Bacteriology* **193**(1), 313-314.
- Yang SC, CH Lin, CT Sung and JY Fang. 2014. Antibacterial Activities of Bacteriocins: Application In Foods and Pharmaceuticals. *Frontiers in Microbiology* **5** (241), 1-10.
- Zhang ZY, C Liu, YZ Zhu, Y Zhong, YQ Zhu, HJ Zheng, GP Zhao, SY Wang and XK Guo. 2009. Complete Genome Sequence of *Lactobacillus plantarum* JDM1. *Journal of Bacteriology* **191**(15), 5020-5021.

Pedoman Penulisan Naskah Berita Biologi

Berita Biologi adalah jurnal yang menerbitkan artikel kemajuan penelitian di bidang biologi dan ilmu-ilmu terkait di Indonesia. Berita Biologi memuat karya tulis ilmiah asli berupa makalah hasil penelitian, komunikasi pendek dan tinjauan kembali yang belum pernah diterbitkan atau tidak sedang dikirim ke media lain. Masalah yang diliput, diharuskan menampilkan aspek atau informasi baru.

Tipe naskah

1. **Makalah lengkap hasil penelitian (*original paper*)**
Naskah merupakan hasil penelitian sendiri yang mengangkat topik yang *up-to-date*. Tidak lebih dari 15 halaman termasuk tabel dan gambar. Pencantuman lampiran seperlunya, namun redaksi berhak mengurangi atau meniadakan lampiran.
2. **Komunikasi pendek (*short communication*)**
Komunikasi pendek merupakan makalah hasil penelitian yang ingin dipublikasikan secara cepat karena hasil temuan yang menarik, spesifik dan baru, agar dapat segera diketahui oleh umum. Artikel yang ditulis tidak lebih dari 10 halaman. Hasil dan pembahasan boleh digabung.
3. **Tinjauan kembali (*review*)**
Tinjauan kembali merupakan rangkuman tinjauan ilmiah yang sistematis-kritis secara ringkas namun mendalam terhadap topik penelitian tertentu. Hal yang ditinjau meliputi segala sesuatu yang relevan terhadap topik tinjauan yang memberikan gambaran '*state of the art*', meliputi temuan awal, kemajuan hingga issue terkini, termasuk perdebatan dan kesenjangan yang ada dalam topik yang dibahas. Tinjauan ulang ini harus merangkum minimal 30 artikel.

Struktur naskah

1. **Bahasa**
Bahasa yang digunakan adalah bahasa Indonesia atau Inggris yang baik dan benar.
2. **Judul**
Judul harus singkat, jelas dan mencerminkan isi naskah diikuti oleh nama dan alamat surat menyurat penulis. Nama penulis untuk korespondensi diberi tanda amplop cetak atas (*superscript*).
3. **Abstrak**
Abstrak dibuat dalam dua bahasa, bahasa Indonesia dan Inggris. Abstrak memuat secara singkat tentang latar belakang, tujuan, metode, hasil yang signifikan, kesimpulan dan implikasi hasil penelitian. Abstrak berisi maksimum 200 kata, spasi tunggal. Di bawah abstrak dicantumkan kata kunci yang terdiri atas maksimum enam kata, dimana kata pertama adalah yang terpenting. Abstrak dalam bahasa Inggris merupakan terjemahan dari bahasa Indonesia. Editor berhak untuk mengedit abstrak demi alasan kejelasan isi abstrak.
4. **Pendahuluan**
Pendahuluan berisi latar belakang, permasalahan dan tujuan penelitian. Sebutkan juga studi terdahulu yang pernah dilakukan.
5. **Bahan dan cara kerja**
Pada bagian ini boleh dibuat sub-judul yang sesuai dengan tahapan penelitian. Metoda harus dipaparkan dengan jelas sesuai dengan standar topik penelitian dan dapat diulang oleh peneliti lain. Apabila metoda yang digunakan adalah metoda yang sudah baku cukup ditulis sitasi dan apabila ada modifikasi harus dituliskan dengan jelas bagian mana dan apa yang dimodifikasi.
6. **Hasil**
Sebutkan hasil-hasil utama yang diperoleh berdasarkan metoda yang digunakan. Apabila ingin mengacu pada tabel/grafik/diagram atau gambar uraikan hasil yang terpenting dan jangan menggunakan kalimat 'Lihat Tabel 1'. Apabila menggunakan nilai rata-rata harus menyebutkan standar deviasi.
7. **Pembahasan**
Jangan mengulang isi hasil. Pembahasan mengungkap alasan didapatkannya hasil dan apa arti atau makna dari hasil yang didapat tersebut. Bila memungkinkan, bandingkan hasil penelitian ini dengan membuat perbandingan dengan studi terdahulu (bila ada).
8. **Kesimpulan**
Menyimpulkan hasil penelitian, sesuai dengan tujuan penelitian, dan penelitian berikut yang bisa dilakukan.
9. **Ucapan terima kasih**
10. **Daftar pustaka**
Tidak diperkenankan untuk mensitasi artikel yang tidak melalui proses peer review. Apabila harus menyitir dari "Laporan" atau "komunikasi personal" dituliskan '*unpublished*' dan tidak perlu ditampilkan di daftar pustaka. Daftar pustaka harus berisi informasi yang *up to date* yang sebagian besar berasal dari *original papers*. Penulisan terbitan berkala ilmiah (nama jurnal) tidak disingkat.

Format naskah

1. Naskah diketik dengan menggunakan program Word Processor, huruf New Times Roman ukuran 12, spasi ganda kecuali Abstrak. Batas kiri-kanan atas-bawah masing-masing 2,5 cm. Maksimum isi naskah 15 halaman termasuk ilustrasi dan tabel.
2. Penulisan bilangan pecahan dengan koma mengikuti bahasa yang ditulis menggunakan dua angka desimal di belakang koma. Apabila menggunakan bahasa Indonesia, angka desimal menggunakan koma (,) dan titik (.) bila menggunakan bahasa Inggris. Contoh: Panjang buku adalah 2,5cm. Length of the book is 2.5 cm. Penulisan angka 1-9 ditulis dalam kata kecuali bila bilangan satuan ukur, sedangkan angka 10 dan seterusnya ditulis dengan angka. Contoh lima orang siswa, panjang buku 5 cm.
3. Penulisan satuan mengikuti aturan *international system of units*.
4. Nama takson dan kategori taksonomi merujuk kepada aturan standar termasuk yang diakui. Untuk tumbuhan *International Code of Botanical Nomenclature* (ICBN), untuk hewan *International Code of Zoological Nomenclature* (ICZN), untuk jamur *International Code of Nomenclature for Algae, Fungi and Plant* (ICFAFP), *International Code of Nomenclature of Bacteria* (ICNB), dan untuk organisme yang lain merujuk pada kesepakatan Internasional. Penulisan nama takson lengkap dengan nama author hanya dilakukan pada bagian deskripsi takson, misalnya pada naskah taksonomi. Sedangkan penulisan nama takson untuk bidang lainnya tidak perlu menggunakan nama author.
5. Tata nama di bidang genetika dan kimia merujuk kepada aturan baku terbaru yang berlaku.
6. Ilustrasi dapat berupa foto (hitam putih atau berwarna) atau gambar tangan (*line drawing*).
7. **Tabel**
Tabel diberi judul yang singkat dan jelas, spasi tunggal dalam bahasa Indonesia dan Inggris, sehingga Tabel dapat berdiri sendiri. Tabel diberi nomor urut sesuai dengan keterangan dalam teks. Keterangan Tabel diletakkan di bawah Tabel. Tabel tidak dibuat tertutup dengan garis vertikal, hanya menggunakan garis horisontal yang memisahkan judul dan batas bawah. Paragraf pada isi tabel dibuat satu spasi.
8. **Gambar**
Gambar bisa berupa foto, grafik, diagram dan peta. Judul ditulis secara singkat dan jelas, spasi tunggal. Keterangan yang menyertai gambar harus dapat berdiri sendiri, ditulis dalam bahasa Indonesia dan Inggris. Gambar dikirim dalam bentuk .jpeg dengan resolusi minimal 300 dpi.
9. **Daftar Pustaka**
Sitasi dalam naskah adalah nama penulis dan tahun. Bila penulis lebih dari satu menggunakan kata 'dan' atau *et al*. Contoh: (Kramer, 1983), (Hamzah dan Yusuf, 1995), (Premachandra *et al.*, 1992). Bila naskah ditulis dalam bahasa Inggris yang menggunakan sitasi 2 orang penulis

maka digunakan kata 'and'. Contoh: (Hamzah and Yusuf, 1995).

a. Jurnal

Nama jurnal ditulis lengkap.

Premachandra GS, H Saneko, K Fujita and S Ogata. 1992. Leaf Water Relations, Osmotic Adjustment, Cell Membrane Stability, Epicuticular Wax Load and Growth as Affected by Increasing Water Deficits in Sorghum. *Journal of Experimental Botany* **43**, 1559-1576.

b. Buku

Kramer PJ. 1983. *Plant Water Relationship*, 76. Edisi ke-(bila ada). Academic, New York.

c. Prosiding atau hasil Simposium/Seminar/Lokakarya.

Hamzah MS dan SA Yusuf. 1995. Pengamatan Beberapa Aspek Biologi Sotong Buluh (*Sepioteuthis lessoniana*) di Sekitar Perairan Pantai Wokam Bagian Barat, Kepulauan Aru, Maluku Tenggara. *Prosiding Seminar Nasional Biologi XI*, Ujung Pandang 20-21 Juli 1993. M Hasan, A Mattimu, JG Nelwan dan M Litaay (Penyunting), 769-777. Perhimpunan Biologi Indonesia.

d. Makalah sebagai bagian dari buku

Leegood RC and DA Walker. 1993. Chloroplast and Protoplast. In: *Photosynthesis and Production in a Changing Environment*. DO Hall, JMO Scurlock, HR Bohlar Nordenkamp, RC Leegood and SP Long (Eds), 268-282. Chapman and Hall. London.

e. Thesis dan skripsi.

Keim AP. 2011. Monograph of the genus *Orania* Zipp. (Arecaceae; Oraniinae). University of Reading, Reading. [PhD. Thesis].

f. Artikel online.

Artikel yang diunduh secara online mengikuti format yang berlaku misalnya untuk jurnal, buku atau thesis, serta dituliskan alamat situs sumber dan waktu mengunduh. Tidak diperkenankan untuk mensitasi artikel yang tidak melalui proses *peer review* atau artikel dari laman web yang tidak bisa dipertanggung jawabkan kebenarannya seperti wikipedia.

Forest Watch Indonesia[FWI]. 2009. Potret keadaan hutan Indonesia periode 2000-2009. <http://www.fwi.or.id>. (Diunduh 7 Desember 2012).

Formulir persetujuan hak alih terbit dan keaslian naskah

Setiap penulis yang mengajukan naskahnya ke redaksi Berita Biologi akan diminta untuk menandatangani lembar persetujuan yang berisi hak alih terbit naskah termasuk hak untuk memperbanyak artikel dalam berbagai bentuk kepada penerbit Berita Biologi. Sedangkan penulis tetap berhak untuk menyebarkan edisi cetak dan elektronik untuk kepentingan penelitian dan pendidikan. Formulir itu juga berisi pernyataan keaslian naskah, yang menyebutkan bahwa naskah adalah hasil penelitian asli, belum pernah dan sedang diterbitkan di tempat lain.

Penelitian yang melibatkan hewan

Untuk setiap penelitian yang melibatkan hewan sebagai obyek penelitian, maka setiap naskah yang diajukan wajib disertai dengan 'ethical clearance approval' terkait *animal welfare* yang dikeluarkan oleh badan atau pihak berwenang.

Lembar ilustrasi sampul

Gambar ilustrasi yang terdapat di sampul jurnal Berita Biologi berasal dari salah satu naskah. Oleh karena itu setiap naskah yang ada ilustrasi harap mengirimkan ilustrasi dengan kualitas gambar yang baik disertai keterangan singkat ilustrasi dan nama pembuat ilustrasi.

Proofs

Naskah *proofs* akan dikirim ke author dan diwajibkan membaca dan memeriksa kembali isi naskah dengan teliti. Naskah proofs harus dikirim kembali ke redaksi dalam waktu tiga hari kerja.

Naskah cetak

Setiap penulis yang naskahnya diterbitkan akan diberikan 1 eksemplar majalah Berita Biologi dan reprint. Majalah tersebut akan dikirimkan kepada *corresponding author*.

Pengiriman naskah

Naskah dikirim dalam bentuk .doc atau .docx.

Alamat kontak: Redaksi Jurnal Berita Biologi, Pusat Penelitian Biologi-LIPI

Cibinong Science Centre, Jl. Raya Bogor Km. 46 Cibinong 16911

Telp: +61-21-8765067

Fax: +62-21-87907612, 8765063, 8765066

Email: jurnalberitabiologi@yahoo.co.id

berita.biologi@mail.lipi.go.id

BERITA BIOLOGI

Vol. 16 (1)

Isi (*Content*)

April 2017

MAKALAH HASIL RISET (ORIGINAL PAPERS)

INDUKSI BIAK KALUS DAN BIAK SUSPENSI SEL *Aquilaria malaccensis* Lam. [Induction of Callus Culture and Cell Suspension Culture of *Aquilaria malaccensis* Lam.]

Aryani Leksonowati, Witjaksono dan Diah Ratnadewi 1 - 11

BAKTERI ENTOMOPATOGEN SEBAGAI AGEN BIOKONTROL TERHADAP LARVA *Spodoptera litura* (F.) [Entomopathogenic Bacteria as Biocontrol Agent Against *Spodoptera litura* (F.) Larvae]

Ni Putu Ratna Ayu Krishanti, Bramantyo Wikantyo, Apriwi Zulfitri dan Deni Zulfiana 13 - 21

PENINGKATAN PERTUMBUHAN PADI VAR. CIHERANG SETELAH DIINOKULASI DENGAN *Azospirillum* MUTAN MULTIFUNGSI PENAMBAT N₂, PELARUT P DAN PENGHASIL FITOHORMON INDOLE ACETIC ACID (IAA) [The growth enhancement of rice var. Ciherang after inoculated with *Azospirillum* mutants multifunction capable of N₂-fixation, P solubilization, and producing phytohormone indole acetic acid (IAA)]

Eny Ida Riyanti dan Edy Listanto 23 - 30

KUALITAS SEMEN BEKU DOMBA GARUT (*Ovis aries*) PADA PENAMBAHAN SUKROSA DALAM PENGECER SEMEN TRIS KUNING TELUR [The Quality of Garut Ram (*Ovis aries*) Frozen Semen In Tris Egg Yolk Extender to The Sucrose Supplementation]

Herdis Suharman 31 - 38

PENGELOLAAN AIR, BAHAN ORGANIK DAN VARIETAS ADAPTIF UNTUK MENINGKATKAN HASIL PADI DI LAHAN RAWA PASANG SURUT [Water Management, Organic Matter Application and Using Adaptable Variety to Increase Rice (*Oryza sativa* L.) Productivity on Tidal Swamp Land]

Koesrini dan Khairil Anwar 39 - 46

POTENSI SERAPAN CO₂ PADA BEBERAPA JENIS KANTONG SEMAR (*Nepenthes* spp.) DATARAN RENDAH [Potency of CO₂ Absorption of Lowland Pitcher Plants (*Nepenthes* spp.)]

Muhammad Mansur 47 - 57

CLONING, EXPRESSION, AND PARTIAL PURIFICATION OF PLANTARICIN W LOCUS PRODUCED BY *Lactobacillus plantarum* S34 [Kloning, Ekspresi, dan Purifikasi Parsial Lokus Plantarisin W Diproduksi oleh *Lactobacillus plantarum* S34]

Rifqiyah Nur Umami, Apon Zaenal Mustopa, Linda Sukmarini, Hasim Danuri, Andini Setyanti Putri, and Krisna Dwi Aria Wibowo 59 - 67

MIKROBA ENDOFIT DARI TANAMAN SRIKAYA (*Annona squamosa* L.) SEBAGAI PENGHASIL ANTI-MIKROBA *Staphylococcus aureus* DAN *Candida albicans* [Antimicrobial activity of endophytic microbes from sugar-apple (*Annona squamosa* L.) plant against *Staphylococcus aureus* and *Candida albicans*]

Ruth Melliawati dan Sunifah 69 - 83

KARAKTERISASI PISANG REJANG TETRAPLOID HASIL INDUKSI DENGAN ORYZALIN [Characterization of tetraploid Pisang Rejang induced by oryzalin]

Yuyu S. Poerba, T Handayani dan Witjaksono 85 - 93

KOMUNIKASI PENDEK

CATATAN KEKAYAAN JENIS GASTROPODA DI PESISIR PULAU LETI, KAWASAN BANDA SELATAN [Note on Species Richness of Gastropoda in Coastal Area of Leti Island, Southern Banda]

Muhammad Masrur Islami 95 - 99

KEANEKARAGAMAN KEONG DI PULAU ENGGANO, BENGKULU UTARA [The snails diversity in Enggano Island, Northern Bengkulu]

Heryanto 101 - 110