

# Peroxisome Proliferator – Activated Receptors and The Metabolic Syndrome

Anna Meiliana<sup>1,2\*</sup> and Andi Wijaya<sup>1,2\*</sup>

## Abstract

**BACKGROUND:** Obesity is a growing threat to global health by virtue of its association with insulin resistance, inflammation, hypertension, and dyslipidemia, collectively known as the metabolic syndrome (MetS). The nuclear receptors PPAR $\alpha$  and PPAR $\gamma$  are therapeutic targets for hypertriglyceridemia and insulin resistance, respectively, and drugs that modulate these receptors are currently in clinical use. More recent work on the PPAR $\delta$  has uncovered a dual benefit for both hypertriglyceridemia and insulin resistance, highlighting the broad potential of PPARs in the treatment of metabolic disease.

**CONTENT:** We have learned much about PPARs, the metabolic fat sensors, and the molecular pathways they regulate. Through their distinct tissue distribution and specific target gene activation, the three PPARs together control diverse aspects of fatty acid metabolism, energy balance, insulin sensitivity glucose homeostasis, inflammation, hypertension and atherosclerosis. These studies have advanced our understanding of the etiology for the MetS. Mechanisms revealed by these studies highlight the importance of emerging concepts, such as the endocrine function of adipose tissue, tissue-tissue cross-talk and lipotoxicity, in the pathogenesis of type 2 diabetes mellitus and CVD.

**SUMMARY:** The elucidation of key regulators of energy balance and insulin signaling have revolutionized our understanding of fat and sugar

metabolism and their intimate link. The three 'lipid-sensing' (PPAR $\alpha$ , PPAR $\gamma$  and PPAR $\delta$ ) exemplify this connection, regulating diverse aspects of lipid and glucose homeostasis, and serving as bonafide therapeutic targets.

**KEYWORDS:** Peroxisome Proliferator, Activated Receptor, Metabolic Syndrome.

## Introduction

The metabolic syndrome, is characterized by abdominal obesity, atherogenic dyslipidemia, hypertension, insulin resistance, inflammation, and prothrombotic states (1).

The major sequelae are cardiovascular disease and type 2 diabetes mellitus, but the syndrome also increases the risk of polycystic ovary syndrome, fatty liver, cholesterol gallstones, asthma, sleep disturbances, and some forms of cancer (2). The pathogenesis of the MetS is thought to involve a complex interaction of multiple factors, which include obesity and abnormal fat distribution; insulin resistance; hepatic, vascular, and immunologic factors; and lifestyle and genetic contributions (3).

Recent advances in our understanding of adipose tissue biology and, in particular, its endocrine function and the dysregulated state associated with obesity characterized by enlarged adipose cells have provided insight into the mechanisms involved (4).

<sup>1</sup>Post Graduate Program in Clinical Biochemistry, Hasanuddin University

<sup>2</sup>Prodia Clinical laboratory

\*Address correspondence to this author at: Prodia Clinical Laboratory, Jl. Cisangkuy No. 2 Bandung; e-mail: anna\_m@prodia.co.id, andi\_w@prodia.co.id

Increased adipose tissue mass contributes to augmented secretion of proinflammatory adipokines, particularly tumor necrosis factor- $\alpha$  (TNF $\alpha$ ), along with diminished secretion of the “protective” adiponectin. TNF $\alpha$  and adiponectin are antagonistic in stimulating nuclear transcription factor- $\kappa$ B (NF- $\kappa$ B) activation. Through this activation, TNF $\alpha$  induces oxidative stress, which exacerbates pathological processes leading to oxidized low-density lipoprotein and dyslipidemia, glucose intolerance, insulin resistance, hypertension, endothelial dysfunction, and atherogenesis. Elevated free fatty acid, glucose, and insulin levels enhance this NF- $\kappa$ B activation and further downstream modulate specific clinical manifestations of metabolic syndrome (5).

Although there has been a debate on the criteria and concept of the metabolic syndrome, the current definition by the National Cholesterol Education Program-Adult Treatment Panel (NCEP-ATP III) and the International Diabetes Federation (IDF) provide adequate screening tools to identify the subjects with high cardiometabolic risk. With these tools in hand the stage is set for attempts to discover the pathophysiology underlying these metabolic abnormalities. The identification of intracellular signaling elements and regulating factors at crossroad steps that direct the metabolic fate of lipids are critical for the understanding of atherogenic dyslipidemia in the MetS. Importantly, several lipid metabolites seem to play a crucial role in the regulation of insulin signaling and action influencing endothelial function and initiating vascular injury. The concept that dysfunctional adipose tissue cannot properly handle the energy surplus derived from excessive calorie consumption combined with sedentary lifestyle sets the stage to identify the main determinants of the MetS in different populations. Resolving these issues is crucial for the optimal management of the MetS and reduction of global CVD risk (6).

Peroxisome proliferator-activated receptor (PPAR)s are a family of 3 (PPAR $\alpha$ ,  $\beta/\delta$ , and  $\gamma$ ) nuclear receptor/ligand-activated transcription factors that work in concert as heterodimers with the retinoid X receptors (7). In recent years, there has been great scientific and clinical interest in the actions of PPAR $\alpha$  and PPAR $\gamma$  because they are the molecular targets for the clinically used lipid-lowering fibrates and insulin-sensitizing thiazolidinedione classes of drugs, respectively (7,8).

The recent development of highly selective ligands and PPAR $\beta/\delta$  knockout and transgenic mice, however, have now implicated roles for PPAR $\beta/\delta$  in adipose tissue formation, metabolism, wound healing, brain development, placental function, colorectal carcinogenesis, and skeletal muscle function. PPAR $\beta/\delta$  ligands appear highly effective in regulating lipid metabolism, particularly in skeletal muscle, and are currently in phase II clinical trials for treatment of dyslipidemia, aimed particularly at individuals with low HDL levels (9,10,11). All of the PPARs, therefore, appear to be able to target aspects of the MetS (11). Because the MetS represents a major risk factor for cardiovascular diseases, there has been an increasing interest in the roles of PPARs, in particular most recently PPAR $\beta/\delta$ , in vascular biology. Indeed, in addition to the treatment of dyslipidemia, PPAR $\beta/\delta$  ligands may reduce the development in atherosclerosis (11,12).

The discovery cycle involving nuclear receptors has elucidated the molecular and physiological basis for a new class of pharmacophores that show promise for treating the MetS (3).

---

## The Biology of PPARs

---

PPARs are members of the nuclear receptor superfamily which includes the steroid, retinoid, and thyroid hormone receptors. These receptors are transcription factors that regulate gene expression in response to certain endogenous and exogenous ligands. It is now well established that the PPARs act as central transcriptional mediators in the regulation of several important metabolic processes that influence adipogenesis, insulin sensitivity, glucose homeostasis, lipid metabolism, vascular endothelial function, atherosclerosis progression, and ultimately cardiovascular risk (13).

### MOLECULAR BIOLOGY OF PPARS

The three PPARs are ligand-activated transcription factors of the nuclear hormone receptor superfamily. They share a high degree of structural homology with all members of the superfamily, particularly in the DNA-binding domain and ligand- and cofactor-binding domain. Many cellular and systemic roles have been attributed to these receptors, reaching far

beyond the stimulation of peroxisome proliferation in rodents after which they were initially named. PPARs exhibit broad, isotype-specific tissue expression patterns (11).

PPAR $\alpha$ (NR1C1)(NuclearReceptorsNomenclature Committee, 1999) was first described as a receptor that is activated by peroxisome proliferators, hence its name (14). Two additional related isotypes, PPAR $\beta$ / $\delta$  (NR1C2) and PPAR $\gamma$  (NR1C3), were then found and characterized. The PPAR $\beta$ / $\gamma$  isotype was called PPAR $\beta$  when it was first isolated from a *Xenopus* oocyte library (15). Because the mammalian PPAR $\beta$  protein sequence was not highly homologous to the *Xenopus* PPAR $\beta$  protein sequences, it was named PPAR $\beta$  when identified in the mouse with the view that there may be four members of this nuclear receptor family (16). PPAR $\beta$  was also designated FAAR (fatty acid activated receptor) (17) in rats and NUC1 in humans (18). Sequencing of mammalian genomes indicated that there are only three PPAR isotypes. Characterization of PPARs in the chick and comparison with the PPARs of mouse and *Xenopus* demonstrated that the mammalian PPAR $\beta$  is the ortholog of the amphibian PPAR $\beta$ . For reasons of clarity, we propose that this receptor be designated herein as PPAR $\beta$ / $\delta$  (16).

PPAR $\alpha$  is expressed at high levels in organs that carry out significant catabolism of fatty acids such as the brown adipose tissue, liver, heart, kidney, and intestine (19). Of the three isotypes, PPAR $\beta$ / $\delta$  has the broadest expression pattern, and the levels of expression in certain tissues depend on the extent of cell proliferation and differentiation. Important functions have been assigned to this isotype in the skin, gut, placenta, skeletal muscle, adipose tissue, and brain (10,11,20,21,22). PPAR $\gamma$  is expressed as two isoforms,  $\gamma$ 1 and  $\gamma$ 2, that differ at their N terminus. PPAR $\gamma$ 2 is found at high levels in the different adipose tissues (15,23,24), whereas PPAR $\gamma$ 1 has a broader expression pattern that extends to settings such as the gut, brain, vascular cells, and specific kinds of immune and inflammatory cells (25,26).

PPARs requires heterodimerization with the retinoid X receptor (RXR; NR2B), which belongs to the same receptor superfamily (27,28). This PPAR/RXR heterodimer can form in the absence of a ligand. When activated by a ligand, it modulates transcription via binding to a specific DNA sequence element frequently called a peroxisome proliferator response element (PPRE) (15,27,29,30). This response element, generally of the direct repeat 1 (DR-1) type, is composed of two half-sites that occur as a direct

repetition of the consensus sequence AGGTCA with a single nucleotide spacing between the two repeats. The PPRE is usually present in one or multiple copies in the promoter region of target genes but may also be located in the proximal transcribed region of certain PPAR-responsive genes (31). PPAR and RXR bind to the 5' and 3' half-sites of this element, respectively, and the 5'-flanking region mediates the selectivity of binding between different PPAR isotypes (32,33,34). Transcriptional control by PPAR/RXR heterodimers requires interaction with coregulator complexes—either a coactivator for stimulation or a corepressor for inhibition of target gene expression (35,36,37,38).

Thus, selective action of PPARs *in vivo* results from the interplay at a given time point between expression levels of each of the three PPAR and RXR isotypes, affinity for a specific promoter PPRE, and ligand and cofactor availabilities (11).

A wide variety of natural or synthetic compounds was identified as PPAR ligands. Among the synthetic ligands, the lipid – lowering drugs, fibrates, and the insulin sensitizers, thiazolidinediones, are PPAR $\alpha$  and PPAR $\gamma$  agonists, respectively, which underscores the important role of PPARs as therapeutic targets (11).

The prevalent point of view today is that PPARs act as lipid sensors that translate changes in lipid / fatty acid levels from the diet or from food deprivation into metabolic activity, leading to either fatty acid catabolism or lipid storage. The endogenous ligands or mediators of these changes have not been characterized but are probably generated by fatty acid metabolism. Their activities are likely to be influenced by their binding specificities toward the different PPARs and by cell-, tissue-, or organ- specific effects (39,40,41).

The possible pathways that generate lipid mediators from fatty acids, which also serve as PPAR ligands, are recapitulated in Fig. 1. In addition, specific lipolytic pathways, for example the action of lipoprotein lipase and endothelial lipase, can hydrolyze certain circulating lipoproteins to generate PPAR ligands and PPAR activation (43,44,45). Given the variety and distribution pattern in the body of fatty acids and fatty acid derivatives with a wide range of affinity to PPARs, it has been difficult thus far to thoroughly evaluate the contribution of each of these endogenous ligands to the biology of PPARs. However, it is not surprising, based on the characteristics of these endogenous ligands with their broad spectrum of activation efficiency, that PPARs are involved in functions as diverse as lipid and carbohydrate metabolism, immune/inflammatory responses, vascular biology, tissue repair, and cell

differentiation and proliferation. The distribution and abundance of the ligands also depend on a variety of pathophysiological situations associated with hyperlipidemia, hypertension, diabetes, chronic inflammation, cancer, and atherosclerosis. It is important to note that some of these endogenous lipid mediators also signal through the classic cell surface G-protein-linked receptors and therefore have many PPAR-independent effects (11).

PPAR $\gamma$  is expressed in adipocytes, macrophages, and muscle, where it regulates development, lipid homeostasis, and glucose metabolism. Endogenous PPAR $\gamma$  agonists include fatty acids and eicosanoids (47,48,49). The PPAR $\gamma$  genetic program includes target genes involved in the uptake of glucose in muscle (c-Cbl associated protein and glucose transporter 4), lipid metabolism (scavenger receptor, adipocyte-fatty-acid-binding protein, lipoprotein lipase, fatty-acid-binding




Fig. 1. Endogenous pathways for PPAR ligand production (Adapted from Michalik L, et al. 2006).

Among synthetic ligands, compounds of the fibrate family (clofibrate, fenofibrate, and bezafibrate) and their derivatives have been widely used to characterize PPAR $\alpha$  functions (43). Taken together, mouse models suggest that PPAR $\alpha$  functions to increase fatty-acid use in the fasting state and that in the pathophysiological context of a high-fat diet, PPAR $\alpha$ -induced fatty-acid catabolism might prevent hypertriglyceridemia. Consistent with this prediction, an activated variant of PPAR $\alpha$  (Leu162Val) is associated with low serum TG levels and reduced adiposity (46).

protein, a cy1-CoA synthetase, and CYP4B1), and energy expenditure (glycerol kinase and uncoupling proteins 2 and 3) (50,51,52,63,54,55,56,57,58).

In addition to regulating glucose and lipid metabolism, PPAR $\gamma$  is a potential modifier of atherogenesis. Signaling through PPAR $\gamma$ , components of oxidized low-density lipoprotein (LDL) increase expression of the scavenger receptor CD36, resulting in lipid accumulation in macrophages (59,60). PPAR $\gamma$  also activates the macrophage LXR-ABCA1 cholesterol efflux pathway (61), which may explain

the finding that PPAR $\gamma$  ligands inhibit the formation of atherosclerotic lesions in LDL-receptor-deficient mice (62).

The landmark finding that the thiazolidinedione class of insulin sensitizers, including rosiglitazone and pioglitazone, function as high-affinity PPAR $\gamma$  agonists has validated the efficacy of PPAR $\gamma$  modulation in treating the MetS (63).

PPAR $\delta$  is expressed ubiquitously and is activated by fatty acids and components of very-low-density lipoprotein (VLDL) (43,47). PPAR $\delta$  target genes control  $\beta$ -oxidation in murine brown fat (long-chain and very-long-chain acyl-CoA synthetase, long-chain and very-long-chain acyl-CoA dehydrogenase, and acyl-CoA oxidase), energy expenditure (uncoupling proteins 1 and 3), and lipid storage (macrophage adipose differentiation-related protein) (64,65). In the pathophysiological context of a high-fat diet, PPAR $\delta$  could function to increase adipose fatty-acid catabolism and may play a role in VLDL-induced

lipid accumulation in atherosclerotic foam cells. A high-affinity synthetic PPAR $\delta$  agonist has been shown to increase HDL and decrease LDL, TG, and fasting insulin in obese rhesus monkeys. These studies suggest that therapeutic activation of PPAR $\delta$  has the potential to decrease diet-induced obesity without activating the PPAR $\gamma$  dependent adipogenic program (66).

PPARs AND CONTROL OF METABOLISM

Lipids are essential for energy homeostasis, reproductive and organ physiology, and numerous aspects of cellular biology. They are also linked to many pathological processes, such as obesity, diabetes, heart disease, and inflammation. To meet the different demands from a variety of tissues, the human body has evolved a sophisticated lipoprotein transport system to deliver cholesterol and fatty acids to the periphery (Fig. 2) (67).




Fig. 2 Circulating lipoproteins deliver both energy substrates and endogenous activators for PPARs (Adapted from Lee CH, et al, 2003).

Disturbances in this system are integral components of life-threatening diseases, best exemplified by the metabolic syndrome, which refers to patients who are insulin-resistant (hyperinsulinemic), dyslipidemic (elevated TG and decreased HDLC levels) (68).

Lipids and their derivatives have a role in the genetic control of their own systemic transport, cellular uptake, storage, mobilization, and use. Fine tuning of these metabolic processes is a hallmark of healthy organisms. Lipid homeostasis depends on factors that are able to transduce metabolic parameters into regulatory events representing the fundamental components of the general control system (69).

The identification of fatty acids as endogenous ligands for PPARs has provided a unique approach to study lipid homeostasis at the molecular level (68).

Liver is the key site of metabolic integration where fatty acids are mobilized and, depending on the body's needs, either stored or used as an energy source. In the fasting state, the fuel sources of the body shift from carbohydrates and fats to mostly fats, and fatty acids that were stored during feeding are released from the adipocyte and taken up by liver. There they are either reesterified to TGs and assembled into VLDL or broken down through  $\beta$ -oxidation and used to generate ketone bodies. Earlier studies have demonstrated that in the liver, PPAR $\alpha$  directly regulates genes involved in fatty acid uptake [fatty acid binding protein (FABP)],  $\beta$ -oxidation (acyl-CoA oxidase) and  $\omega$ -oxidation (cytochrome P450). Gene targeting studies confirmed that PPAR $\alpha$  is essential for the up-regulation of these genes caused by fasting or by pharmacological stimulation with synthetic ligands such as the fibrates (70,71,72).

Fasting also results in severe hypoglycemia, hypoketonemia, and elevated plasma levels of nonesterified fatty acid, indicating a defect in fatty acid uptake and oxidation caused by dysregulation of these genes (73,74). In line with these observations, the fibrate class of drugs including fenofibrate and gemfibrozil, which are synthetic ligands for PPAR $\alpha$ , lower serum TGs and slightly increase HDL cholesterol levels in patients with hyperlipidemia (75), most likely due to induction of fatty acid oxidation through activation of PPAR $\alpha$ . PPAR $\alpha$  has also been shown to down-regulate apolipoprotein C-III, a protein which inhibits TG hydrolysis by LPL. This activity of PPAR $\alpha$  ligands further contributes to the lipid-lowering effect (68).

Apolipoprotein A-V (apoA-V) is now recognized as a key regulator of serum TG levels. Administration

of the PPAR $\alpha$  agonist caused a 50% decrease in TG that reversed at washout. Serum apoA-V concentrations increased 2-fold, correlated inversely with TG, and were reversible at washout. The apoA-V/apoC-III ratio increased 2-fold, with this increase also reversible at washout. These data demonstrate for the first time that a PPAR $\alpha$  agonist increases circulating apoA-V protein levels and the apoA-V/apoC-III ratio (76).

Unlike its function in the adaptive response to fasting, the role of PPAR $\alpha$  in cardiovascular pathogenesis appears to be detrimental. Cardiac-specific PPAR $\alpha$  overexpression increases fatty acid oxidation and concomitantly decreases glucose transport and use, a phenotype similar to that of the diabetic heart, indicate that PPAR $\alpha$  senses fatty acids and induces their use, and thus plays a causative role in cardiomyopathy. The net effect, however, of fibrate intervention in cardiovascular disease is likely beneficial because systemic TG reduction should result in less fat accumulation in the heart and at the vessel wall (68).

Adipocytes are the main site for lipid storage and modulate the levels of lipids in the blood stream in response to hormonal signals. PPAR $\gamma$  has high expression in this tissue and has been shown to potentiate adipocyte differentiation from fibroblasts (54). The PPAR $\gamma$ 2 isoform prevents lipotoxicity by (a) promoting adipose tissue expansion, (b) increasing the lipid-buffering capacity of peripheral organs, and (c) facilitating the adaptive proliferative response of  $\beta$ -cells to insulin resistance (77).

PPAR $\gamma$  also promotes cholesterol efflux through the induction of a transcriptional cascade involving the nuclear sterol receptor LXR $\alpha$  and its downstream target ABCA1, a membrane transporter that is important for HDL-mediated reverse cholesterol transport (61,78,79,80). In this view, one would predict that in the absence of proportionately increased ox-LDL, pharmacological activation of PPAR $\gamma$  should shift the balance from lipid loading to lipid efflux and improve the status of the atherosclerotic lesion (68).

PPAR $\delta$  has recently been shown to mediate VLDL signaling in the macrophage (43). VLDL treatment in cultured macrophages results in lipid accumulation and up-regulation of adipose differentiation-related protein, a lipid droplet-coating protein that has been implicated in lipid storage (81). Adipose differentiation-related protein was subsequently identified as a direct PPAR $\delta$  target gene, and components of VLDL released by LPL serve as ligands for the receptor (68).

Exercise increases fatty acid oxidation (FAO), improves serum HDLC and TG, and upregulates skeletal muscle PPAR $\delta$  expression. In parallel, PPAR $\delta$  agonist-upregulated FAO would induce fatty-acid uptake (via peripheral lipolysis), and influence HDLC and TG-rich lipoprotein particle metabolism, as suggested in preclinical models (82). In their report, Oliver et al demonstrate that a selective PPAR $\delta$  agonist increases ABCA1 expression and cholesterol efflux from cells and increases HDLC in primates. PPAR $\delta$  agonists may provide a new approach to the treatment of cardiovascular disease by promoting reverse cholesterol transport (66).

Molecular and functional analyses suggest that PPAR $\delta$  activation reduces hepatic glucose output by increasing glycolysis and the pentose phosphate shunt and promoting fatty acid synthesis in the liver. This uncovered hepatic activity thus constitutes the earliest component of the regulatory mechanism by which PPAR $\delta$  regulates insulin sensitivity, in addition to its known function in fatty acid  $\beta$ -oxidation (83).

Coupling increased hepatic carbohydrate catabolism with its ability to promote  $\beta$ -oxidation in muscle allows PPAR $\delta$  to regulate metabolic homeostasis and enhance insulin action by complementary effects in distinct tissues. The combined hepatic and peripheral actions of PPAR $\delta$  suggest new therapeutic approaches to treat type II diabetes (83).

It is now evident that PPARs, which are activated by various lipid molecules, function in distinct target tissues and coordinately regulate different metabolic pathways. PPAR $\alpha$  and PPAR $\delta$  potentiate fatty acid use in liver and muscle, respectively, whereas PPAR $\gamma$  promotes lipid storage in adipocytes. In this dynamic system, lipids are shuttled between these tissues according to the needs of the body by lipoproteins. In this view, lipoproteins not only deliver energy substrates but also carry endogenous activators for these receptors (68).

#### PPARs AND OBESITY

The cluster of medical sequelae, collectively referred to as the MetS, poses one of the most serious threats to public health that our society faces. Why does obesity carry with it a stereotyped collection of medical problems? That is, why is adipose tissue unable to store excess calories in a safe way? The answer to this question is not clear, but it may reflect the reality that fat is not simply a storage depot, but rather a dynamic tissue that constantly communicates with

other key tissues in the body, including liver, muscle and the appetite centers in the brain. In the past decade, tremendous advances in understanding this system of signals and sensors have been made, and the emergence of PPARs as key regulators of obesity and metabolism shedding light on how problems begin and how they may be therapeutically approached (41).

The initial suggestion that PPAR $\gamma$  stimulated adipogenesis was based on the observation that overexpression of PPAR $\gamma$  in cells was by itself sufficient to induce adipocyte differentiation (54). Consistent with this, PPAR $\gamma$  increases the expression of genes that promote fatty acid storage, whereas it represses genes that induces lipolysis in adipocytes (84).

PPAR- $\gamma$  is a master regulator in the formation of fat cells and their ability to function normally in the adult (85). PPAR $\gamma$  is induced during adipocyte differentiation, and forced expression of PPAR $\gamma$  in nonadipogenic cells effectively converts them into mature adipocytes (54). In addition, PPAR $\gamma$  knockout mice fail to develop adipose tissue (86,87,88). Consistent with these findings, humans with dominant-negative mutations in a single allele PPARG (the gene encoding PPAR $\gamma$ ) have partial lipodystrophy and insulin resistance (89,90,91). In vitro studies suggest that PPAR $\gamma$  is the ultimate effector of adipogenesis in a transcriptional cascade that also involves members of the C/EBP transcription factor family (92,93).

Fat cells develop from a fibroblast-like preadipocyte to a mature, lipid-enriched adipocyte. The underlying transcriptional regulatory network that controls the maturation of adipocytes has been the focus of intense research and is reviewed elsewhere (94).

The highest levels of PPAR $\gamma$  are expressed in adipose tissue (23,24). In 1994, Spiegelman and colleagues discovered that expression and activation of PPAR $\gamma$  was sufficient to induce adipogenesis (54). The essential role of PPAR $\gamma$  in adipogenesis has been clearly demonstrated in functional and genetic knockdown experiments (86,87).

In addition to its importance in adipogenesis, PPAR $\gamma$  plays an important role in regulating lipid metabolism in mature adipocytes. Much of what is known about this role of PPAR $\gamma$ , and indeed many other aspects of its biology, followed the discovery that thiazolidinedione (TZD) antidiabetic drugs are highaffinity agonist ligands for PPAR $\gamma$  (63). TZDs appear to coordinately activate gene expression leading to an increase in net lipid partitioning into adipocytes. Target genes directly regulated by PPAR $\gamma$  that are involved in this pathway include lipoprotein lipase

(55), fatty-acid transport protein (95), and oxidized LDL receptor 1 (96), which all favor adipocyte uptake of circulating fatty acids; phosphoenolpyruvate carboxykinase (97,98), glycerol kinase (58), and the glycerol transporter aquaporin 7 (99), which promote recycling rather than export of intracellular fatty acids. Together, these pathways lead to the net flux of fatty acids from the circulation and other tissues into adipocytes. Although increased fat storage would be expected to boost the size of adipocytes, TZD treatment actually leads to smaller adipocytes (100). This is partly due to increased adipocyte differentiation, leading to new smaller cells. In addition, TZDs induce the coactivator PPAR $\gamma$ -coactivator 1a (PGC-1 $\alpha$ ), which promotes mitochondrial biogenesis (101), leading to an increase in fatty-acid oxidation that further protects against adipocyte hypertrophy (102).

At a molecular level, how does activation of PPAR $\gamma$ , a protein that is mainly present in adipose cells, lead to systemic insulin sensitization? Two plausible mechanisms should be considered. First, pharmacologic activation of PPAR $\gamma$  in adipose tissue improves its ability to store lipids, thereby reducing lipotoxicity in muscle and liver. This model involves activation of genes encoding molecules that promote a combination of lipid storage and lipogenesis, such as aP2 (fatty-acid binding protein), CD36 (receptor for lipoproteins), lipoprotein lipase (hydrolysis of lipoproteins), FATP-1 (fatty acid transporter), glycerol kinase, and SREBP-1 and SCD-1 (regulators of sterol and fatty acid synthesis, respectively). Activation of this metabolic pathway causes body-wide lipid

repartitioning by increasing the TG content of adipose tissue and lowering free fatty acids and TG in the circulation, liver and muscle, thereby improving insulin sensitivity (58, 103). Second, PPAR $\gamma$ -specific drugs alter the release of signaling molecules from fat, including leptin, TNF $\alpha$ , resistin and adiponectin, which by virtue of serum transport have far-reaching metabolic effects in other tissues. For example, PPAR $\gamma$  agonists inhibit the expression of TNF $\alpha$  and resistin, which both promote insulin resistance (58, 104, 111, 105, 106). On the other hand, PPAR $\gamma$  agonists stimulate the production of adiponectin, which promotes fatty acid oxidation and insulin sensitivity in muscle and liver. As a result, hepatic glucose production is reduced and muscle glucose use is increased (107, 108, 109). This pathway is mediated by adiponectin receptors and downstream AMP-activated protein kinase (110, 111), making the intracellular AMP-activated kinase pathway another potential pharmacologic target for type 2 diabetes mellitus. Besides the aforementioned mechanisms, it is possible that PPAR $\gamma$  directly modulates the insulin signal transduction pathway in adipose tissue by increasing the expression of intracellular proteins such as c-Cbl-associated protein (CAP), which stimulates glucose transport (51).

There may be an appropriate level of PPAR $\gamma$ /RXR activity for insulin sensitivity. This raises the possibility that there may be a hitherto unrecognized U-shaped relationship between PPAR $\gamma$ /RXR activity and insulin resistance within physiologically "normal" limits (Figure 3) (112).




Fig. 3. One possible model for the relationship between PPAR $\gamma$ /RXR activity and insulin sensitivity on the HF diet (Adapted from Yamauchi T et al, 2001).

Yamauchi et al have shown herein that PPAR $\gamma$ /RXR promotes fat storage in the body by a combination of direct induction of molecules involved in TG accumulation and suppression of leptin gene expression as well as inactivation of PPAR $\alpha$ -signaling pathways. In times of fasting, this PPAR $\gamma$ /leptin/PPAR $\alpha$  network maximizes energy storage, which is quite advantageous for survival. In times of feast, which are the norm in industrialized nations nowadays, however, this network causes excessive adiposity, insulin resistance, and obesity related diseases such as diabetes. Thus appropriate antagonism of PPAR $\gamma$ /RXR, which simultaneously leads to appropriate agonism of leptin and PPAR $\alpha$ , may be a logical approach to protection against obesity and related diseases such as type 2 diabetes (112).

PPAR $\gamma$ , function as an important adipocyte determination factor. In contrast, TNF $\alpha$  inhibits adipogenesis, cause dedifferentiation of mature adipocytes, and reduces the expression of several adipocyte – specific genes. Reduced PPAR $\gamma$  gene expression is likely to represent an important component of the mechanism by which TNF $\alpha$  exerts its antiadipogenic effects (113).

Insulin receptor substrate (IRS)-1 and IRS-2 have dominant roles in the action of insulin (114), but other substrates of the insulin receptor kinase, such as Gab1, c-Cbl, SH2-B and APS, are also of physiological relevance (115,116,117,118). Although the protein downstream of tyrosine kinases-1 (Dok1) is known to function as a multisite adapter molecule in insulin signaling (119,120,121), its role in energy homeostasis has remained unclear. Hosooka et al show that Dok1 regulates adiposity. Dok1 promotes adipocyte hypertrophy by counteracting the inhibitory effect of ERK on PPAR $\gamma$  and may thus confer predisposition to diet-induced obesity. (122).

The nuclear receptor corepressors NCoR and SMRT repress gene transcription by recruiting a histone deacetylase complex. Their roles in PPAR $\gamma$  action have been controversial. Recent evidence, however, suggests that NCoR and SMRT repress PPAR $\gamma$ -mediated transcriptional activity on specific promoters in the adipocyte. In addition, by repressing PPAR $\gamma$  action, these corepressors inhibit the ability of adipocyte differentiation to proceed (123).

PPAR $\delta$  initially received much less attention than the other PPARs because of its ubiquitous expression and the unavailability of selective ligands. However, genetic studies and recently developed synthetic PPAR $\delta$  agonists have helped reveal its role as a powerful regulator of

fatty acid catabolism and energy homeostasis (9,22). The PPAR $\delta$  agonist GW501516 was shown to lower plasma TG levels in obese monkeys while raising high-density lipoprotein levels, prompting the initiation of clinical trials to assess its efficacy in hyperlipidemic patients (66). Studies in adipose tissue and muscle have begun to uncover the metabolic functions of PPAR $\delta$ . Transgenic expression of an activated form of PPAR $\delta$  in adipose tissue produces lean mice that are resistant to obesity, hyperlipidemia and tissue steatosis induced genetically or by a highfat diet. The activated receptor induces genes required for fatty acid catabolism and adaptive thermogenesis. Interestingly, the transcription of PPAR $\gamma$  target genes for lipid storage and lipogenesis remain unchanged. In parallel, PPAR $\delta$ -deficient mice challenged with a high-fat diet show reduced energy uncoupling and are prone to obesity. Together, these data identify PPAR $\delta$  as a key regulator of fat-burning, a role that opposes the fat-storing function of PPAR $\gamma$  (64). Thus, despite their close evolutionary and structural kinship, PPAR- $\gamma$  and PPAR $\delta$  regulate distinct genetic networks. The thermogenic function of PPAR $\delta$  is markedly similar to that of the nuclear cofactor PGC-1 $\alpha$  (124,125). Indeed, PPAR strongly associates with PGC-1 $\alpha$  both in cultured cells and in tissue, so it is possible that many metabolic effects of PGC-1 $\alpha$  may be mediated through PPAR $\delta$ (41).

Upon agonist stimulation of PPAR $\delta$ , KLF5 was deSUMOylated, and became associated with transcriptional activation complexes containing both the liganded PPAR $\delta$  and CREB binding protein (CBP). This activation complex increased the expression of Cpt1b, Ucp2 and Ucp3. Thus, SUMOylation seems to be a molecular switch affecting function of KLF5 and the transcriptional regulatory programs governing lipid metabolism (126).

Activation of PPAR $\delta$  with the specific agonist GW501516, however, coordinately upregulates expression of genes involved in fatty acid oxidation and energy uncoupling, which markedly diminishes weight gain while increasing O<sub>2</sub> consumption in mice fed a high-fat diet (64, 127,128). Likewise, skeletal muscle-specific overexpression of an activated form of PPAR $\delta$  protects mice from diet-induced obesity. As a result, PPAR $\delta$  is now considered to be an attractive target for treatment of metabolic syndrome (126). PPAR $\delta$  as a nuclear receptor, represents a bonafide target. In particular, its ability to induce adaptive thermogenesis and protection against both dietary and genetic obesity (24).

## PPARs AND CONTROL OF INFLAMMATORY RESPONSES

In addition to the regulation of lipid metabolism, PPARs and LXRs play roles in influencing inflammatory and immune responses. PPARs can be activated by eicosanoids, which are produced by metabolism of arachidonic acids and other long-chain PUFAs during inflammatory responses (36,48,49,59,129). For example, ligands for PPAR $\alpha$  are leukotriene LTB<sub>4</sub> and 8(S)-hydroxyeicosatetraenoic acid (HETE), whereas 15deoxy-prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>), 15-HETE, and 13-hydroxyoctadecadienoic acid (HODE) act as ligands for PPAR $\gamma$ . Interestingly, the expression of PPARs is differentially regulated by factors that control the development of immune responses. PPAR $\gamma$  expression is dramatically upregulated in macrophages and T cells during the inflammatory response and can be induced *in vitro* by interleukin (IL)-4 and other immunoregulatory molecules (130,131). In contrast, IFN $\gamma$  and lipopolysaccharide (LPS) repress the expression of PPAR $\gamma$  (132). PPAR $\gamma$  is highly expressed in elicited peritoneal macrophages, while low levels of PPAR $\alpha$  are present (133). The opposite pattern is observed in primary human monocytes (134). PPAR $\alpha$  and PPAR $\gamma$  have been shown to inhibit the expression of proinflammatory genes, suggesting that they might inhibit inflammatory responses *in vivo*. Activation of PPAR $\alpha$  resulted in the induction of genes involved in fatty acid oxidation with the subsequent degradation of fatty acids and fatty acid derivatives like LTB<sub>4</sub>. In addition, the response to LTB<sub>4</sub> and arachidonic acid was prolonged in mice lacking the PPAR $\alpha$  gene as compared with wild-type mice (135). However, some *in vivo* studies show proinflammatory effects for PPAR $\alpha$  ligands, such as an increase in the plasma levels of TNF $\alpha$  during endotoxemia (136) and in the production of monocyte chemoattractant protein (MCP-1) by endothelial cells (137). Thus, PPARs and LXRs negatively regulate transcriptional programs involved in the development of inflammatory responses (138).

PPAR $\alpha$  activators reduced cytokine-induced expression of VCAM-1 and ICAM-1 in human carotid artery endothelial cells in culture. Pasceri *et al.* (139) reported that fibrates reduced C-reactive protein (CRP)-induced expression of monocyte chemoattractant protein (MCP)-1 in human umbilical vein endothelial cells. With regard to clinical studies, Marchesi *et al.* observed that 3-month therapy with fenofibrate in hyperTGmic patients decreased VCAM-1 and ICAM-1 levels in the fasting state.

A molecular rationale for suppression of IL-1-induced CRP transcription is provided by the fact that fenofibrate upregulates I- $\kappa$ B protein expression. This reduces nuclear translocation of p50-NF- $\kappa$ B, resulting in decreased amounts of nuclear p50-NF- $\kappa$ B and CCAAT/enhancer binding protein- $\beta$  complexes, the major determinants of CRP transcription. These results provide strong evidence for a direct suppressive effect of fenofibrate on CRP expression, independent of cholesterol lowering and atherogenesis (140,141).

Endothelial dysfunction associated with the metabolic syndrome and other insulin-resistant states is characterized by impaired insulin-stimulated nitric oxide production from the endothelium and decreased blood flow to skeletal muscle. Thus, improvement in insulin sensitivity leads to improved endothelial function (142).

Teissier *et al.* extend our insight into the connections between PPARs, LDL metabolism, and oxidative stress (143). The oxidized form of LDL (ox-LDL) promotes inflammation in part via ox-LDL uptake by scavenger receptors and subsequent nuclear factor  $\kappa$ B activation. They find that synthetic PPAR agonists induce the production of reactive oxygen species (ROS) in a PPAR $\alpha$ -dependent manner by inducing NADPH oxidase, a key enzyme in oxidative stress (144). Moreover, these investigators offer the intriguing notion that ROS interact with LDL to activate PPAR $\alpha$  and subsequently limit inflammation, as indicated by PPAR-dependent repression of inducible nitric oxide synthase (iNOS) gene transcription (145).

The PPAR $\alpha$ -induced increase in ROS was attributable to the induction of NADPH oxidase, because (1) preincubation with the NADPH oxidase inhibitor diphenyleneiodonium prevented the increase in ROS production (2) PPAR $\alpha$  agonists increased O<sub>2</sub><sup>-</sup> production measured by superoxide dismutase-inhibitable cytochrome c reduction (3) PPAR $\alpha$  agonists induced mRNA levels of the NADPH oxidase subunits p47<sup>phox</sup>, p67<sup>phox</sup>, and gp91<sup>phox</sup> and membrane p47<sup>phox</sup> protein levels (4) induction of ROS production was abolished in p47<sup>phox</sup><sup>-/-</sup> and gp91<sup>phox</sup><sup>-/-</sup> macrophages. Finally, induction of NADPH oxidase by PPAR $\alpha$  agonists resulted in the formation of oxidized LDL metabolites that exert PPAR $\alpha$ -independent proinflammatory and PPAR $\alpha$ -dependent decrease of lipopolysaccharide-induced inducible nitric oxide synthase expression in macrophages. These data identify a novel mechanism of autogeneration of endogenous PPAR $\alpha$  ligands via stimulation of NADPH oxidase activity (143).

Several reports (146,147,148) have indicated that PPAR $\gamma$  agonists reduce macrophage inflammatory responses, such as the elaboration of cytokines, nitric oxide, and matrix metalloproteinases (MMPs). In contrast, other reports have indicated that PPAR $\gamma$  agonists stimulate the expression of proinflammatory receptors and cytokines (59,60,149,150,151). Finally, there are even reports that PPAR $\gamma$  agonists have no impact at all on macrophage inflammatory cytokine production (150,152).

PPAR $\gamma$  generally inhibits inflammatory response genes by negatively interfering with the NF- $\kappa$ B, STAT, and AP-1 signaling pathways in a DNA-binding independent manner (153). Indeed, PPAR $\gamma$  modulates chemokine gene expression by inhibiting expression of MCP-1 and its receptor CCR2 in monocyte/macrophages (154,155). PPAR $\gamma$  also plays a role in the regulation of CXC chemokine pathway. CXCR2 upregulation by PPAR $\gamma$  leads to the acquisition of IL-8/Gro $\beta$  responsiveness, as measured by enhanced reactive oxygen species (ROS) formation (156).

PPAR $\gamma$  expression in human ECs has been demonstrated by reverse transcription polymerase chain reaction (157) and more definitively using Western blotting and immunohistochemistry (158). Subsequent data suggested that PPAR $\gamma$  activation can influence target genes and processes that are of central relevance to endothelial biology. One such example is the role of chemokines (chemoattractant cytokines), signals for inflammatory cell recruitment (158).

Recently, Huang et al demonstrated that the PPAR $\gamma$  ligands, either 15-deoxy- $\Delta$  prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>) or ciglitazone, increased endothelial nitric oxide ( $\bullet$ NO) release without altering endothelial nitric oxide synthase (eNOS) expression. However, the precise molecular mechanisms of PPAR $\gamma$ -stimulated endothelial  $\bullet$ NO release remain to be defined. Superoxide anion radical (O<sub>2</sub> $\bullet^-$ ) combines with  $\bullet$ NO to decrease  $\bullet$ NO bioavailability. NADPH oxidase, which produces O<sub>2</sub> $\bullet^-$ , and Cu/Zn-superoxide dismutase (Cu/Zn-SOD), which degrades O<sub>2</sub> $\bullet^-$ , thereby contribute to regulation of endothelial cell  $\bullet$ NO metabolism. These findings further elucidate the molecular mechanisms by which PPAR $\gamma$  ligands directly alter vascular endothelial function (159).

CCAAT/enhancer-binding proteins (C/EBPs) upregulate transcription of various inflammatory cytokines and acute phase proteins, such as interleukin (IL)-1 $\beta$ , IL-6, TNF $\alpha$ , and cyclooxygenase-2. Recent studies have demonstrated that PPAR $\gamma$  is present in atherosclerotic lesions, and negatively regulates

expression of these genes. Interestingly, PPAR $\gamma$  gene promoter has tandem repeats of C/EBP-binding motif, and C/EBP- $\delta$  plays a pivotal role in transactivation of PPAR $\gamma$  gene. Recent findings strongly suggest that C/EBP- $\delta$  is negatively autoregulated via transactivation of PPAR $\gamma$ . This feedback mechanism probably downregulates transcription of inflammatory cytokines and acute phase proteins, and modulates inflammatory responses in the early process of atherosclerosis (160).

PPAR $\delta$  controls the inflammatory status of the macrophage. Deletion of PPAR $\delta$  from foam cells increased the availability of inflammatory suppressors, which in turn reduced atherosclerotic lesion area by more than 50%. Lee et al propose an unconventional ligand-dependent transcriptional pathway in which PPAR $\delta$  controls an inflammatory switch through its association and disassociation with transcriptional repressors. PPAR $\delta$  and its ligands may thus serve as therapeutic targets to attenuate inflammation and slow the progression of atherosclerosis (161).

PPAR $\delta$ , but not PPAR $\gamma$ , is the major nuclear VLDL sensor in the macrophage, which is a crucial component of the atherosclerotic lesion. In addition to  $\beta$ -oxidation and energy dissipation, activation of PPAR $\beta$  by VLDL particles induces key genes involved in carnitine biosynthesis and lipid mobilization mediated by a recently identified TG lipase, transport secretion protein 2 (also named desnutrin, iPLA2 $\zeta$ , and adipose TG lipase), resulting in increased fatty acid catabolism. Unexpectedly, deletion of PPAR $\delta$  results in derepression of target gene expression, a phenotype similar to that of ligand activation, suggesting that unliganded PPAR $\delta$  suppresses fatty acid utilization through active repression, which is reversed upon ligand binding. This unique transcriptional mechanism assures a tight control of the homeostasis of VLDL-derived fatty acid and provides a therapeutic target for other lipid-related disorders, including dyslipidemia and diabetes, in addition to coronary artery disease (162).

Ligand activation of PPAR $\delta$  in ECs has a potent antiinflammatory effect, probably via a binary mechanism involving the induction of antioxidative genes and the release of nuclear corepressors. PPAR $\delta$  agonists may have a potential for treating inflammatory diseases such as atherosclerosis and diabetes (163).

Administration of GW501516 to mice upregulated TGF- $\beta$ 1, whereas the expression of proinflammatory genes including MCP-1 was significantly attenuated in the thoracic aorta. Taken together, these results demonstrate the presence of a novel TGF- $\beta$ 1

mediated pathway in the antiinflammatory activities of PPAR $\delta$  (164).

#### PPARs AND HYPERTENSION

The quest for better understanding for the pathophysiological basis of hypertension and atherosclerosis is ongoing. The complexity of hypertension and atherosclerosis and of the underlying mechanisms is becoming increasingly apparent (165).

Tordjman *et al* (166) explore the role of the candidate gene, PPAR $\alpha$ , in the regulation of blood pressure and atherogenesis. PPAR $\alpha$  is believed to impart direct protection in the vessel wall by intervening at essentially every level of the atherogenic process: inflammation, monocyte recruitment and adhesion, cholesterol transport, plaque formation, and thrombosis, mostly through downregulation of NF- $\kappa$ B and AP-1 (167,168,169). In contrast to the plethora of data regarding the atherogenic process, the role of PPAR $\alpha$  in the regulation of vascular tone and blood pressure is still unclear. Several studies conducted in different rat models of hypertension yielded inconsistent results (170,171).

Evidence that PPAR $\alpha$  regulates the expression of the RAS starting with its rate-limiting step, renin. Based on the findings of the current study and on some published data, we suggest that, in the setting of an activated RAS and high endogenous levels of Ang II, PPAR $\alpha$  activation could be detrimental. In this context, PPAR $\alpha$  activation is likely to promote even higher levels of Ang II, increase oxidative stress, raise blood pressure, and ultimately hasten atherosclerosis. Further investigations aimed at delineating the relationships between the transcription factor PPAR $\alpha$  and the components of the RAS and their respective molecular effectors are needed (166).

Evidence demonstrate that short-term clofibrate treatment modulates vasomotor tone, producing effects that are consistent with antagonism of AT1-mediated but promotion of AT2-mediated renal and systemic hemodynamic effects of angiotensin II. These effects appear to involve nitric oxide production and are accompanied by concomitant changes in the expression of AT receptors – increased expression of AT2 receptors but diminished expression of AT1 receptors. PPAR $\alpha$  ligands also protect against pathological damage especially resulting from angiotensin II hypertension (172).

Endothelin-1 (ET-1) is a vasoactive peptide that causes vasoconstriction and vasodilation by binding to ETA receptors on vascular smooth muscle cells

and ETB receptors on endothelial cells, respectively (173,174,175).

The role of cytochrome P450 (CYP450) hydroxylase metabolites in the maintenance of arterial pressure is complex because of their contrasting hypertensive and antihypertensive properties (176). Clofibrate, a PPAR $\alpha$  agonist, has been shown to reduce arterial pressure in Dahl salt-sensitive rats on high salt diet by inducing the genes that code for CYP4504A (CYP4A) enzymes in the renal cortex (177,178). CYP4A is the enzyme responsible for the synthesis of 20-HETE. Interestingly, 20-HETE has actions to reduce sodium transport and, like ETB receptor activation, has been implicated in salt-dependent hypertension (179,180,181,182,183). Therefore, chronic PPAR $\alpha$  agonist treatment reduces salt-dependent hypertension produced by ETB receptor blockade in male and female Sprague-Dawley rats. This suggests a possible relationship between ETB receptor activation and the maintenance of CYP4A protein expression in the kidney of rats fed a high-salt diet (184).

Docosahexaenoic acid (DHA), PPAR $\alpha$  activator, reduces blood pressure in some hypertensive models by unclear mechanisms. PPAR $\alpha$  activator DHA attenuated the development of hypertension, corrected structural abnormalities, and improved endothelial dysfunction induced by Ang II. These effects are associated with decreased oxidative stress and inflammation in the vascular wall (180).

In the absence of dexamethasone, fenofibrate lowered fasting TG and cholesterol but unexpectedly increased systolic blood pressure by ambulatory monitoring. These data suggest that PPAR $\alpha$  activation in humans does not correct insulin resistance induced by glucocorticoids and may adversely affect blood pressure (185).

The current study by Tordjman *et al* is of prime importance, because the investigators have successfully focused our attention on a controversy that involves PPAR $\alpha$ . This central molecule which action had been considered until now beneficial and a prime therapeutic target, may in fact turn out to be a candidate gene for hypertension and for atherosclerosis and, thus, a foe to human health. More in-depth research is required to establish if, when, and how PPAR $\alpha$  might indeed be involved in the generation of high blood pressure and atherosclerosis in humans, issues that remain, at present, unresolved (165).

PPAR $\gamma$  is an attractive molecule given its role as a fatty acid sensor and the increased incidence of hypertension in obese patients. It remains possible that

PPAR $\gamma$  may represent a link among obesity, metabolic dysfunction, and activity of either the circulating or tissue renin-angiotensin system in hypertension. There is no doubt that the mechanisms regulating the renin-angiotensin system by PPAR $\gamma$  will prove to be quite complex given the positive stimulation on renin transcription reported in this issue, along with the negative impact on the AT1 receptor reported previously. That TZDs are generally thought to modestly lower blood pressure suggests that a delicate balance must exist between the effects of PPAR $\gamma$  on the renin-angiotensin system (renin versus AT1 receptor) and other vasoconstrictors, such as ET-1 (186).

Among the transcription factor binding sites in the enhancer is the hormone response element (HRE). Several members of the nuclear hormone receptor superfamily, including retinoic acid receptor and RXR, have been shown to bind to the HRE and to regulate renin gene expression (187). In addition, vitamin D has been reported to negatively modulate renin gene expression through a vitamin D receptor–dependent mechanism, which may involve the HRE (187,188). Because the HRE is homologous to a PPRE, and PPAR $\gamma$ , retinoic acid receptor- $\alpha$ , RXR $\alpha$ , and vitamin D receptor are all members of the same subfamily of ligand-activated transcription factors, it should not be surprising that PPAR $\gamma$  may have to be included among those transcription factors thought to regulate renin expression.

PPAR $\gamma$  activation by TZDs causes a down-regulation of AT1 receptor gene expression via a PPAR $\gamma$ -dependent mechanism in vascular smooth muscle cells. Therefore, PPAR $\gamma$  may play a role in the regulation of Ang II action (189).

#### PPARs AND THE METABOLIC SYNDROME

Metabolic syndrome (MetS) is a cluster of metabolic abnormalities, which is characterized by abdominal obesity, insulin resistance, dyslipidemia, elevated blood pressure, and a proinflammatory and prothrombotic milieu (200,201).

MetS appears to affect a significant proportion of the population. While up to 80% of the almost 200 million adults worldwide with diabetes will die of CVD, people with MetS are also at increased risk, being twice as likely to die from and three times as likely to have a heart attack or stroke compared to people without the syndrome (192). Subjects with MetS have a five – fold greater risk of developing T2D if not already present (193).

Whether or not it is accepted that MetS is a specific disease entity or just a constellation of symptoms, the prevalence of this condition is increasing worldwide and patients need to tackle these risk factor through either lifestyle or pharmacological approaches, in order to reduce the odds of developing diabetes and cardiovascular disease (CVD) (194).

PPARs are intimately involved in nutrient sensing and the regulation of carbohydrate and lipid metabolism. PPAR $\alpha$  and PPAR $\delta$  appear primarily to stimulate oxidative lipid metabolism, while PPAR $\gamma$  is principally involved in the cellular assimilation of lipids via anabolic pathways. These may nevertheless have much greater significance for the public health burden in the abnormal lipid and carbohydrate metabolism of the MetS (195).

One of the hallmarks of MetS is depressed levels of HDLC, which occurs in 37% of patients with MetS (194). Whereas HDLC is not the primary target of lipid-modulating therapy, it is recognized as an important secondary target of therapy, and, thus, treatments that raise HDLC may be important in reducing CVD risk (197).

The identification of PPARs as molecular targets for drugs to treat hypertriglyceridemia and type 2 diabetes mellitus has fueled interest in their biology and potential as targets to treat the metabolic syndrome (198). In keeping with their roles as lipid sensors, ligand-activated PPARs turn on feed-forward metabolic cascades to regulate lipid homeostasis via the transcription of genes involved in lipid metabolism, storage, and transport. Additionally, PPARs may suppress inflammation through mechanisms involving the release of antiinflammatory factors or the stabilization of repressive complexes at inflammatory gene promoters (161,199).

The pathophysiology underlying the MetS is incompletely understood, but insulin resistance appears to be an important component (199,200). Insulin resistance is marked by hyperinsulinemia, enhanced hepatic gluconeogenesis, and impaired insulin-stimulated glucose uptake into skeletal muscle and fat. Elevated levels of circulating FFAs, associated with obesity and insulin resistance, increase fat accumulation in insulin target tissues and contribute to defective insulin action. Obese adipose tissue–derived inflammation and altered adipokine secretion may also inhibit insulin signals and affect systemic metabolism (201). The resulting hyperglycemia, dyslipidemia, hypertension, and inflammation of the MetS cause endothelial dysfunction and hasten atherogenesis (198).

Hypertriglyceridemia and abdominal obesity are key components of the MetS. They may result from an inability of adipose tissue to sequester fatty acids appropriately for storage (202). Instead, fatty acids are deposited as ectopic fat in skeletal muscle (203), liver (204), and other organs (205). It is thought that such fat accumulation is linked to impaired metabolic function of the tissue (206,207). Obesity-related insulin resistance involves the release of mediators, such as FFAs, TNF $\alpha$ , or resistin from adipocytes and decreased production of adiponectin, all of which impair insulin action in skeletal muscle (207).

Adipose tissue as an endocrine organ also releases proinflammatory mediators that promote vascular damage and atherosclerosis. TNF $\alpha$  inhibits insulin signaling contributing to insulin resistance and activates multiple mechanisms of inflammation via NF- $\kappa$ B (208). Leptin can alter insulin action and has recently been recognized to be an important mediator of obesity-related hypertension (209). Angiotensinogen, the precursor of angiotensin II, a key mediator of vascular injury, can be produced and secreted by adipose tissue (210). Plasminogen activator inhibitor 1 (PAI-1) is typically increased in the obesity/insulin-resistance state and plays an important role in atherothrombosis (211,212). In contrast, excessive visceral adipose tissue has been shown to be associated with decreased adiponectin levels (111), an important hormone that exerts antidiabetic (109,110) and antiatherogenic functions (213,214). Adiponectin activates AMP-activated protein kinase, which promotes skeletal muscle glucose uptake and suppresses hepatic glucose production (110). Importantly, adiponectin also inhibits NF- $\kappa$ B activation, thus, attenuating inflammation (215). Visfatin, a growth factor with insulin mimetic action, was recently cloned from fat (216). Unlike adiponectin, plasma levels of visfatin increase in parallel with visceral fat in both mice and humans (216), so the role of visfatin in insulin resistance needs additional investigation. Taken together, these observations suggest that the adipocyte is an integral coordinator of the relationship among obesity, diabetes, and CVD (207).

Interestingly, it is possible that the intra-myocellular lipid (IMCL) impact on insulin sensitivity might be modulated by the oxidative disposal of long-chain fatty acyl-CoAs, depending on the efficiency of the regulation of its flux and metabolism within the mitochondrion. These latest findings represent a fascinating and promising novel understanding linking small abnormalities of tissue energy metabolism to a

steady tendency toward both weight gain and ectopic fat accumulation – particularly in skeletal muscle – as a common denominator inducing both obesity and insulin resistance (203).

The PPAR $\delta$  agonist GW501516 attenuated multiple metabolic abnormalities normally associated with the MetS in humans, and this was probably due to an increase in skeletal muscle fatty acid oxidation. Presently, the individual components of the MetS are treated separately; i.e., statins are used for elevated cholesterol, fibrates are used to reduce TG, and metformin and thiazolidinediones are used for hyperglycemia. The wide range of beneficial effects suggested by the response to GW501516 calls for a larger study in patients to evaluate the clinical efficacy of PPAR $\delta$  agonists for the treatment of hyperlipidemia, liver fat accumulation, obesity, and insulin resistance (217). Thus, PPAR $\delta$  is pivotal to control the program for fatty acid oxidation in the skeletal muscle, thereby ameliorating obesity and insulin resistance through its activation in obese animals (128).

Accelerated atherosclerosis is a major cause of morbidity and death in insulin-resistant states such as obesity and the MetS, macrophages from obese (ob/ob) mice have increased binding and uptake of oxidized LDL, in part due to a posttranscriptional increase in CD36 protein. Macrophages from ob/ob mice are also insulin resistant, as shown by reduced expression and signaling of insulin receptors. Defective macrophage insulin signaling predisposes to foam cell formation and atherosclerosis in insulin-resistant states and that this is reversed in vivo by treatment with PPAR $\gamma$  activators (218).

Obesity and insulin resistance, the cardinal features of MetS, are closely associated with a state of low-grade inflammation (219,220). In adipose tissue chronic overnutrition leads to macrophage infiltration, resulting in local inflammation that potentiates insulin resistance (220,222).

PPAR $\gamma$  is required for maturation of alternatively activated macrophages. Disruption of PPAR $\gamma$ sw in myeloid cells impairs alternative macrophage activation, and predisposes these animals to development of diet-induced obesity, insulin resistance, and glucose intolerance. Furthermore, gene expression profiling revealed that down regulation of oxidative phosphorylation gene expression in skeletal muscle and liver leads to decreased insulin sensitivity in these tissues. These suggest that resident alternatively activated macrophages have a beneficial role in regulating nutrient homeostasis and suggest

that macrophage polarization towards the alternative state might be a useful strategy for treating T2D (223).

#### PPARs AS THERAPEUTICS TARGETS FOR MetS

Because these nuclear receptors are activated by extracellular signals and control multiple gene targets, PPARs can be seen as nodes that control multiple inputs and outputs involved in energy balance, providing insight into how metabolism and the vasculature may be integrated. The ongoing clinical use of fibrates, which activate PPAR $\alpha$ , and thiazolidinediones, which activate PPAR $\gamma$ , establishes these receptors as viable drug targets, whereas considerable *in vitro* animal model and human surrogate marker studies suggest that PPAR activation may limit inflammation and atherosclerosis. Together, these various observations have stimulated intense interest in PPARs as therapeutic targets and led to large-scale cardiovascular end-point trials with PPAR agonists (224).

The current clinical approach to MetS is to focus on appropriate management of accompanying risk factors. While priority should be given to management of underlying risk factors with therapeutic lifestyle changes, associated major risk factors should be treated according to evidence-based medicine goals and principles, and appropriate clinical attention should be given to the presence of emerging risk factors. Clearly, there are multiple targets for therapy to reduce the high risk of the MetS. While no single treatment for the MetS as a whole yet exist, it is well established that lifestyle changes, for example, changes in diet and increased physical exercise, form the first-line strategy of intervention (194).

In MetS, a condition of impaired fatty acid metabolism in adipose tissue generally results in the increased release of free fatty acids into the circulation. This in turn leads to multiple abnormalities in the circulating lipoprotein profile including low HDL, high TG, and VLDL remnants, with average LDL. The triad of elevated TG, reduced HDL and small dense LDL, along with concomitant increased in TG-rich remnant particles, comprises the atherogenic dyslipidemia of MetS. MetS, with or without progression to T2D, is therefore a major atherogenic factor. CHD risk reduction in MetS requires not only aggressive LDL-C lowering but also management of each aspect of dyslipidemia, including lowering TG levels, increasing HDL-C levels, and increasing the size of the average LDL-C particle. Based on available evidence, LDL-C is the primary target of therapy in MetS (223). In

patients with atherogenic dyslipidemia with elevated TG, non-HDL-C represents a secondary target of treatment after the LDL-C goal is achieved. The tertiary goal in these patients is to raise HDL-C when it is reduced after attaining goals for LDL-C and non-HDL-C (194).

PPAR activation can improve metabolic parameters like glucose and lipid levels but also alter directly vascular responses by regulating target genes including those encoding adhesion molecules, The ATP-binding cassette transporter 1 (ABCA 1), lipoprotein lipase, cytokines and chemokines (226). PPAR activation has many anti-atherosclerotic effects with distinct and overlapping targets. For instance, lipoprotein lipase acts on circulating lipoproteins to activate PPAR $\alpha$  and initiates PPAR $\alpha$ -dependent positive feedback loop for TG-rich lipoprotein catabolism (44). Exogenous PPAR $\alpha$  agonist such as fenofibrate reduce TNF $\alpha$ -induced activation of NF- $\kappa$ B (227), a transcription factor integrating inflammation and atherosclerosis. There have been indications that PPAR $\alpha$  is involved in the pathogenesis of insulin resistance. In this regard, PPAR $\alpha$  knockout mice are protected from diet-induced insulin resistance probably because of inhibition of PPAR $\alpha$ -dependent fatty acid oxidation (227). By contrast, genetic defects in PPAR $\gamma$  can recapitulate all the salient features of MetS in humans (96). Therefore, PPARs are likely to be involved in the development of MetS and, accordingly, can serve as potential therapeutic targets for the prevention and treatment of MetS (228).

Activation of PPAR $\alpha$  by fibrates stimulates the oxidation of free fatty acids in the liver (229), diverting them away from TG synthesis and thus reducing the hepatic synthesis of TG-rich lipoproteins (230). The activation of PPAR $\alpha$  also induces expression of the gene for lipoprotein lipase (LPL), the enzyme responsible for hydrolyzing TG and phospholipids in TG-rich lipoproteins in plasma (231). Thus, activators of PPAR $\alpha$  lower the concentration of plasma TG by reducing its rate of synthesis and increasing its rate of hydrolysis. An additional effect of activating PPAR $\alpha$  is the inhibition of synthesis of apolipoprotein (apo) C-III (232). Because apoC-III delays the catabolism of TG-rich lipoproteins, its inhibition provides a further mechanism by which PPAR $\alpha$ -activators such as fibrates lower the concentration of plasma TG. Fibrates also impact on the concentration of HDLC in humans, although the precise mechanisms are not known. Fibrates increase expression of the genes for both apo A-I and apo A-II, the 2 main apolipoproteins of HDL

## Possible anti-atherogenic mechanisms of fibrates




Fig. 4. Effects of fibrates with the potential to protect against cardiovascular disease (Adapted from Barter PJ, 2008).

(233,234), although the magnitude of the increases in the plasma concentrations of these proteins tends to be small. Generally, the increase in plasma apoA-II is greater than that of apoA-I (234,235,236). This results in an increase in the concentration of HDL particles containing both apoA-I and apoA-II but a decrease in those containing apoA-I without apoA-II (236). Other potential mechanisms by which fibrates increase the level of HDLC include an enhancement of cell cholesterol efflux secondary to an induction of cell ABCA1 expression (237), although others have not been able to confirm this (238). Fenofibrate has been reported to decrease SR-B1 in the liver (239), an action that would contribute to an elevation of HDLC but which may not necessarily translate into a reduction in atherosclerosis. Fibrates also increase LDL particle size (142,240) and decrease the concentration of TG-rich lipoprotein remnants. These latter effects are potentially antiatherogenic and appear to be mediated equally well by all fibrates. Post hoc analyses of several of the fibrate studies have shown that people with features of the MetS, particularly overweight people with high plasma TG levels and low levels of HDL cholesterol, derive a disproportionately large reduction in cardiovascular events when treated with these agents. Thus, there is a strong case for the use of a fibrate to reduce the cardiovascular risk in overweight people with high TG and low HDLC (241).

PPAR $\gamma$  is essential for normal adipocyte differentiation and proliferation as well as fatty acid uptake and storage. Thiazolidinediones increase the

number of small adipocytes and the subcutaneous adipose-tissue mass in studies in animal models (100,242,243). These observations, plus the high level of PPAR $\gamma$  expression in adipose tissue, have led to the hypothesis that thiazolidinediones exert their insulin-sensitizing actions either directly (the “fatty acid steal” hypothesis) or indirectly, by means of altered adipokine release, modulating insulin sensitivity outside adipose tissue. According to the “fatty acid steal” hypothesis, thiazolidinediones promote fatty acid uptake and storage in adipose tissue. In this way, they increase adipose-tissue mass and spare other insulin-sensitive tissues such as skeletal muscle and the liver, and possibly pancreatic beta cells, from the harmful metabolic effects of high concentrations of free fatty acids. Thiazolidinediones thus keep fat where it belongs.

Various adipokines, such as adiponectin (244,245), TNF $\alpha$  (246), resistin (247), and 11  $\beta$ -hydroxysteroid dehydrogenase 1, the enzyme that produces cortisol locally in adipose tissue (242,248), are among the genes that are regulated by PPAR $\gamma$  agonists in rodents. Of these, adiponectin increases insulin sensitivity, and TNF $\alpha$ , resistin, and 11  $\beta$ -hydroxysteroid dehydrogenase 1 (249) induce insulin resistance in rodents. Adiponectin, an adipocytokine produced exclusively by adipose tissue, has both insulin-sensitizing and antiatherogenic properties in mice (244,245). PPAR $\gamma$  agonists increase adiponectin expression in vitro in adipose tissue (250).

In the liver and in adipose tissue,  $11\beta$ -hydroxysteroid dehydrogenase 1 catalyzes the interconversion of cortisone to cortisol (251). A full-blown MetS characterized by obesity and the accumulation of visceral fat, as well as increased concentrations of cortisol in the portal vein but not of systemic cortisol, develops in mice that overexpress  $11\beta$ -hydroxysteroid dehydrogenase 1 in adipose tissue (249). Thiazolidinediones down-regulate  $11\beta$ -hydroxysteroid dehydrogenase 1 expression in adipose tissue (250) and might thereby alleviate features of the MetS.

Study of Samaha et al (252) found overall favorable effects of rosiglitazone on adipokines, including a significant increase in adiponectin levels. This increase in adiponectin may be attributable to a direct effect of rosiglitazone on adipocytes, and possibly macrophages (253). Adiponectin has been shown to play a role in modulating insulin sensitivity and to be increased by rosiglitazone in patients with diabetes (254). Furthermore, increasing quintiles of adiponectin levels have been associated with decreased risk of myocardial infarction (255).

Study also demonstrates that rosiglitazone lowers resistin levels in patients with MetS, as recently demonstrated in one small study of 14 patients with T2D (256). In humans, however, resistin is predominantly produced by macrophages in response to inflammatory stimuli and is almost undetectable in adipose tissue. Rosiglitazone may thus have a direct effect on resistin expression, such as through macrophage  $PPAR\gamma$  activation. We have recently shown that resistin levels independently correlate with degree of coronary artery calcification (257).

Rosiglitazone also significantly reduced levels of CRP, IL-6, and sTNF $\alpha$  R2. (258). Similar responses for CRP and IL-6 were previously found in patients with diabetes (259). A significant correlation between the rosiglitazone-induced changes in resistin and changes in the inflammatory markers IL-6 and sTNF $\alpha$  R<sup>2</sup>. These findings are consistent with our previous findings of a significant correlation between baseline levels of resistin and sTNF $\alpha$  R<sup>2</sup>. (258). Rosiglitazone has been shown to lower CRP in nondiabetic patients with coronary artery disease (260) and in nonobese Taiwanese patients with MetS (261).

These findings suggest that rosiglitazone, presumably through its  $PPAR\gamma$  agonist properties, has direct effects on inflammatory markers and adipokines in the absence of favorable lipid effects. These findings may help explain the mechanism underlying the possible antiatherosclerotic effects of rosiglitazone (252).

Current pharmacologic approaches are unsatisfactory in improving such consequences of insulin resistance as hyperglycemia, diabetic dyslipidemia, abnormal coagulation and fibrinolysis, and hypertension, each of which may require the use of at least one medication. Thus, the development of drugs targeted to reverse insulin resistance is important. The insulin-sensitizing thiazolidinediones, which are selective ligands of the  $PPAR\gamma$ , are the first drugs to address the basic problem of insulin resistance in patients with MetS and T2D (262).

It is noteworthy that TZDs show intracellular antioxidant activity. This property may reflect 'preventative' action since these agents do not show direct antioxidant scavenging activity on free radicals, but block several mechanisms that in hyperglycaemic or hyperlipidaemic conditions lead to the generation of oxidative stress. It has been observed that  $PPAR\gamma$  ligands inhibit the expression of inducible NO synthase (iNOS) and, consequently, ONOO<sup>-</sup> production, in mesangial cells and in cerebellar granule cells (133).

The antagonising activity of NF- $\kappa$ B is also of relevance in this setting, since it is through this mechanism that rosiglitazone and pioglitazone can reduce iNOS over-expression, and the related over-production of nitrotyrosine during experimental myocardial infarction (263).

Such effects may explain the ability of pioglitazone to reduce oxidative stress in type 2 diabetic patients after a meal (263). It is not surprising therefore that pioglitazone can improve endothelial dysfunction in cardiovascular disease patients, both with and without T2D and MetS (265,266).

$PPAR\gamma$  activation by the TZDs results in an array of effects on traditional and non-traditional cardiovascular risk factors that are independent of their effects on glycaemic control. These include reduction of IMT progression and circulating platelet activity, attenuation of PAI-1 expression, inhibition of glycation, increase in plasma adiponectin and reduction of CRP, IL-6 and MMP-9 levels. There is evidence that many of the anti-atherogenic and anti-inflammatory effects of the TZDs may be attributable to the glitazone class, although no head-to-head studies have been carried out to establish comparative efficacy (267).

There continues to be uncertainty about the risk of ischemic heart disease in patients with T2D associated with the use of rosiglitazone (268). There was no evidence of any increase in death from either cardiovascular causes or all causes. Rosiglitazone was associated with an increased risk of heart failure.




Fig. 5. TZDs affect oxidative stress and inflammation at multiple levels (adapted from Ceriello A, 2008).

The data were insufficient to determine whether the drug was associated with an increase in the risk of myocardial infarction (269). Caution should be used in patients with underlying heart disease using nitrates, and when added to ongoing insulin-based therapy (268).

A hallmark of the MetS is dyslipidemia, marked by elevated TG and low levels of HDLC. HDL is a driving force in the process of reverse cholesterol transport, reclaiming excess peripheral tissue cholesterol to the liver for excretion. Accordingly, low levels of HDL are associated with an increased risk of coronary artery disease and cardiovascular death in afflicted patients, while overexpression of apoA-I, the major apolipoprotein composing HDL particles, retards atherogenesis in animal models (270,271,272). Despite clear therapeutic need, currently marketed cholesterol-modifying drugs raise serum HDL levels only modestly.

High-affinity PPAR $\delta$  ligands have revealed an important role for PPAR $\delta$  in lipoprotein metabolism. Treatment of insulin-resistant obese rhesus monkeys with the PPAR $\delta$ -selective agonist GW501516 resulted in a dramatic 79% increase in HDLC, a 56% decrease in TG, and a 29% decrease in LDLC (66). The profound increase in HDLC levels correlated with an increase in number, not size, of HDL particles and was accompanied by increased serum levels of the HDL-associated apolipoproteins apoA-I, apoA-II, and apoC-III. In addition, fasting insulin levels declined by up to 48% in the PPAR $\delta$  drug-treated animals (66). Obese and nonobese mice similarly develop an increase of up to 50% in HDLC levels when treated with PPAR $\delta$  agonists (273,274). The mechanism by which PPAR $\delta$  activation

raises HDLC levels remains to be elucidated, but studies to date indicate that expression of the reverse cholesterol transporter ABCA1 is enhanced in some tissues upon exposure to PPAR $\delta$  agonists, including human and mouse macrophages as well as human intestinal cells and fibroblasts (66,273). Additional work suggests that PPAR $\delta$  activation reduces intestinal cholesterol absorption via downregulation of the Niemann-Pick C1-like 1 gene (NPC1L1) (273). NPC1L1 is a key mediator of intestinal cholesterol absorption and a putative target for the clinically used cholesterol absorption inhibitor ezetimibe. In light of these findings, PPAR $\delta$  drugs are now in clinical trial for the treatment of human dyslipidemia.

Once viewed as a bland storage depot, adipose tissue has emerged as a dynamic endocrine organ (201). Adiposity correlates with insulin resistance and is believed by some to be of primacy in the MetS (275). Even mild weight loss may improve serum lipid profiles, glycemic control, and hypertension, yet currently available weight-loss drugs are of limited effectiveness (276,277). Genetic models and ligand treatment studies have uncovered powerful regulatory functions for PPAR $\delta$  in adipose tissue metabolism and weight control. Transcriptional analysis of brown fat in fat-specific VP16-PPAR $\delta$  mice revealed upregulation of genes involved in TG hydrolysis (hormone-sensitive lipase), fatty acid oxidation (long-chain acyl-CoA synthetase, very-long-chain acyl-CoA synthetase, acyl-CoA oxidase), and uncoupling of oxidative phosphorylation (UCP1 and UCP3). UCP1 expression was likewise elevated in white adipose tissue. Conversely, PPAR $\delta$ -null mice are more susceptible to weight gain and have blunted expression of brown




Fig. 6. Therapeutic targets of PPAR $\delta$  in the MetS (Adapted from Barish GD, 2006).

fat UCP1 on a high-fat diet. These genetic models collectively suggest that activation of PPAR $\delta$  protects against obesity (164).

Administration of the synthetic PPAR $\delta$  agonist GW501516 to genetically obese (db/db) mice reduced intracellular TG accumulation in the brown fat and liver, analogous to the effects of VP16-PPAR $\delta$ . Moreover, PPAR $\delta$  agonists enhanced  $\beta$ -oxidation in 3T3-L1 preadipocytes by 50% (164). Most importantly, PPAR $\beta$  ligands retard weight gain in models of high-fat diet-induced obesity (128,164). These results suggest that PPAR $\beta$  synthetic drugs may be therapeutic as antiobesity agents.

PPAR $\beta$  has emerged as a powerful metabolic regulator in diverse tissues including fat, skeletal muscle, and the heart. Its transcriptional program enhances fatty acid catabolism and energy uncoupling, resulting in decreased TG stores, improved endurance performance, and enhanced cardiac contractility, respectively. PPAR $\delta$  receptor activation mitigates macrophage inflammatory responses and modulates lipoprotein metabolism to lower TG and robustly raise HDL-C. Additionally, recent studies reveal that PPAR $\delta$  activation in the liver suppresses hepatic glucose output, contributing to improved glucose homeostasis (278). These aggregate effects suggest that high-affinity PPAR $\delta$  synthetic drugs may uniquely target multiple components of the MetS, including obesity, insulin resistance, hyperglycemia, dyslipidemia, and atherosclerosis, or other diseases such as cardiomyopathy (Figure 6) (198).

Research programs to develop agonists that combine the therapeutic effects of both PPAR $\alpha$  and

PPAR $\gamma$  selective agonists, creating the expectation of synergy in the treatment of T2D and the MetS, have therefore been undertaken. Among these dual PPAR $\alpha$ / $\gamma$  agonists, compounds belonging to the glitazar class have been advanced to clinical development (Phases II and III). These trials clearly demonstrate superior efficacy of these compounds in correcting abnormalities associated with T2D and the MS, improving both lipid and glucose homeostasis to a greater extent than did selective PPAR agonists. However, safety remains a critical issue for glitazars, as the development of several promising candidates was halted because of adverse toxicity profiles (279).

The discovery and preliminary biological and pharmacokinetic properties of muraglitazar (BMS-298585), a novel oxybenzylglycine dual ( $\alpha$ / $\gamma$ ) PPAR activator, have been recently described. Muraglitazar binds with high affinity to both human PPAR $\gamma$  and PPAR $\alpha$  ligand binding domain protein (IC<sub>50</sub> for binding = 0.19 and 0.25  $\mu$ mol/l, respectively) and potently transactivates full-length human PPAR $\gamma$ - or PPAR $\alpha$ -mediated reporter gene activity (EC<sub>50</sub> for transactivation = 0.11 and 0.32  $\mu$ mol/l, respectively). Harrity et al (280) assessed the effects of muraglitazar treatment on diabetes and other metabolic abnormalities in genetically obese, diabetic, and hyperlipidemic db/db mice.

Tesaglitazar, a Novel Dual PPAR  $\alpha$ / $\gamma$  agonist, dose – dependently improves the metabolic abnormalities associated with insulin resistance in a non – diabetic population (281). Macelignan, a natural compound isolated from *Myristica fragrans*, enhanced insulin sensitivity and improved lipid metabolic disorders

by activating PPAR $\alpha$ / $\gamma$  and attenuating ER stress, suggesting that it has potential as an antidiabetes agent for the treatment of T2D and MetS (282).

However, current safety standards with compounds for T2D and the MetS are high, and few adverse effects are currently tolerated. We believe that the best solution would be to screen for compounds that combine an intermediate PPAR $\alpha$  affinity with selective PPAR $\gamma$  - modulating activity (279).

## Conclusion

The MetS is diagnosed by a cluster of clinical parameters including central obesity, atherogenic dyslipidemia, raised blood pressure and hyperglycemia. Visceral obesity, hepatic steatosis and insulin resistance have been proposed as unifying mechanisms, yielding a prothrombotic and proinflammatory state. Consequently, patients with MetS are at increased risk of CAD and T2D.

The discovery of the crucial role of peroxisome proliferator-activated receptors (PPARs) as regulators of lipid and glucose metabolism has raised interest in the development of synthetic ligands as potential tools for therapeutic intervention in T2D and the MetS.

## References:

1. Grundy SM, Brewer HB Jr, Cleeman JI, Smith SC Jr, Lenfant C. Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation* 2004; 109: 433 – 438.
2. Lakka HM, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, et al. The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* 2002; 288: 2709 – 2216.
3. Shulman AI, Mangelsdorf DJ. Retinoid X Receptor Heterodimers in the Metabolic Syndrome. *N Engl J Med* 2005; 353: 604 – 615.
4. Gustafson B, Hammarstedt A, Andersson CX, Smith U. Inflamed Adipose Tissue. A Culprit Underlying the Metabolic Syndrome and Atherosclerosis. *Arterioscler Thromb Vasc Biol* 2007; 27: 2276 – 2283.
5. Sonnenberg G, Krakower R, Kissebah AH. A Novel Pathway to the Manifestations of Metabolic Syndrome. *Obes Res* 2004; 12: 180 – 186.
6. Taskinen MR. Is Metabolic Syndrome the Main Threat to Human health in the Twenty – First Century? *Arterioscler Thromb Vasc Biol* 2007; 27: 2275.
7. Morales LA, Piqueras L, Bishop-Bailey D. Peroxisome proliferator-activated receptors and inflammation. *Pharmacol Ther* 2006; 110: 371 – 385.
8. Bishop-Bailey D, Wray J. Peroxisome proliferator-activated receptors: a critical review on endogenous pathways for ligand generation. *Prostaglandins Other Lipid Mediat* 2003; 71: 1 – 22.
9. Barak Y, Liao D, He W, Ong ES, Nelson MC, Olefsky JM, et al. Effects of peroxisome proliferator-activated receptor delta on placentation, adiposity, and colorectal cancer. *Proc Natl Acad Sci U S A* 2002; 99: 303 – 308.
10. Michalik L, Desvergne B, Tan NS, Basu-Modak S, Escher P, Rieusset J, et al. Impaired skin wound healing in peroxisome proliferator-activated receptor (PPAR)alpha and PPARbeta mutant mice. *J Cell Biol* 2001; 154: 799 – 814.
11. Michalik L, Auwerx J, Berger JP, Chatterjee VK, Glass CK, Gonzalez FJ, et al. International Union of Pharmacology. LXI. Peroxisome proliferator-activated receptors. *Pharmacol Rev* 2006; 58:726 –741.
12. Bishop –Bailey D. Peroxisome Proliferator – Activated Receptor  $\beta/\gamma$  Goes Vascular. *Circ Res* 2008; 102: 146 – 147.
13. Mudaliar S, Henry RR. PPAR Agonist in Health and Disease: a Pathophysiologic and Clinical Overview. *Curr Opin Endocrinol Metab* 2002; 9: 285 – 302.
14. Issemann I and Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature (Lond)* 1990; 347: 645–650.
15. Dreyer C, Krey G, Keller H, Givel F, Helftenbein G, and Wahli W. Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell* 1992; 68: 879 – 887.
16. Kliewer SA, Forman BM, Blumberg B, Ong ES, Borgmeyer U, Mangelsdorf DJ, Umesonon K, and Evans RM. Differential expression and activation of a family of murine peroxisome proliferator-activated receptors. *Proc Natl Acad Sci USA* 1994; 91: 7355 – 7359.
17. Amri EZ, Bonino F, Ailhaud G, Abumrad NA, and Grimaldi PA. Cloning of a protein that mediates transcriptional effects of fatty acids in preadipocytes. Homology to peroxisome proliferator- activated receptors. *J Biol Chem* 1995; 270: 2367 – 2371.
18. Schmidt A, Endo N, Rutledge SJ, Vogel R, Shinar D, and Rodan GA. Identification of a new member of the steroid hormone receptor superfamily that is activated by a peroxisome proliferator and fatty acids. *Mol Endocrinol* 1992; 6:1634 – 1641.
19. Mandart S, Muller M, and Kersten S. Peroxisome proliferator-activated receptor alpha target genes. *Cell Mol Life Sci* 2004; 61: 393 – 416.
20. Braissant O, Fougelle F, Scotto C, Dauca M, and Wahli W. Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat. *Endocrinology* 1996; 137: 354 – 366.
21. Bastie C, Holst D, Gaillard D, Jehl-Pietri C, and Grimaldi PA. Expression of peroxisome proliferator-activated receptor PPARdelta promotes induction of PPARgamma and adipocyte differentiation in 3T3C2 fibroblasts. *J Biol Chem* 1999; 274:21920 – 21925.
22. Peters JM, Lee SS, Li W, Ward JM, Gavrilova O, Everett C, et al. Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(delta). *Mol Cell Biol* 2000; 20: 5119 – 5128.
23. Chawla A, Schwarz EJ, Dimaculangan DD, and Lazar MA. Peroxisome proliferator-activated receptor (PPAR) gamma: adipose-predominant expression and induction early in adipocyte differentiation. *Endocrinology* 1994; 135: 798 – 800.
24. Tontonoz P, Hu E, Graves RA, Budavari AI, and Spiegelman BM. Mppar gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes Dev* 1994a; 8: 1224 – 1234.
25. Tontonoz P, Graves RA, Budavari AI, Erdjument-Bromage H, Lui M, Hu E, Tempst P, and Spiegelman BM. Adipocyte-specific transcription factor ARF6 is a heterodimeric complex of two nuclear hormone receptors. PPAR gamma and RXR alpha. *Nucleic Acids Res* 1994b; 22: 5628 – 5634.
26. Zhu Y, Qi C, Korenberg JR, Chen XN, Noya D, Rao MS, et al. Structural organization of mouse peroxisome proliferator-activated receptor gamma (mPPAR gamma) gene: alternative promoter use and different splicing yield two mPPAR gamma isoforms. *Proc Natl Acad Sci USA* 1995; 92: 7921 – 7925.
27. Kliewer SA, Umesonon K, Noonan DJ, Heyman RA, and Evans RM. Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature (Lond)* 1992; 358: 771 – 774.

28. Keller H, Dreyer C, Medin J, Mahfoudi A, Ozato K, and Wahli W. Fatty acids and retinoids control lipid metabolism through activation of peroxisome proliferator-activated receptor-retinoid X receptor heterodimers. *Proc Natl Acad Sci USA* 1993; 90: 2160 – 2164.
29. Tugwood JD, Issemann I, Anderson RG, Bundell KR, McPheat WL, and Green S. The mouse peroxisome proliferator activated receptor recognizes a response element in the 5' flanking sequence of the rat acyl CoA oxidase gene. *EMBO (Eur Mol Biol Organ) J* 1992; 11: 433 – 439.
30. Feige JN, Gelman L, Tudor C, Engelborghs Y, Wahli W, and Desvergne B. Fluorescence imaging reveals the nuclear behavior of peroxisome proliferator-activated receptor/retinoid X receptor heterodimers in the absence and presence of ligand. *J Biol Chem* 1995; 280: 17880 – 17890.
31. Di-Poi N, Tan NS, Michalik L, Wahli W, and Desvergne B. Antiapoptotic role of PPARbeta in keratinocytes via transcriptional control of the Akt1 signaling pathway. *Mol Cell* 2002; 10: 721 – 733.
32. DiRenzo J, Soderstrom M, Kurokawa R, Ogliastro MH, Ricote M, Ingrey S, et al. Peroxisome proliferator-activated receptors and retinoic acid receptors differentially control the interactions of retinoid X receptor heterodimers with ligands, coactivators, and corepressors. *Mol Cell Biol* 1997; 17: 2166 – 2176.
33. Ijpenberg AI, Jeannin E, Wahli W, and Desvergne B. Polarity and specific sequence requirements of peroxisome proliferator-activated receptor (PPAR)/retinoid X receptor heterodimer binding to DNA. A functional analysis of the malic enzyme gene PPAR response element. *J Biol Chem* 1997; 272: 20108 – 20117.
34. Juge-Aubry C, Pernin A, Favez T, Burger AG, Wahli W, Meier CA, et al. DNA binding properties of peroxisome proliferator-activated receptor subtypes on various natural peroxisome proliferator response elements. Importance of the 5'-flanking region. *J Biol Chem* 1997; 272: 25252 – 25259.
35. Dowell P, Ishmael JE, Avram D, Peterson VJ, Nevrivy DJ, and Leid M. Identification of nuclear receptor corepressor as a peroxisome proliferator-activated receptor alpha interacting protein. *J Biol Chem* 1999; 274: 15901 – 15907.
36. Stanley TB, Leesnitzer LM, Montana VG, Galardi CM, Lambert MH, Holt JA, et al. Subtype specific effects of peroxisome proliferator-activated receptor ligands on corepressor affinity. *Biochemistry* 2003; 42: 9278 – 9287.
37. Guan HP, Ishizuka T, Chui PC, Lehrke M, and Lazar MA. Corepressors selectively control the transcriptional activity of PPARgamma in adipocytes. *Genes Dev* 2005; 19: 453 – 461.
38. Yu C, Markan K, Temple KA, Deplewski D, Brady MJ, and Cohen RN. The nuclear receptor corepressors NCoR and SMRT decrease peroxisome proliferator-activated receptor gamma transcriptional activity and repress 3T3-L1 adipogenesis. *J Biol Chem* 2005; 280: 13600 – 13605.
39. Bishop-Bailey D and Wray J. Peroxisome proliferator-activated receptors: a critical review on endogenous pathways for ligand generation. *Prostaglandins Other Lipid Mediat* 2003; 71: 1 – 22.
40. Desvergne B, Michalik L, and Wahli W. Be fit or be sick: peroxisome proliferator-activated receptors are down the road. *Mol Endocrinol* 2004; 18: 1321 – 1332.
41. Evans RM, Barish GD, and Wang YX. PPARs and the complex journey to obesity. *Nat Med* 2004; 10: 1 – 7.
42. Ferré. The Biology of Peroxisome Proliferator-Activated Receptors. *Diabetes* 2004; 53 (suppl. 1): S43 – S50.
43. Chawla A, Lee CH, Barak Y, He W, Rosenfeld J, Liao D, et al. PPARdelta is a very low-density lipoprotein sensor in macrophages. *Proc Natl Acad Sci USA* 2003; 100: 1268 – 1273.
44. Ziouzenkova O, Perrey S, Asatryan L, Hwang J, MacNaul KL, Moller DE, et al. Lipolysis of TG-rich lipoproteins generates PPAR ligands: evidence for an antiinflammatory role for lipoprotein lipase. *Proc Natl Acad Sci USA* 2003; 100: 2730 – 2735.
45. Ahmed W, Orasanu G, Nehra V, Asatryan L, Rader DJ, Ziouzenkova O, et al. High-density lipoprotein hydrolysis by endothelial lipase activates PPARalpha: a candidate mechanism for high-density lipoprotein-mediated repression of leukocyte adhesion. *Circ Res* 2006; 98: 490 – 498.
46. Bosse Y, Despres JP, Bouchard C, Perusse L, Vohl MC. The peroxisome proliferator-activated receptor alpha L162V mutation is associated with reduced adiposity. *Obes Res* 2003; 11: 809 – 816.
47. Kliewer SA, Sundseth SS, Jones SA, et al. Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferator-activated receptors alpha and gamma. *Proc Natl Acad Sci U S A* 1997; 94: 4318 – 4323.
48. Kliewer SA, Lenhard JM, Willson TM, Patel I, Morris DC, Lehmann JM. A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 1995; 83: 813 – 819.
49. Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, Evans RM. 15-Deoxydelta 12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma. *Cell* 1995; 83: 803 – 812.
50. Way JM, Harrington WW, Brown KK, Gottschalk WK, Sundseth SS, Mansfield TA, et al. Comprehensive messenger ribonucleic acid profiling reveals that peroxisome proliferator-activated receptor gamma activation has coordinate effects on gene expression in multiple insulin-sensitive tissues. *Endocrinology* 2001; 142: 1269 – 1277.
51. Ribon V, Johnson JH, Camp HS, Saltiel AR. Thiazolidinediones and insulin resistance: peroxisome proliferator-activated receptor gamma activation stimulates expression of the CAP gene. *Proc Natl Acad Sci U S A* 1998; 95: 14751 – 14756.
52. Young PW, Cawthorne MA, Coyle PJ, Holder JC, Holman GD, Kozka IJ, et al. Repeat treatment of obese mice with BRL 49653, a new potent insulin sensitizer, enhances insulin action in white adipocytes: association with increased insulin binding and cell-surface GLUT4 as measured by photoaffinity labeling. *Diabetes* 1995; 44: 1087 – 1092.
53. Sfeir Z, Ibrahim A, Amri E, Grimaldi P, Abumrad N. Regulation of FAT/CD36 gene expression: further evidence in support of a role of the protein in fatty acid binding/transport. *Prostaglandins Leukot Essent Fatty Acids* 1997; 57: 17 – 21.
54. Tontonoz P, Hu E, Spiegelman BM. Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor. *Cell* 1994; 79: 1147-56. [Erratum, *Cell* 1995; 80: following 957.
55. Schoonjans K, Peinado-Onsurbe J, Lefebvre AM, Heyman RA, Briggs M, Deeb S, et al. PPARalpha and PPARgamma activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene. *EMBO J* 1996; 15: 5336 – 5348.
56. Martin G, Schoonjans K, Lefebvre AM, Staels B, Auwerx J. Coordinate regulation of the expression of the fatty acid transport protein and acyl-CoA synthetase genes by PPARalpha and PPARgamma activators. *J Biol Chem* 1997; 272: 28210 – 28217.
57. Aubert J, Champigny O, Saint-Marc P, Negrel L, Collins S, Ricquier D, et al. Up-regulation of UCP-2 gene expression by PPAR agonists in preadipose and adipose cells. *Biochem Biophys Res Commun* 1997; 238: 606 – 11.
58. Guan HP, Li Y, Jensen MV, Newgard CB, Stepan CM, Lazar MA. A futile metabolic cycle activated in adipocytes by antidiabetic agents. *Nat Med* 2002; 8: 1122 - 1128.
59. Nagy L, Tontonoz P, Alvarez JG, Chen H, Evans RM. Oxidized LDL regulates macrophage gene expression through ligand activation of PPARgamma. *Cell* 1998 93: 229 – 240.
60. Tontonoz P, Nagy L, Alvarez JG, Thomazy VA, Evans RM. PPARgamma promotes monocyte/macrophage differentiation and uptake of oxidized LDL. *Cell* 1998; 93: 241 – 252.
61. Chawla A, Boisvert WA, Lee CH, Laffitte BA, Barak Y, Joseph SB, et al. A PPAR gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Mol Cell* 2001; 7: 161 – 171.
62. Li AC, Brown KK, Silvestre MJ, Willson TM, Palinski W, Glass CK. Peroxisome proliferator-activated receptor gamma ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. *J Clin Invest* 2000; 106: 523 – 531.
63. Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM, Kliewer SA. An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 1995; 270: 12953 – 12956.

64. Wang YX, Lee CH, Tiep S, [Yu RT](#), [Ham J](#), [Kang H](#), [Evans RM](#). Peroxisome-proliferator-activated receptor delta activates fat metabolism to prevent obesity. *Cell* 2003; 113: 159 – 170.
65. Muoio DM, MacLean PS, Lang DB, Li S, Houmard JA, Way JM, et al. Fatty acid homeostasis and induction of lipid regulatory genes in skeletal muscles of peroxisome proliferator-activated receptor (PPAR) alpha knock-out mice: evidence for compensatory regulation by PPAR delta. *J Biol Chem* 2002; 277: 26089 – 26097.
66. Oliver WR Jr, Shenk JL, Snaith MR, Russel CS, Plunket KD, Bodkin NL, et al. A selective peroxisome proliferator-activated receptor delta agonist promotes reverse cholesterol transport. *Proc Natl Acad Sci U S A* 2001; 98: 5306 – 5311.
67. Reaven GM. Syndrome X: 6 years later. *J Intern Med* 1994; Suppl 736: 13 – 22.
68. Lee CH, Olson P, Evans RM. Minireview: Lipid Metabolism, Metabolic Disease, and Peroxisome Proliferator-Activated Receptors. *Endocrinology* 2003; 144: 2201 – 2207.
69. Desvergne B, Wahli W. Peroxisome Proliferator-Activated Receptors: Nuclear Control of Metabolism. *Endocrinol Rev* 1999; 20: 649 – 688.
70. Peters JM, Hennuyer N, Staels B, Fruchart JC, Fievet C, Gonzalez FJ, et al. Alterations in lipoprotein metabolism in peroxisome proliferator-activated receptor  $\alpha$ -deficient mice. *J Biol Chem* 1997; 272: 27307 – 27312.
71. Lee SS, Pineau T, Drago J, Lee EJ, Owens JW, Kroetz DL, et al. Targeted disruption of the  $\alpha$  isoform of the peroxisome proliferator-activated receptor gene in mice results in abolishment of the pleiotropic effects of peroxisome proliferators. *Mol Cell Biol* 1995; 15: 3012 – 3022.
72. Aoyama T, Peters JM, Iritani N, Nakajima T, Furihata K, Hashimoto T, et al. Altered constitutive expression of fatty acid-metabolizing enzymes in mice lacking the peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ). *J Biol Chem* 1998; 273: 5678 – 5684.
73. Leone TC, Weinheimer CJ, Kelly DP. A critical role for the peroxisome proliferator-activated receptor alpha (PPARalpha) in the cellular fasting response: the PPARalpha null mouse as a model of fatty acid oxidation disorders. *Proc Natl Acad Sci U S A* 1999; 96: 7473 – 7478.
74. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W. Peroxisome proliferator-activated receptor  $\alpha$  mediates the adaptive response to fasting. *J Clin Invest* 1999; 103:1489 – 1498.
75. Watts GF, Dimmitt SB. Fibrates, dyslipoproteinaemia and cardiovascular disease. *Curr Opin Lipidol* 1999; 10: 561 – 574.
76. Schultz AE, Alborn WE, Newton RK, Konrad RJ. Administration of PPAR $\alpha$  agonist increases serum apolipoprotein A-V levels and the apolipoprotein A-V/apolipoprotein C-III ratio. *J Lipid Res* 2005; 46: 1591 – 1595.
77. Medina-Gomez G, Gray SL, Yetukuri L, Shimomura K, Virtue S, Campbell M, et al. PPAR Gamma 2 Prevents Lipotoxicity by Controlling Adipose Tissue Expandability and Peripheral Lipid Metabolism. *Plos Genet* 2007; 3: 634 – 647.
78. Repa JJ, Turley SD, Lobaccaro JA, Medina J, Li L, Lustig K, et al. Regulation of absorption and ABC1-mediated efflux of cholesterol by RXR heterodimers. *Science* 2000; 289: 1524 – 1529.
79. Venkateswaran A, Laffitte BA, Joseph SB, Mak PA, Wilpitz DC, Edwards PA, et al. Control of cellular cholesterol efflux by the nuclear oxysterol receptor LXR  $\alpha$ . *Proc Natl Acad Sci USA* 2000; 97: 12097 – 12102.
80. Costet P, Luo Y, Wang N, Tall AR. Sterol-dependent transactivation of the ABC1 promoter by the liver X receptor/retinoid X receptor. *J Biol Chem* 2000; 275: 28240 – 28245.
81. Londos C, Brasaemle DL, Schultz CJ, Segrest JP, Kimmel AR. Perilipins, ADRP, and other proteins that associate with intracellular neutral lipid droplets in animal cells. *Semin Cell Dev Biol* 1999; 10: 51 – 58.
82. Sprecher DL, Massien C, Pearce G, Billin AN, Perlstein I, Willson TM, et al. TG: High – Density Lipoprotein Cholesterol Effects in Healthy Subjects Administered a Peroxisome Proliferator—Activated Receptor  $\delta$  Agonist. *Arterioscler Thromb Vasc Biol* 2007; 27: 359 – 365.
83. Lee CH, Olson P, Havener A, Mehl I, Chong LW, Olefsky JM, et al. PPAR $\delta$  Regulates Glucose Metabolism and Insulin Sensitivity. *Proc Natl Acad Sci U S A* 2006; 103: 3444 – 3449.
84. Cock TA, Auwerx J. PPAR $\gamma$  Fundamental Role in Adipogenesis. *Int Congress Series* 2004; 1262: 47 – 50.
85. Rosen E.D., Walkey, C.J., Puigserver, P. & Spiegelman, B.M. Transcriptional regulation of adipogenesis. *Genes Dev* 2000; 14: 1293 – 1307.
86. Barak Y, Nelson M, Ong E, Jones Y, Ruiz-Lozano P, Chien K, A. et al. PPAR $\gamma$  is required for placental, cardiac, and adipose tissue development. *Mol Cell* 1999; 4: 585 – 595.
87. Rosen ED, Sarraf P, Troy A, Bradwin G, Moore K, Milstone D, et al. PPAR $\gamma$  is required for the differentiation of adipose tissue in vivo and in vitro. *Mol Cell* 1999; 4: 611 – 617.
88. Kubota N, Terauchi Y, Miki H, Tamemoto H, Yamauchi T, Komeda K, et al. PPAR $\gamma$  mediates high-fat diet-induced adipocyte hypertrophy and insulin resistance. *Mol Cell* 1999; 4: 597 – 609.
89. Agarwal AK & Garg A. A novel heterozygous mutation in peroxisome proliferator activated receptor- $\gamma$  gene in a patient with familial partial lipodystrophy. *J Clin Endocrinol Metab* 2002; 87: 408 – 411.
90. Hegele RA, Cao H, Frankowski C, Mathews ST & Leff T. PPAR $\gamma$  F388L, a transactivation-deficient mutant, in familial partial lipodystrophy. *Diabetes* 2002; 51: 3586 – 3590.
91. Savage DB, Tan GD, Acerini CL, Jebb SA, Agostini M, Gurnell M, et al. Human metabolic syndrome resulting from dominant-negative mutations in the nuclear receptor peroxisome proliferator-activated receptor- $\gamma$ . *Diabetes* 2003; 52: 910 – 917.
92. Wu Z, Rosen E, Brun R, Hauser S, Adelmant A, Troy A, et al. Cross-regulation of C/EBP and PPAR $\gamma$  controls the transcriptional pathway of adipogenesis and insulin sensitivity. *Mol Cell* 1999; 3: 151 – 158.
93. Rosen ED, Hsu CH, Wang X, Sakai S, Freeman MW, Gonzalez FJ, et al. C/EBP $\alpha$  induces adipogenesis through PPAR $\gamma$ : a unified pathway. *Genes Dev* 2002; 16: 22 – 26.
94. MacDougald OA, and Mandrup S. Adipogenesis: forces that tip the scales. *Trends Endocrinol Metab* 2002; 13: 5 – 11.
95. Frohnert BI, Hui TY, and Bernlohr DA. Identification of a functional peroxisome proliferator-responsive element in the murine fatty acid transport protein gene. *J Biol Chem* 1999; 274: 3970 – 3977.
96. Chui, PC, Guan HP, Lehrke M, and Lazar M.A. PPARgamma regulates adipocyte cholesterol metabolism via oxidized LDL receptor 1. *J Clin Invest* 2005; 115: 2244 – 2256.
97. Tontonoz P, Hu E, Devine J, Beale EG, and Spiegelman BM. PPAR gamma 2 regulates adipose expression of the phosphoenolpyruvate carboxykinase gene. *Mol Cell Biol* 1995; 15: 351 – 357.
98. Tordjiman J, Chauvet G, Ueuette J, Beale EG, Forest C, and Antoine B. Thiazolidinediones block fatty acid release by inducing glyceroneogenesis in fat cells. *J Biol Chem*. 2003; 278: 18785 – 18790.
99. Hibuse T, Maeda N, Funahashi T, Yamamoto K, Nagasawa A, Mizunoya W, et al. Aquaporin 7 deficiency is associated with development of obesity through activation of adipose

- glycerol kinase. *Proc Natl Acad Sci USA* 2005; 102: 10993 – 10998.
100. Okuno A, Tamemoto H, Tobe K, Ueki K, Mori Y, Iwamoto K, et al. Troglitazone increases the number of small adipocytes without the change of white adipose tissue mass in obese Zucker rats. *J Clin. Invest* 1998; 101: 1354 – 1361.
  101. Wilson-Fritch L, Nicoloso S, Chouinard M, Lazar MA, Chui PC, Leszyk J, Straubhaar J, et al Mitochondrial remodeling in adipose tissue associated with obesity and treatment with rosiglitazone. *J Clin Invest* 2004; 114: 1281 – 1289.
  102. Lehrke M, Lazar MA. The Many Faces of PPAR $\gamma$ . *Cell* 2005; 123: 993 – 999.
  103. Yamauchi T, Kamon J, Waki H, Murakami K, Motojima J, Komeda K, et al. The mechanisms by which both heterozygous peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) deficiency and PPAR $\gamma$  agonist improve insulin resistance. *J Biol Chem* 2001; 276: 41245 – 41254.
  104. Peraldi P, Xu M, Spiegelman BM. Thiazolidinediones block tumor necrosis factor- $\alpha$ -induced inhibition of insulin signaling. *J Clin. Invest* 1997; 100: 1863 – 1869.
  105. Stepan CM, [Bailey ST](#), [Bhat S](#), [Brown EJ](#), [Banerjee RR](#), [Wright CM](#), et al. The hormone resistin links obesity to diabetes. *Nature* 2001; 409: 307 – 312.
  106. Rajala MW, Obici S, Scherer PE, Rossetti L. Adipose-derived resistin and gut-derived resistin-like molecule- $\beta$  selectively impair insulin action on glucose production. *J Clin Invest* 2003; 111: 225 – 300.
  107. Combs TP, Berg AH, Obici S, Scherer PE, Rossetti L. Endogenous glucose production is inhibited by the adipose-derived protein Acrp30. *J Clin Invest* 2001; 108: 1875 – 1881.
  108. Berg AH, Combs TP, Du X, Brownlee M, Scherer PE. The adipocytosecreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 2001; 7: 947 – 953.
  109. Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med* 2001; 7: 941 – 946.
  110. Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, et al. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 2002; 8: 1288 – 1295.
  111. Yamauchi T, Kamon J, Ito Y, Tsuchida A, Yokomizo T, Kita S, et al. Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 2003; 423: 762 – 769.
  112. Yamauchi T, Waki H, Kamin J, Murakami K, Motojima K, Komeda K, et al. Inhibition of RXR and PPAR $\gamma$  Ameliorates Diet – Induced Obesity and Type 2 Diabetes. *J Clin Invest* 2001; 108: 1001 – 1013.
  113. Zhang B, Berger J, Hu E, Szalkowski D, White – Carrington S, Spiegelman BM, et al. Negative Regulation of Peroxisome Proliferator – Activated Receptor -  $\gamma$  Gene Expression Contributes to the Antiadipogenic Effects of Tumor Necrosis Factor –  $\alpha$ . *Mol Endocrinol* 1996; 10: 1457 – 1466.
  114. White, M.F. Insulin signaling in health and disease. *Science* 2003; 302: 1710 – 1711.
  115. Minami A, Iseki M, Kishi K, Wang M, Ogura M, Furukawa N, et al. Increased insulin sensitivity and hypoinsulinemia in APS knockout mice. *Diabetes* 2003; 52: 2657 – 2665.
  116. Duan C, Yang H, White MF, Rui L. Disruption of the SH2-B gene causes agedependent insulin resistance and glucose intolerance. *Mol Cell Biol* 2004; 24: 7435 – 7443.
  117. Molero JC, Jensen TE, Withers PC, Couzens M, Herzog , Thien CBF, et al. c-Cbl-deficient mice have reduced adiposity, higher energy expenditure and improved peripheral insulin action. *J Clin Invest* 2004; 114: 1326 – 1333.
  118. Bard-Chapeau EA, Hevener AL, Long S, Zhang EE, Olefsky JM, Feng GS. Deletion of Gab1 in the liver leads to enhanced glucose tolerance and improved hepatic insulin action. *Nat Med* 2005; 11: 567 – 571.
  119. Hosomi Y, Ogawa KS, Matsuba H, Yoshida M, Okada Y, Yokono K, et al. Characterization of a 60-kilodalton substrate of the insulin receptor kinase. *J Biol Chem* 1994; 269: 11498 – 11502.
  120. Carpino N, [Wisniewski D](#), [Strife A](#), [Marshak D](#), [Kobayashi R](#), [Stillman B](#), et al. p62dok: a constitutively tyrosine-phosphorylated, GAP-associated protein in chronic myelogenous leukemia progenitor cells. *Cell* 1997; 88: 197 – 204.
  121. Yamanashi Y, Baltimore D. Identification of the Abl- and rasGAP-associated 62-kDa protein as a docking protein. *Dok. Cell* 1997; 88: 205 – 211.
  122. Hosooka T, Noguchi T, Kotani K, Nakamura T, Sakaue H, Inoue H, et al. Dok1 Mediates High – Fat Diet – Induced Adipocyte Hypertrophy and Onesity Through Modulation of PPAR -  $\alpha$  Phosphorylation. *Nat Med* 2008; 14: 188 – 193.
  123. Cohen RN. Nuclear Receptor Corepressors and PPAR $\alpha$ . *NRS* 2006; 4: 1 – 4.
  124. Wu Z, [Andersson U](#), [Zhang C](#), [Adelmant G](#), [Mootha V](#), [Troy A](#), et al. Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell* 1999; 98: 115 – 124.
  125. Puigserver, P. [Wu Z](#), [Park CW](#), [Graves R](#), [Wright M](#), [Spiegelman BM](#). A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 1998; 92: 829 – 839.
  126. Oishi Y, Manabe I, Tobe K, Ohsugi M, Kubota T, Fujiu K, et al. SUMOylation of Kr $\mu$ ppel – like Transcription Factor 5 Acts As A Molecular Switch in Transcriptional Programs of Lipid Metabolism Involving PPAR -  $\delta$ . *Nat Med* 2008; 14: 656 – 666.
  127. Dressel U, Allen TL, Pippal JB, Rohde PR, Lau P, Muscat GEO. The peroxisome proliferator-activated receptor b $\mu$ d agonist GW501516, regulates the expression of genes involved in lipid catabolism and energy uncoupling in skeletal muscle cells. *Mol Endocrinol* 2003; 17: 2477 – 2493.
  128. Tanaka T, Yamamoto J, Iwasaki S, Asaba H, Jamura H, Ikeda Y, et al. Activation of peroxisome proliferator-activated receptor d induces fatty acid b-oxidation in skeletal muscle and attenuates metabolic syndrome. *Proc Natl Acad Sci USA* 2003; 100: 15924 – 15929.
  129. Krey G, Braissant O, L $\mu$ HorsetF, Kalkhoven E, Perroud M, Parker MG, et al. Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay. *Mol Endocrinol* 1997; 11: 779 – 791.
  130. Huang JT, Welch JS, Ricote M, Binder CJ, Willson TM, Kelly C, et al. Interleukin-4-dependent production of PPAR- $\gamma$  ligands in macrophages by 12 $\mu$ 15-lipoxygenase. *Nature* 1999; 400: 378 – 382.
  131. Cunard R, Ricote M, DiCampli D, Archer DC, Kahn DA, Glass CK, et al. Regulation of cytokine expression by ligands of peroxisome proliferator activated receptors. *J Immunol* 2002; 168: 2795 – 2802.
  132. Welch JS, Ricote M, Akiyama TE, Gonzalez FJ, Glass CK. PPARgamma and PPARdelta negatively regulate specific subsets of lipopolysaccharide and IFN-gamma target genes in macrophages. *Proc Natl Acad Sci U S A* 2003; 100: 6712 – 6717.
  133. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome proliferator-activated receptor- $\alpha$  is a negative regulator of macrophage activation. *Nature* 1998; 391: 79 – 82.
  134. Chinetti G, Griglio S, Antonucci M, Torra IP, Delerive P, Majd Z, et al. Activation of proliferatoractivated receptors  $\alpha$

- and  $\gamma$  induces apoptosis of human monocytederived macrophages. *J Biol Chem.* 1998; 273: 25573 – 25580.
135. Devchand PR, Keller H, Peters JM, Vazquez M, Gonzalez FJ, Wahli W. The PPAR $\alpha$ -leukotriene B4 pathway to inflammation control. *Nature* 1996; 384: 39 – 43.
  136. Hill MR, Clarke S, Rodgers K, Thornhill B, Peters JM, Gonzalez FJ, et al. Effect of peroxisome proliferator-activated receptor alpha activators on tumor necrosis factor expression in mice during endotoxemia. *Infect Immun* 1999; 67: 3488 – 3493.
  137. Lee H, Shi W, Tontonoz P, Wang S, Subbanagounder G, Hedrick CC, et al. Role for peroxisome proliferator-activated receptor alpha in oxidized phospholipid-induced synthesis of monocyte chemotactic protein-1 and interleukin-8 by endothelial cells. *Circ Res* 2000; 87: 516 – 521.
  138. Ricote M, Valledor AF, Glass CK. Decoding Transcriptional Programs Regulated by PPARs and LXRs in the Macrophage: Effects on Lipid Homeostasis, Inflammation, and Atherosclerosis. *Arterioscler Thromb Vasc Biol* 2004; 24: 230 – 239.
  139. Pasceri V, Chang J, Willerson JT, Yeh ET. Modulation of C-reactive protein-mediated monocyte chemoattractant protein-1 induction in human endothelial cells by antiatherosclerosis drugs. *Circulation* 2001; 103: 2531 – 2534.
  140. Ikewaki K, Tohyama J, Nakata Y, Wakikawa T, Kido T, Mochizuki S. Fenofibrate effectively reduces remnants and small dense LDL and increases HDL particle number in hyperTGmic men—a nuclear magnetic resonance study. *J Atheroscler Thromb.* 2004; 11: 278 – 285.
  141. Wang TD, Chen WJ, Lin JW, Cheng CC, Chen MF, Lee YT. Efficacy of fenofibrate and simvastatin on endothelial function and inflammatory markers in patients with combined hyperlipidemia: relations with baseline lipid profiles. *Atherosclerosis* 2003; 170: 315 – 323.
  142. Han SH, Quon MJ, Koh KK. Beneficial Vascular and Metabolic Effects of Peroxisome Proliferator – Activated Receptor –  $\alpha$  Activators. *Hypertension* 2005; 46: 1086 – 1092.
  143. Teissier E, Nohara A, Chonetti G, Paumelle R, Cariou B, Fruchart JC, et al. Peroxisome proliferator-activated receptor  $\alpha$  induces NADPH oxidase activity in macrophages, leading to the generation of LDL with PPAR- $\alpha$  activation properties. *Circ Res* 2004; 95: 1174 – 1182.
  144. Boullier A, Bird DA, Chang MK, Dennis EA, Friedman P, Gillotre-Taylor K, et al. Scavenger receptors, oxidized LDL, and atherosclerosis. *Ann N Y Acad Sci.* 2001; 947: 214 – 222; discussion 222 – 223.
  145. Devchand PR, Ziouzenkova O, Plutzky J. Oxidative Stress and Peroxisome Proliferator – Activated Receptors. Reversing or Curse? *Circ Res* 2004; 95: 1137 – 1139.
  146. Chawla A, Barak Y, Nagy L, [Liao D](#), [Tontonoz P](#), [Evans RM](#). PPAR-gamma dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation. *Nature Med* 2001; 7: 48 – 52.
  147. Jiang C, Ting AT, Seed B. PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 1998; 391: 82 – 86.
  148. Ricote M, Huang J, Fajas L, Li A, Welch J, Najib J, et al. Expression of the peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoproteins. *Proc Natl Acad Sci USA* 1998; 95: 7614 – 7619.
  149. Moore KJ, Rosen ED, Fitzgerald ML, [Rando F](#), [Andersson LP](#), [Altschuler D](#), et al. The role of PPAR-gamma in macrophage differentiation and cholesterol uptake. *Nature Med* 2001; 7: 41 – 47.
  150. Thieringer R, Fenyk-Melody JE, Le Grand CB, Shelton BA, Detmers PA, Somers EP, et al. Activation of peroxisome proliferator-activated receptor gamma does not inhibit IL-6 or TNF-alpha responses of macrophages to lipopolysaccharide in vitro or in vivo. *J Immunol* 2000; 164: 1046 – 1054.
  151. Bamba H, Ota S, Kato A, Kawamoto C, Fujiwara K. Prostaglandins up-regulate vascular endothelial growth factor production through distinct pathways in differentiated U937 cells. *Biochem Biophys Res Commun* 2000; 273: 485 – 491.
  152. Moore KJ, Fitzgerald ML, Freeman MW. Peroxisome Proliferator – Activated Receptors in Macrophage Biology: Friend or Foe? *Curr Opin Lipidol* 2001; 12: 519 – 527.
  153. Chinetti G, Fruchart JC, Staels B. Peroxisome proliferator-activated receptors (PPARs): nuclear receptors at the crossroads between lipid metabolism and inflammation. *Inflamm Res* 2000; 49: 497 – 505.
  154. Kintscher U, Goetze S, Wakino S, Kim S, Nagpal S, Chandraratna RA, et al. Peroxisome proliferator-activated receptor and retinoid X receptor ligands inhibit monocyte chemotactic protein-1-directed migration of monocytes. *Eur J Pharmacol* 2000; 401: 259 – 270.
  155. Tanaka T, Fukunaga Y, Itoh H, Doi K, Yamashita J, Chun TH, et al. Therapeutic potential of thiazolidinediones in activation of peroxisome proliferator-activated receptor gamma for monocyte recruitment and endothelial regeneration. *Eur J Pharmacol* 2005; 508: 255 – 265.
  156. Rigamonti E, Fontaine C, Lefebvre B, Duhem C, Lefebvre P, Marx N, et al. Induction of CXCR2 Receptor by Peroxisome Proliferator – Activated Receptor  $\gamma$  in Human Macrophages. *Arterioscler Thromb Vasc Biol* 2008; 28: 932 – 939.
  157. Marx N, Bourcier T, Sukhova GK, Libby P, Plutzky J. PPARgamma activation in human endothelial cells increases plasminogen activator inhibitor type-1 expression: PPARgamma as a potential mediator in vascular disease. *Arterioscler Thromb Vasc Biol* 1999; 19: 546 – 551.
  158. Plutzky J. Peroxisome Proliferator-Activated Receptors in Endothelial Cell Biology. *Curr Opin Lipidol* 2001; 12: 511.
  159. Hwang J, Kleinhenz DJ, Lassègue B, Griendling KK, Dikalov S, Hart CM. Peroxisome Proliferator-Activated Receptors -  $\gamma$  Ligands Regulate Endothelial Membrane Superoxide Production. *Am J Physiol Cell Physiol* 2005; 288: C899 – C905.
  160. Takata Y, Kitami Y, Yang ZH, Nakamura M, Okura T, Hiwada K. Vascular Inflammation Is negatively Autoregulated by Interaction Between CCAAT/Enhancer – Binding Protein -  $\delta$  and Peroxisome Proliferator – Activated Receptor -  $\gamma$ . *Circ Res* 2002; 91: 427 – 433.
  161. Lee CH, Chawla A, Urbiztondo N, Liao D, Boisvert W, Evans EM. Transcriptional Repression of Atherogenic Inflammation: Modulation by PPAR $\delta$ . *Science* 2003; 302: 453 – 457.
  162. Lee CH, Kang K, Mehl IR, Nofsinger R, Alaynick WA, Chong LW, et al. Peroxisome Proliferator-Activated Receptor  $\delta$  Promotes Very Low – Density Lipoprotein – Derived Fatty Acid Catabolism in the macrophage. *Proc Natl Acad Sci U S A* 2006; 103: 2434 – 2439.
  163. Fan Y, Wang Y, Tang Z, Zhang H, Qin X, Zhu Y, et al. Suppression pro-Inflammatory Adhesion Molecules by PPAR- $\delta$  in Human Vascular Endothelial Cells. *Arteriol Thromb Vasc Biol* 2008; 28: 315 – 321.
  164. Kim HJ, Ham SA, Kim SU, Hwang JY, Kim JH, Chang KC, et al. Transforming Growth factor -  $\beta$ 1 Is a Molecular Target for the Peroxisome Proliferator – Activated Receptor  $\delta$ . *Circ Res* 2008; 102: 193 – 200.
  165. Yagil C and Yagil Y. Peroxisome Proliferator – Activated Receptors –  $\alpha$ . Friend or Foe? *Hypertension* 2007; 50: 847 – 850.
  166. Tordjman K, Semenkovich CF, Coleman T, Yudovich R, Bak S, Osher E, et al. Absence of peroxisome proliferator-activated

- receptor- $\alpha$  abolishes hypertension and attenuates atherosclerosis in the Tsukuba hypertensive mouse. *Hypertension* 2007; 50: 945 – 951.
167. Marx N, Duez H, Fruchart JC, Staels B. Peroxisome Proliferator – Activated Receptors and Atherogenesis. *Regulators of Gene Expression in Vascular Cells. Circ Res* 2004; 94: 1168 – 1178.
168. Duez H, Chao YS, Hernandez M, Torpier G, Poulain P, Mundt S, et al. Reduction of atherosclerosis by the peroxisome proliferator-activated receptor agonist fenofibrate in mice. *J Biol Chem* 2002; 277: 48051 – 48057.
169. Vosper H, Khoudoli GA, Graham TL, Palmer CNA. Peroxisome proliferator-activated agonists, hyperlipidaemia, and atherosclerosis. *Pharm Ther* 2002; 95: 47 – 62.
170. Diep QN, Amiri F, Touyz RM, Cohn JS, Endemann D, Schiffrin EL. PPAR $\alpha$  activator effects on ANG II-induced vascular oxidative stress and inflammation. *Hypertension* 2002; 40: 866 – 871.
171. Iglarz M, Touyz RM, Amiri F, Lavoie MF, Diep QN, Schiffrin EL. Effect of peroxisome proliferator-activated receptor- $\alpha$  and - $\gamma$  activators on vascular remodeling in endothelin-dependent hypertension. *Arterioscler Thromb Vasc Biol* 2003; 23: 45 – 51.
172. Banks T, Oyekan A. Peroxisome proliferator-activated receptor  $\alpha$  activation attenuated angiotensin type 1-mediated but enhanced angiotensin type 2-mediated hemodynamic effects to angiotensin II in the rat. *J Hypertension* 2008; 26: 468 – 477.
173. Granger JP. Endothelin. *Am J Physiol Regul Integr Comp Physiol* 2003; 285: R298–R301.
174. Rubanyi GM, Polokoff MA. Endothelins: molecular biology, biochemistry, pharmacology, physiology, and pathophysiology. *Pharmacol Rev* 1994; 46: 325 – 415.
175. Pollock DM. Renal endothelin in hypertension. *Curr Opin Nephrol Hypertens* 2000; 9: 157 – 164.
176. Roman RJ. P-450 metabolites of arachidonic acid in the control of cardiovascular function. *Physiol Rev* 2002; 82: 131 – 185.
177. Alonso-Galicia M, Frohlich B, Roman RJ. Induction of P4504A activity improves pressure-natriuresis in Dahl S rats. *Hypertensio*. 1998; 31: 232 – 236.
178. Roman RJ, Ma YH, Frohlich B, Markham B. Clofibrate prevents the development of hypertension in Dahl salt-sensitive rats. *Hypertension* 1993; 21: 985 – 988.
179. Dos Santos EA, Dahly-Vernon AJ, Hoagland KM, Roman RJ. Inhibition of the formation of EETs and 20-HETE with 1-aminobenzotriazole attenuates pressure natriuresis. *Am J Physiol Regul Integr Comp Physiol* 2004; 287: R58 – R68.
180. Zhang YB, Magyar CE, Holstein-Rathlou NH, McDonough AA. The cytochrome P-450 inhibitor cobalt chloride prevents inhibition of renal Na<sup>+</sup>,K<sup>+</sup>-ATPase and redistribution of apical NHE-3 during acute hypertension. *J Am Soc Nephrol* 1998; 9: 531 – 537.
181. Hoagland KM, Flasch AK, Roman RJ. Inhibitors of 20-HETE formation promote salt-sensitive hypertension in rats. *Hypertension* 2003; 42: 669 – 673.
182. Hoagland KM, Flasch AK, Dahly-Vernon AJ, dos Santos EA, Knepper MA, Roman RJ. Elevated BSC-1 and ROMK expression in Dahl salt-sensitive rat kidneys. *Hypertension* 2004; 43: 860 – 865.
183. Wilson TW, Alonso-Galicia M, Roman RJ. Effects of lipid-lowering agents in the Dahl salt-sensitive rat. *Hypertension* 1998; 31: 225 – 231.
184. Williams JM, Zhao X, Wang MH, Imig JD, Pollock DM. Peroxisome Proliferator-Activated Receptor- $\alpha$  Activation Reduces Salt-Dependent Hypertension During Chronic Endothelin B Receptor Blockade. *Hypertension* 2005; 46: 366 – 371.
185. Subramanian S, DeRosa MA, Bernal – Mizrahi C, Laffely N, Cade WT, Yarasheski KE, et al. PPAR $\alpha$  Activation Elevates Blood Pressure and Does Not Correct Glucocorticoid – Induced Insulin Resistance in Humans. *Am J Physiol Endocrinol Metab* 2006; 291: E1365 – E1371.
186. Weatherford ET, Itani H, Keen HL, Sigmund CD. Is Peroxisome Proliferator-Activated Receptor- $\gamma$  a New “Pal” of Renin? *Hypertension* 2007; 50: 844 – 846.
187. Shi Q, Gross KW, Sigmund CD. Retinoic acid-mediated activation of the mouse renin enhancer. *J Biol Chem* 2001; 276: 3597 – 3603.
188. Li YC, Kong J, Wei M, Chen Z-F, Liu SQ, Cao L-P. 1,25-Dihydroxyvitamin D3 is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest* 2002; 110: 229 – 238.
189. Sugawara A, Takeuchi K, Iruno A, Ikeda Y, Arima S, Kudo M, et al. Transcriptional suppression of type 1 angiotensin II receptor gene expression by peroxisome proliferator-activated receptor-gamma in vascular smooth muscle cells. *Endocrinology*. 2001; 142: 3125 – 3134.
190. Grundy SM, Hansen B, Smith SC Jr, Cleeman JI, Kahn RA. Clinical management of metabolic syndrome: report of the Am Heart Association/National Heart, Lung, and Blood Institute/AmDiabetes Association conference on scientific issues related to management. *Circulation* 2004; 109: 551 – 556.
191. Szapary PO, Bloedon LT, Samaha FF, Duffy D, Wolfe ML, Soffer D, et al. Effects of Pioglitazone on Lipoproteins, Inflammatory Markers, and Adipokines in Nondiabetic Patients with Metabolic Syndrome. *Arterioscler Thromb Vasc Biol* 2006; 26: 182 – 188.
192. Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nissen M, et al. Cardiovascular Morbidity and Mortality Associated with the metabolic Syndrome. *Diab Care* 2001; 24: 683.
193. Stern MP, Williams K, Gonzalez – Villalpando C, Hunt KJ, Haffner SM. Does the Metabolic Syndrome Improve Identification of Individuals at Risk of Type 2 Diabetes and/or Cardiovascular Disease? *Diab Care* 2004; 27: 2676 – 2681.
194. Cignarella A, Bellocchio S, Corsini A, Bolego C. Hypolipidemic Therapy for The Metabolic Syndrome. *Pharmacol Res* 2006; 53: 492 – 500.
195. Semple RK, Chatterjee VK, O'Rahilly S. PPAR $\gamma$  and human metabolic disease. *J Clin Invest* 2006; 116: 581 – 589.
196. Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 2002; 287: 356 – 359.
197. Szapary PO, Rader DJ. The TG-high-density lipoprotein axis: an important target of therapy? *Am Heart J* 2004; 148: 211 – 221.
198. Barish GD, Narkar VA, Evans RM. PPAR $\delta$ : A Dagger in The Heart of The Metabolic Syndrome. *J Clin Invest* 2006; 116: 590 – 597.
199. Eckel RH, Grundy SM, Zimmet PZ. The metabolic syndrome. *Lancet* 2005; 365: 1415 – 1428.
200. Grundy SM. Point: the metabolic syndrome still lives. *Clin Chem* 2005; 51: 1352 – 1354.
201. Kershaw EE, Flier JS. Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab* 2004; 89: 2548 – 2556.
202. Frayn KN. Adipose tissue as a buffer for daily lipid flux. *Diabetologia* 2002; 45: 1201 – 1210.
203. Perseghin G. Muscle lipid metabolism in the metabolic syndrome. *Curr Opin Lipidol* 2005; 16: 416 – 420.

204. Yki-Jarvinen H: Fat in the liver and insulin resistance. *Ann Med* 2005; 37: 347 – 356.
205. Heilbronn L, Smith SR, Ravussin E: Failure of fat cell proliferation, mitochondrial function and fat oxidation results in ectopic fat storage, insulin resistance and type II diabetes mellitus. *Int J Obes Relat Metab Disord* 2004; 28 (Suppl 4): S12 – S21.
206. Goodpaster BH, Wolf D: Skeletal muscle lipid accumulation in obesity, insulin resistance, and type 2 diabetes. *Pediatr Diabetes* 5:219 –226, 2004 6. Petersen KF, Shulman GI: Etiology of insulin resistance. *Am J Med* 2006; 119: S10 – S16.
207. Blaschke F, Takata Y, Caglayan E, Law RE, Hsueh WA. Obesity, Peroxisome Proliferator-Activated Receptor, and Atherosclerosis in Type 2 Diabetes. *Arteriocler Throm Vasc Biol* 2006; 26: 28 – 40.
208. Ruan H, Lodish HF. Insulin resistance in adipose tissue: direct and indirect effects of tumor necrosis factor- $\alpha$ . *Cytokine Growth Factor Rev* 2003; 14: 447 – 455.
209. Aizawa-Abe M, Ogawa Y, Masuzaki H, Ebihara K, Satoh N, Iwai H, et al. Pathophysiological role of leptin in obesity-related hypertension. *J Clin Invest* 2000; 105: 1243 – 1252.
210. Cassis LA, Saye J, Peach MJ. Location and regulation of rat angiotensinogen messenger RNA. *Hypertension* 1988; 11: 591 – 596.
211. Cigolini M, Tonoli M, Borgato L, Frigotto L, Manzato F, Zeminian S, et al. Expression of plasminogen activator inhibitor-1 in human adipose tissue: a role for TNF- $\alpha$ ? *Atherosclerosis* 1999; 143: 81 – 90.
212. Lijnen HR. Pleiotropic functions of plasminogen activator inhibitor-1. *J Thromb Haemost* 2005; 3: 35 – 44.
213. Kubota N, Terauchi Y, Yamauchi T, Kubota T, Moroi M, Matsui J, et al. Disruption of adiponectin causes insulin resistance and neointimal formation. *J Biol Chem* 2002; 277: 25863 – 25866.
214. Yamauchi T, Kamon J, Waki H, Imai Y, Shimozawa N, Hioki K, et al. Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis. *J Biol Chem* 2003; 278: 2461 – 2468.
215. Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, et al. Adiponectin, an adipocytederived plasma protein, inhibits endothelial NF- $\kappa$ B signaling through a cAMP-dependent pathway. *Circulation* 2000; 102: 1296 – 1301.
216. Fukuhara A, Matsuda M, Nishizawa M, Segawa K, Tanaka M, Kishimoto K, et al. Visfatin: a protein secreted by visceral fat that mimics the effects of insulin. *Science* 2005; 307: 426 – 430.
217. Rise'rus U, Sprecher D, Johnson T, Olson E, Hirschberg S, Liu A, et al. Activation of Peroxisome Proliferator-Activated Receptor (PPAR) $\gamma$  Promotes Reversal of Multiple Metabolic Abnormalities, Reduces Oxidative Stress, and Increases Fatty Acid Oxidation in Moderately Obese Men. *Diabetes* 2008; 57: 332 – 229.
218. Liang CP, Han S, Okamoto H, Carnemolla R, Tabas I, Accili D, et al. Increased CD36 protein as a response to defective insulin signaling in macrophages. *J Clin Invest* 2004; 113: 764 – 773.
219. Shoelson SE, Lee J, Goldfine AB. Inflammation and insulin resistance. *J Clin Invest* 2006; 116: 1793 – 1801.
220. Hotamisligil GS. Inflammation and metabolic disorders. *Nature* 2006; 444: 860 – 867.
221. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante, Jr AW.. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003; 112: 1796 – 1808.
222. Xu, H, [Barnes GT](#), [Yang Q](#), [Tan G](#), [Yang D](#), [Chou CJ](#), et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 2003; 112: 1821 – 1830.
223. Odegaard Ricardo-Gonzalez RR, Goforth MH, Morel CR, Subramanian V, Mukundan L, Eagle AR, et al. Macrophage-specific PPAR $\gamma$  controls alternative activation and improves insulin resistance. *Nature* 2007; 447: 1116 – 1121.
224. Brown JD, Plutzky JP. Peroxisome Proliferator-Activated Receptor as Transcriptional Nodal Points and Therapeutic targets. *Circulation* 2007; 115: 518 – 533.
225. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and Management of the Metabolic Syndrome – an American Heart Association/ National Heart, lung, and Blood Institute Scientific Statement. *Circulation* 2005; 112: 2735 – 2752.
226. Chinetti – Gbaguidi G, Fruchart JC, Staels B, Role of PPAR family of Nuclear Receptors in the Regulation of Metabolic and Cardiovascular Homeostasis: New Approach to Therapies. *Curr Opin Pharmacol* 2005; 5: 177 – 183.
227. Tordjman K, Bernal – Mizrachi C, Zemanly L, Weng S, Feng C, Zhang F, Lene TC, et al. PPAR $\alpha$  Deficiency Reduces Insulin Resistance And Atherosclerosis in apoE – null Mice. *J Clin Invest* 2001; 107: 1025 – 1034.
228. Berger JP. Akiyama TE, Meinke PT. PPARs: Therapeutic Targets for Metabolic Disease. *Trends Pharmacol Sci* 2005; 26: 244 – 251.
229. Maragoudakis ME, Hankin H. On the mode of action of lipid-lowering agents. V. Kinetics of the inhibition in vitro of rat acetyl coenzyme A carboxylase. *J Biol Chem* 1971; 246: 348 – 358.
230. Staels B, Dallongeville J, Auwerx J, Schoonjans K, Leitersdorf E, Fruchart JC. Mechanism of action of fibrates on lipid and lipoprotein metabolism. *Circulation* 1998; 98: 2088 – 2093.
231. Heller F, Harvengt C. Effects of clofibrate, bezafibrate, fenofibrate and probucol on plasma lipolytic enzymes in normolipaeamic subjects. *Eur J Clin Pharmacol* 1983; 25: 57 – 63.
232. Staels B, Vu-Dac N, Kosykh VA, Saladin R, Fruchart JC, Dallongeville J, et al/ Fibrates downregulate apolipoprotein C-III expression independent of induction of peroxisomal acyl coenzyme A oxidase. A potential mechanism for the hypolipidemic action of fibrates. *J Clin Invest* 1995; 95: 705 – 712.
233. Malmendier CL, Delcroix C. Effects of fenofibrate on high and low density lipoprotein metabolism in heterozygous familial hypercholesterolemia. *Atherosclerosis* 1985; 55: 161 – 169.
234. Mellies MJ, Stein EA, Khoury P, Lamkin G, Glueck CJ. Effects of fenofibrate on lipids, lipoproteins, and apolipoproteins in 33 subjects with primary hypercholesterolemia. *Atherosclerosis* 1987; 63: 57 – 64.
235. Knopp RH, Walden CE, Warnick GR, Albers JJ, Ginsberg J, McGinnis BM. Effect of fenofibrate treatment on plasma lipoprotein lipids, highdensity lipoprotein cholesterol subfractions, and apolipoproteins B, AI, AII, and E. *Am J Med* 1987; 83: 75 – 84.
236. Bard JM, Parra HJ, Camare R, Luc G, Ziegler O, Datchet C, et I. A multicenter comparison of the effects of simvastatin and fenofibrate therapy in severe primary hypercholesterolemia, with particular emphasis on lipoproteins defined by their apolipoprotein composition. *Metabolism* 1992; 41: 498 – 503.
237. Chinetti G, Lestavel S, Bocher V, Remaley AT, Neve B, Torra IP, et al. PPAR- $\gamma$  and PPAR- $\gamma$  activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway. *Nat Med* 2001; 7: 53 – 58.
238. Li AC, Binder CJ, Gutierrez A, Brown KK, Plotkin CR, Pattison JW, et al. Differential Inhibition of Macrophage Foam – Cell

- Formation and Atherosclerosis in Mice by PPAR $\alpha$ ,  $\beta/\delta$ , and  $\gamma$ . *J Clin Invest* 2004; 114: 1564 – 1576.
239. Mardones P, Pilon A, Bouly M, Duran D, Nishimoto T, Arai H, Kozarsky KF, Altayo M, Miquel JF, Luc G, Clavey V, Staels B, Rigotti A et al. Fibrates down-regulate hepatic scavenger receptor class B type I protein expression in mice. *J Biol Chem* 2003; 278:7 884 – 7890.
240. Tokuno A, Hirano T, Hayashi T, Mori Y, Yamamoto T, Nagashima M, et al. The effects of statin and fibrate on lowering small dense LDL- cholesterol in hyperlipidemic patients with type 2 diabetes. *J Atheroscler Thromb* 2007; 14: 128 – 132.
241. Barter PJ, Rye KA. Is There A Role for Fibrates in the Management of Dyslipidemia in the Metabolic Syndrome? *Arterioscler Thromb Vasc Biol* 2008; 28: 39 – 46.
242. Picard F, Auwerx J. PPAR(gamma) and glucose homeostasis. *Annu Rev Nutr* 2002; 22: 167 – 197.
243. Miyazaki Y, Mahankali A, Matsuda M, Jean H, Kenneth C, Mandarino LJ, et al. Effect of pioglitazone on abdominal fat distribution and insulin sensitivity in type 2 diabetic patients. *J Clin Endocrinol Metab* 2002; 87: 2784 – 2791.
244. Maeda N, Shimomura I, Kishida K, Nishizawa H, Matsuda H, Nagaretani H, et al. Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Nat Med* 2002; 8: 731 – 737.
245. Matsuda M, Shimomura I, Sata M, Arita Y, Nishida M, Maeda N, et al. Role of adiponectin in preventing vascular stenosis: the missing link of adipo-vascular axis. *J Biol Chem* 2002; 277: 37487 – 37491.
246. Peraldi P, Spiegelman B. TNF-alpha and insulin resistance: summary and future prospects. *Mol Cell Biochem* 1998; 182: 169 – 175.
247. Steppan CM, Bailey ST, Bhat S, [Brown EJ](#), [Banerjee RR](#), [Wright CM](#), et al. The hormone resistin links obesity to diabetes. *Nature* 2001; 409: 307 – 312.
248. Berger J, Tanen M, Elbrecht A, **Hermanowski-Vosatka A**, **Moller DE**, **Wright SD**, et al. Peroxisome proliferator-activated receptor gamma ligands inhibit adipocyte 11betahydroxysteroid dehydrogenase type 1 expression and activity. *J Biol Chem* 2001; 276: 12629 – 12635.
249. Masuzaki H, Paterson J, Shinyama H, [Morton NM](#), [Mullins JJ](#), [Seckl JR](#), [Flier JS](#). A transgenic model of visceral obesity and the metabolic syndrome. *Science* 2001; 294: 2166 – 2170.
250. Maeda N, Takahashi M, Funahashi T, **Kihara S**, **Nishizawa H**, **Kishida K**, et al. PPARgamma ligands increase expression and plasma concentrations of adiponectin, an adipose-derived protein. *Diabetes* 2001; 50: 2094 – 2099.
251. Walker BR. How will we know if 11betahydroxysteroid dehydrogenases are important in common diseases. *Clin Endocrinol (Oxf)* 2000; 52: 401 – 402.
252. Samaha FF, Szapary PO, Iqbal N, Williams MM, Bloedon LT, Kochar A, et al. Effects of Rosglitazone of Lipids, Adipokines, and Inflammatory Markers in Nondiabetic Patients With Low High – Density Lipoprotein Cholesterol and Metabolic Syndrome. *Arterioscler Thromb Vasc Biol* 2006; 26: 624 – 630.
253. Chinetti G, Zawadski C, Fruchart JC, Staels B. Expression of adiponectin receptors in human macrophages and regulation by agonists of the nuclear receptors PPARalpha, PPARgamma, and LXR. *Biochem Biophys Res Commun* 2004; 314: 151 – 158.
254. Tiikkainen M, Hakkinen A-M, Korshennikova E, Nyman T, Makimattila S, Yki-Jarvinen H. Effects of rosiglitazone and metformin on liver fat content, hepatic insulin resistance, insulin clearance, and gene expression in adipose tissue in patients with type 2 diabetes. *Diabetes* 2004; 53: 2169 – 2176.
255. Pischon T, Girman CJ, Hotamisligil GS, Rifai N, Hu FB, Rimm EB. Plasma adiponectin levels and risk of myocardial infarction in men. *JAMA* 2004; 291: 1730 – 1737.
256. Junga HS, Younb B-S, Choa YM, Yub K-Y, Parkb HJ, Shina CS, et al. The effects of rosiglitazone and metformin on the plasma concentrations of resistin in patients with type 2 diabetes mellitus. *Metabolism* 2005; 54: 314 – 320.
257. Reilly MP, Lehrke M, Wolfe ML, Rohatgi A, Lazar MA, Rader DJ. Resistin is an inflammatory marker of atherosclerosis in humans. *Circulation* 2005; 111: 932 – 939.
258. Lehrke M, Reilly M, Millington S, Iqbal N, Rader D, and Lazar MA. An inflammatory cascade leading to hyperresistinemia in humans. *PLoS Med* 2004; 1: 161 – 168.
259. Haffner SM, Greenberg AS, Weston WM, Chen H, Williams K, Freed MI. Effect of rosiglitazone treatment on nontraditional markers of cardiovascular disease in patients with type 2 diabetes mellitus. *Circulation* 2002; 106: 679 – 684.
260. Sidhu JS, Cowan D, Kaski J-C. The effects of rosiglitazone, a peroxisome proliferator-activated receptor-gamma agonist, on markers of endothelial cell activation, C-reactive protein, and fibrinogen levels in non-diabetic coronary artery disease patients. *J Am Coll Cardiol* 2003; 42: 1757 – 1763.
261. Wang T-D, Chen W-J, Lin J-W, Chen M-F, Lee Y-T. Effects of rosiglitazone on endothelial function, C-reactive protein, and components of the metabolic syndrome in nondiabetic patients with the metabolic syndrome. *Am J Cardio*. 2004; 93: 362 – 365.
262. Yki-Järvinen H. Thiazolidinediones. *N Engl J Med* 2004; 351: 1106 – 1118.
263. Wayman NS, Hattori Y, McDonald MC, [Mota-Filipe H](#), [Cuzzocrea S](#), [Pisano B](#), et al. Ligands of the peroxisome proliferator-activated receptors (PPAR-gamma and PPAR-alpha) reduce myocardial infarct size. *FASEB J* 2002; 16: 1027 – 1040.
264. Mori Y, Itoh Y, Obata T, Tajima N. Effects of pioglitazone vs glibenclamide on postprandial increases in glucose and TG levels and on oxidative stress in Japanese patients with type 2 diabetes. *Endocrine* 2006; 29: 143 – 148.
265. Campia U, Matuskey LA, Panza JA. Peroxisome proliferator-activated receptor-gamma activation with pioglitazone improves endothelium-dependent dilation in nondiabetic patients with major cardiovascular risk factors. *Circulation* 2006; 113: 867 – 875.
266. Sourij H, Zweiker R, Wascher TC. Effects of pioglitazone on endothelial function, insulin sensitivity, and glucose control in subjects with coronary artery disease and new-onset type 2 diabetes. *Diabetes Care* 2006; 29: 1039 – 1045.
267. Ceriello A. Thiazolidinediones As Anti – inflammatory and Anti – Atherogenic Agents. *Diabetes Metab Res Rev* 2008; 24: 14 – 26.
268. Goldfine AB. The Rough Road for Rosglitazone. *Curr Opin Endocrinol Metab* 2008; 15: 113 – 117.
269. Home PD, Phil D, Pocock SJ, Beck – Nielsen H, Gornis R, Hanefeld M, Jones NP, Komadja M, McMurray JJV. Rosglitazone Evaluated for Cardiovascular Outcomes – An Interim Analysis. *N Engl J med* 2007; 357: 28 – 38.
270. Miller NE, Thelle DS, Forde OH, Mjos OD. The Tromso heart-study. High-density lipoprotein and coronary heart-disease: a prospective case-control study. *Lancet* 1977; 1: 965 – 968.
271. Wilson PW, Abbott RD, Castelli WP. High density lipoprotein cholesterol and mortality. The Framingham Heart Study. *Arteriosclerosis* 1988; 8: 737 – 741.
272. Benoit P, Emmanuel F, Caillaud JM, bassinet L, Castro G, Gallix P, et al. Somatic gene transfer of human ApoA-I inhibits atherosclerosis progression in mouse models. *Circulation* 1999; 99: 105 – 110.

273. van der Veen JN, [Kruit JK](#), [Havinga R](#), [Baller JF](#), [Chimini G](#), [Lestavel S](#), et al. Reduced cholesterol absorption upon PPARdelta activation coincides with decreased intestinal expression of NPC1L1. *J. Lipid Res* 2005; 46: 526 – 534.
274. Staels B, Fruchart JC. Therapeutic Roles of Peroxisome Proliferator-Activated Receptor Agonist. *Diabetes* 2005; 54: 2460 – 2470.
275. Carr DB, **Utzschneider KM**, **Hull RL**, **Kodama K**, **Retzlaff BM**, **Brunzell JD**, et al. Intra-abdominal fat is a major determinant of the National Cholesterol Education Program Adult Treatment Panel III criteria for the metabolic syndrome. *Diabetes* 2004; 53: 2087 – 2094.
276. [Anonymous]. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: the evidence report. National Institutes of Health. *Obes Res* 1988; 6(Suppl. 2): 51S – 209S.
277. Yanovski SZ, Yanovski JA. Obesity. *N Engl J Med* 2002; 346: 591 – 602.
278. Lee CH, et al. PPAR delta is a potent insulin sensitizer. *Proc Natl Acad Sci USA* 2006; in press.
279. Flévet C, Fruchart JC, Staels B. PPAR $\alpha$  and PPAR $\gamma$  Dual Agonists for The Treatment of Type 2 Diabetes and The Metabolic Syndrome. *Curr Opin Pharmacol* 2006; 6: 606.
280. Harrity T, Farrelly D, Tieman A, Chu C, Kunselman L, Gu L, et al. Muraglitazar, a Novel Dual ( $\alpha/\gamma$ ) Peroxisome Proliferator-Activated Receptor Activator, Improves Diabetes and Other Metabolic Abnormalities and Preserves  $\beta$ -Cell Function in db/db Mice. *Diabetes* 2006; 55: 240 – 248.
281. Fagerberg B, Edward S, Halmos T, Lopatynsky J, Schuster H, Stender S, et al. Tesaglitazar, a Novel Dual Peroxisome Proliferator – Activated Receptor  $\alpha/\gamma$  Agonist, Dose – Dependently Improves the Metabolic Abnormalities Associated With Insulin Resistance in A Non – Diabetic Population. *Diabetologia* 2005; 48: 1716 – 1725.
282. Han KL, Choi JS, Lee JY, Song J, Joe MK, Jung MH. Therapeutic Potential of Peroxisome Proliferator – Activated Receptor -  $\alpha/\gamma$  Dual Agonist With Alleviation of Endoplasmic Reticulum Stress for the Treatment of Diabetes. *Diabetes* 2008; 57: 737 – 745.