

Epstein-Barr Virus Infection as One of the Predisposition Factor for Colorectal Cancer

Marcellus Simadibrata

Department of Medical Education, Faculty of Medicine, University of Indonesia
and Division of Gastroenterology, Department of Internal Medicine
Dr. Cipto Mangunkusumo General National Hospital, Jakarta

There have been many studies reported that Epstein-Barr virus (EBV) infection is associated with nasopharyngeal cancer and Hodgkin's lymphoma.^{1,2} Recently, many studies find that Epstein-Barr virus infection is also associated with the occurrence of other cancers such as colorectal cancer.^{1,2} Song et al, reported that EBV LMP1 exon 3 and W fragments were detected in 27.7% and 32.2% of the colorectal carcinoma specimens, which were significantly higher than the positive rate of EBV gene in the adjacent non-cancerous tissues (4.0%, $p < 0.001$).¹

Epstein-Barr virus (EBV) nuclear antigen (EBNA)-1 is the only viral protein expressed in all EBV-carrying malignancies. Gruhne et al, found that EBNA-1 induced chromosomal aberrations, DNA double-strand breaks, and engagement of the DNA damage response (DDR). EBNA-1 promotes genomic instability via induction of reactive oxygen species.³

In this new issue, Simatupang et al, found that EBV viral infection plays a role in the pathology of colon cancer. The findings of EBNA-1 expressions on lymphocytes B membranes proved that there was a chronic EBV infection in patients with colorectal carcinoma. This finding is inline with other studies which found the correlation between Epstein-Barr virus infection and colorectal cancer.

REFERENCES:

1. Song LB, Zhang X, Zhang CQ, Zhang Y, Pan ZZ, Liao WT, et al. Infection of Epstein-Barr virus in colorectal cancer in Chinese. *Ai Zheng* 2006;25:1356-60.
2. Besson C, Amiel C, Le-Pendeven C, Brice P, Ferme C, Carde P, et al. Positive correlation between Epstein-Barr virus viral load and anti-viral capsid immunoglobulin G titers determined for Hodgkin's lymphoma patients and their relatives. *J Clin Microbiol* 2006;44:47-50.
3. Gruhne B, Sompallae R, Marescotti D, Kamranvar SA, Gastaldello S, Masucci MG. The Epstein-Barr virus nuclear antigen-1 promotes genomic instability via induction of reactive oxygen species. *PNAS* 2009;106:2313-8.