

Study of Oxidative Stress in Relation with Antioxidant Status in Chronic Bronchitis

Anita M Raut*, A.N. Suryakar**, Dilip Mhaisekar***

*Dr. Vikhe Patil Institute of Medical Sciences, Ahmednagar

**Prof. & Registrar MUHS, Nashik

***Dr. Shankarrao Chavan Govt. Medical College, Nanded

Article Info

Article history:

Received May 16, 2012

Revised May 28, 2012

Accepted June 4, 2012

Keyword:

Alpha-tocopherol

Chronic bronchitis

Malondialdehyde

Total antioxidant capacity

ABSTRACT

Lipid peroxide plays an important role in inflammatory lung diseases. Increased epithelial permeability produced by cigarette smoke is likely to be mediated through depletion of the Total Antioxidant Capacity. Oxidative stress has been recognized as a central feature of smoke induced chronic bronchitis. Imbalance between oxidants and Total Antioxidant Capacity is also an established fact in these patients. 60 patients with chronic bronchitis included in the study. Their base line clinical examination, malondialdehyde (MDA), nitric oxide, alpha tocopherol and Total Antioxidant Capacity were measured. 100 healthy non-smokers' were served as controls. The mean malondialdehyde levels and nitric oxide in the patients at base line were higher than Controls ($p < 0.001$). Plasma alpha-tocopherol and total antioxidant capacity were lower ($p < 0.001$) in the patients compared to controls. The present study shows that initially the plasma lipid peroxide (MDA) levels were high and antioxidants (alpha-tocopherol, total antioxidant capacity) were low in patients with chronic bronchitis. Our results suggest the presence of oxidative stress and decrease in total antioxidant capacity in chronic bronchitis.

Copyright © 2012 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Dr. Mrs. Anita M. Raut

Dept of Biochemistry

Dr. Vikhe Patil Institute of Medical Sciences

Ahmednagar (M S), India

Email: anitaraut009@gmail.com

M-9270019213, 8421753938

1. INTRODUCTION

Lung is the organ which is constantly exposed to many atmospheric pollutants such as cigarette smoke, ozone and nitrogen dioxide and is also at risk from oxidant injury by inhalation [1]. Inhaled ozone induces toxic processes that impair lung function. Lipid peroxide plays an important role in inflammatory lung diseases. Increased epithelial permeability produced by cigarette smoke is likely to be mediated through depletion Total Antioxidant Capacity [2]. Oxidant-antioxidant balance is essential for the normal lung function. Both an increased oxidants and or decreased antioxidants may reverse the physiologic oxidant-antioxidants balance in favors of oxidants leading to lung injury.

Chronic bronchitis is defined clinically that it is present in any patient who has persistent cough with sputum production for at least two consecutive years, in the absence of any other identifiable cause. In simple chronic bronchitis, patients have a productive cough but no physiologic evidence of airflow obstruction. Where some individual may demonstrate hyperreactive airways with intermittent bronchospasm and wheezing. This condition is called chronic asthmatic bronchitis some patients, especially heavy smokers,

develop chronic airflow obstruction usually with evidence of associated emphysema and are classified as obstructive chronic bronchitis.

The earliest feature of chronic bronchitis is hyper secretion of mucus in the large airways associated with hypertrophy of the submucosal glands in the trachea and bronchi. As chronic bronchitis persist there is also marked increase in goblet cells of small airways-small bronchi and bronchioles-leading to excessive mucus production that contributes to airway obstruction [3].

Now a days attempt towards oxidative stress status is continuing. The study of antioxidant capacity in lung disease patients opens a promising field in prevention of oxidative stress related complications in these patients. The aims and objectives are: to explore the existence of possible peroxidative damage in lung disease patients by estimating the level of serum malondialdehyde as an index of lipid peroxide; to estimate nitric oxide as a marker of oxidative stress; to study possible alteration in Total antioxidant Status in lung disease patients by estimating the total antioxidant capacity; and to study non enzymatic antioxidant vitamin E.

2. MATERIALS AND METHODS

The present study was conducted in the department of Biochemistry Dr. Vikhe Patil Medical College and Hospital Ahmednagar. This study included 60 clinically stable COPD Patients in the age group of 35-60 years, 100 healthy controls who were diagnosed by physicians on the basis of detailed clinical history, clinical examination and relevant biochemical examinations.

Hypertension, malignancy, overt cardiac failure recent surgery, severe endocrine, hepatic or renal diseases and lung disorders other than COPD were excluded from the present study. Informed consent was obtained from each patient in the study. The study was cleared by institutional ethics committee. 10 ml blood was collected from each patient. Serum was separated by centrifugation at 3000 rpm for 10 minutes at room temperature. Following parameters were carried out on the samples on the same day of collection:

1. The level of serum total lipid peroxide in terms of Malondialdehyde (MDA) was determined by Kei Satoh method [4].
2. Serum Nitric oxide (NO^{*}) as nitrite was estimated by Najwa Cortas and Nabil Wakid method [5]
3. Serum Vitamin 'E' (α-Tocopherol) was estimated by the method of Baker and Frank [6]
4. 4-Total antioxidant capacity in plasma (TAC) was assayed by FRAP analysis [7]

3. RESULTS AND DISCUSSIONS

Table 1 Illustrate the levels of MDA, NO^{*}, Vitamin E and TAC in the healthy controls and chronic bronchitis patients.

Table 1. Levels of MDA, NO^{*}, Vitamin E and TAC in the healthy controls and chronic bronchitis patients

Test	Healthy Controls n=100	Chronic Bronchitis patients n=60
Sr. MDA (umol/L)	1.66±0.289	4.61±2.7*
Sr. NO [*] (umol/L)	33.15±6.13	33.58±12*
Sr. Vit E (mg/dl)	0.927±0.12	0.32±0.09*
Sr.TAC (umol/L)	1253.12±170.22	354.43±88*

n = number of cases

All values are expressed in mean ± SD

* = Significant when compared with control group

It was carried out using students unpaired' test. Probability values 0.05 were considered as significant. Also data were expressed in mean ± SD form. Table 1, results of the present study showed significant high levels (p< 0.001) of oxidants serum lipid peroxide (MDA) and serum nitric oxide (NO^{*}) were observed as compared to healthy controls. The mean plasma levels of Vit. E (P<0.001) and total antioxidant capacity (TAC) was lower than controls.

Lung cells, in particular alveolar epithelial type II cells, are susceptible to the injurious effects of oxidants. Lungs are continuously exposed to oxidants, either generated endogenously by metabolic reactions or exogenously, such as air pollutants or cigarette smoke and since Cigarette smoking, another environmental hazard, also delivers oxidants and free radicals to the lungs. Cigarette smoke contains many oxidants and free radicals, both in the gas and the tar phase and causes sequestration of neutrophils into the pulmonary microcirculation and accumulation of macrophages in respiratory bronchioles. Once recruited, these cells

become activated and generate ROS. ROS, which may also be released by lung epithelial cells, may also stimulate inflammatory cells directly, thereby amplifying lung inflammatory and oxidant events. There by increases the MDA significantly in chronic bronchitis patients [8].

In the respiratory tract, NO[•] is generated enzymatically by all three distinct isoforms of NO[•] synthase i.e. NOS-1, NOS-2 and NOS-3. Of these three forms NOS-2 activity is primarily regulated transcriptionally and is commonly induced by bacterial products and pro-inflammatory cytokines. Inflammatory diseases of the respiratory tract such as chronic bronchitis is commonly characterized by an increased expression of NOS-2 within respiratory epithelial and inflammatory immune cells, and markedly elevated local production of NO[•] in the patients with lung diseases [9].

Vitamin E is the most important lipophilic antioxidant in humans in this study we observed the reduced vitamin E level in lung disease patients could be due partly to its overconsumption as an antioxidant subsequent to increased production of free radicals by cigarette smoke and inflammatory reaction.

The total antioxidative potential of the plasma reflects the ability of an individual to resist the oxidative stress. Ferric reducing ability of plasma (FRAP) evaluates plasma total antioxidant capacity due to known and unknown antioxidants in the plasma [10]. Therefore in the present study significant reduction observed in total ferric reducing ability of plasma may be due to increased free radical activity either because of inflammation or complications that results in imbalance between antioxidant capacity and prooxidant affecting lung function. Extensively amplified oxidant burden and declined individual antioxidant levels might be responsible for the observed significant fall in total antioxidant capacity of patients with lung disease.

4. CONCLUSION

Thus evaluating oxidative stress in lung disease patients by measuring lipid peroxidation and antioxidant status can lead to better understanding of free radical mediated damage in chronic bronchitis patients. An inequity between oxidative stress and antioxidative capacity has been proposed to play an important role in the development and progression of chronic bronchitis and it is related to the severity of disease.

ACKNOWLEDGEMENT

We are grateful to our CEO Dr. Sujay Vikhe Patil and Deputy Director Dr. Abhijit Diwate for his support and encouragement for carrying out this study.

REFERENCES

- [1]. Irfan Rahman and William MacNee. Lung glutathione and oxidative stress implication in cigarette smoke induced airway disease. Am. J. Physiol. Lung cell Mol. Physiol. 279:1999 L1067-L 1088.
- [2]. I Rahman and I. M. Adcock. Oxidative stress and redox regulation of lung inflammation in COPD. Eur. Respir J. 2006, 28; 219-242.
- [3]. Thurbleck W.M., pathology of chronic airflow obstruction. In Cherniack NS (ed) chronic obstructive pulmonary Disease. Philadelphia WB. Saunders 1991: P-
- [4]. Estimation of lipid peroxide (MDA) by Kei satoh method. Clinica Chemica Acta, 1998, 90:37-43.
- [5]. Najwa K. Cortas and W. Wakid. Determination of inorganic nitrate in serum and urine by kinetic cadmium-reduction method. Clin. Chem. 1990, 36:18, 1440-1443.
- [6]. Iris F.F. Benzie and J.J. Strain. The Ferric Reducing Ability of plasma (FRAP) as a Measure of "Antioxidant Power". The FRAP assay Article No. 0292 Analytical Biochem. 239: 1996: 70-76.
- [7]. The Tiet estimation of Serum vitamin 'E' (α -Tocopherol) by method of Baker and Frank. Practical Clinical Biochemistry 1968: Page No. 902.
- [8]. Paul Kirkham and Irfan Rahman. Oxidative Stress in asthma and COPD: Antioxidants as a therapeutic strategy pharmacology and Therapeutics volume 1: issue 2: August 2006: 476-494.
- [9]. DC. Chambers, W.S. Tunnicliffe. Acute inhalation of cigarette smoke increases nitric oxide concentration. Thorax: 1998: 53: 677-679.
- [10]. Aysen Agacodiken, Ilknur Basyigit, Meltem Ozden, Fusun Yildiz, Dilek Ural Hale Maral, Hasim Boyaci, Ahmet Ilgazil and Baki Komsuoglu. The effects of antioxidants on exercise-induced lipid peroxidation in patients with COPD: 24 Feb. 2004: Vol 9 issue 1, p.38-42.

