

## CORALS AS SOURCE OF BACTERIA WITH ANTIMICROBIAL ACTIVITY

Ocky Karna Radjasa<sup>1,2\*</sup>, Jutta Wiese<sup>3</sup>, Agus Sabdono<sup>1,2</sup> and Johannes F Imhoff<sup>3</sup>

Department of Marine Science, Diponegoro University, Semarang - 50275, Central Java, Indonesia  
Center for Tropical Coastal and Marine Studies, Diponegoro University, Widya Puraya, Semarang -  
50275, Central Java, Indonesia

Leibniz-Institute of Marine Sciences, IFM-GEOMAR, Düsternbrooker Weg 20, 24105 Kiel, Germany

Received : February, 22, 2008 ; Accepted : April, 25, 2008

### ABSTRACT

*In this study we examined marine bacteria associated with different corals (Porites lutea, Galaxea fascicularis, Acropora sp. and Pavona sp.) collected from vicinity of Panjang island, Jepara, North Java Sea, Indonesia for their antimicrobial activities against the bacteria Echerichia coli, Bacillus subtilis, Staphylococcus lentus and the yeast Candida glabrata. A total of 13 bacterial isolates belonged to the members of Bacillus, Vibrio, Micrococcus, Pseudoalteromonas, Arthrobacter and Pseudovibrio were found to inhibit the growth of at least one test strain. Further examinations among the biologically active strains by using PCR with specific primers of non-ribosomal peptide synthetase (NRPS) and polyketide synthase (PKS) resulted in the presence of NRPS gene fragments in the 2 members of Bacillus and Micrococcus and PKS gene fragments in the 2 members of Bacillus and Vibrio. Following cloning and sequencing of the PCR products, the fragments from Bacillus BM1.5 and Micrococcus BJB showed sequence identity with peptide synthetase genes of Bacillus subtilis (61 %) and Actinoplanes teichomyceticus (62.4%). On the other hand, PKS-amplifying strains Bacillus BJ.7 and Vibrio MJ.5 showed closest sequence identity with polyketide synthase genes of Bacillus subtilis (73%) and Anabaena sp 90 (62%), respectively.*

**Keywords:** Coral associated bacteria, Antimicrobial activity, PKS, NRPS

\***Correspondence:** Phone: +62-24-7460038; Fax: +62-24-7460039; e-mail : ocky\_radjasa@yahoo.com

## INTRODUCTION

Marine invertebrates that are mainly accumulating within coral reef ecosystems such as soft corals, sponges, tunicates, and bryozoans have long been recognised as the sources of structurally unique natural products. However, it has been suggested that natural products from marine invertebrates have striking similarities to metabolites of their associated microorganisms including bacteria (Proksch *et al.* 2002; Imhoff and Stöhr, 2003). Thus,

it is important to highlight the possible role of bacteria in bacteria-invertebrate associations in the production of biologically active substances.

On the other hand, it has been well known that corals harbor diverse microbial communities (William *et al.* 1987; Shashar *et al.* 1994; Kim, 1994; Santavy *et al.* 1995; Kushmaro *et al.* 1996; Rohwer *et al.* 2001). Every surface immersed in the marine environment, including those organisms

(Armstrong *et al.* 2001) such as mucopolysaccharides-covered coral surfaces providing a nutrient rich habitat for heterotrophic bacteria that leading to the formation of biofilm-forming microbial communities (Kushmaro *et al.* 1997).

Inhibitory interactions among coral-associated bacteria that occur on the coral surface then could be of great interest to search for secondary metabolite-producing bacteria. In addition, examination concernin secondary metabolite-producing bacteria among coral colonizers have been strongly neglected in comparison to soft-bodied marine invertebrates.

Polyketides and non-ribosomal peptides are two of the largest groups of multifunctional proteins that create a multitude of secondary metabolites (Hutchinson, 2003), many of them are used as therapeutic agents *et al.* 2001; Piel *et al.* 2003). Products of the microbial non-ribosomal peptide synthesis include the immunosuppressant cyclosporine and other antibiotics such as gramicin S, tyrocin A and surfactins (Kleinkauf and von Doehren, 1996). Among clinically important polyketides are the antibiotibiotic daunorubicin, erythromycin, lovastatin and rapamycin (Due *et al.* 2001).

With advanced techniques of molecular biology such as polymerase chain reaction (PCR), it is now become possible to carry out a screening on the presence of polyketides and non ribosomal peptides by using specific primers of polyketide synthases (PKS) (Piel, 2002) and non ribosomal polypeptide synthetases (NRPS) (Marahiel *et al.* 1997).

The present study focusses on marine bacteria, which are associated with the corals representing different life forms massive (*Porites lutea*), sub-massive (*Galaxea fascicularis*), branching (*Acropora* sp.) and folious (*Pavona* sp.) and produce secondary metabolites with antimicrobial activity. The examinations are coupled with PCR-based analysis for the presence of polyketide synthases and non-ribosomal

polypeptide synthetases of the bacterial isolates.

## MATERIALS AND METHODS

### *Sampling and isolation of coral-associated bacteria*

Four different corals were collected from the vicinity of Panjang island, Jepara, North Java Sea, Indonesia by scuba diving and identified as *Porites lutea*, *Galaxea fascicularis*, *Acropora* sp. and *Pavona* sp. Upon collection coral fragments were put into sterile plastic bags (Whirl-Pak, Nasco, USA) and immediately brought to the Marine Station of the Diponegoro University where it was rinsed with sterile seawater and scraped off with a sterile knife. The resultant tissues were serially diluted in sterile seawater, spread on Marine Broth, (Difco, Becton Dickenson GmbH, Heidelberg, Germany) and incubated at 20°C for 48 hours. On the basis of morphological features, colonies were randomly picked and purified by making streak plates (Madigan *et al.* 2000).

### *Inhibitory interaction tests*

Inhibitory interaction tests of coral-associated bacteria against test bacteria were performed by using an overlay method. The following bacteria were used: *Escherichia coli* (DSM 498), *Staphylococcus lentus* (DSM 6672), *Bacillus subtilis* (DSM 347), and *Candida glabrata* (DSM 6425) obtained from the German Culture Collection (DSMZ, Braunschweig, Germany).

Overnight cultures of each target microorganism in the logarithmic phase (ca.  $10^9$  cells  $ml^{-1}$ ) were mixed with TSB soft agar medium (TSB 3g, NaCl 10g and Bacto-agar 8g per liter). This mixture (1% v/v, except *C. glabrata* of 10% v/v) was poured on to the respective TSB medium surface, which was previously inoculated with coral-

associated bacteria and incubated for 4 d at 20°C. The plates were then incubated at 20°C for 48 hours. Antimicrobial activity was defined by the formation of inhibition zones around the bacterial colonies.

### ***PCR-based analysis of NRPS and PKS producing bacterial strains***

To obtain genomic DNA of secondary metabolite producing-strains for PCR analysis, cell materials were taken from an agar plate, suspended in sterile water (Sigma, Germany) and subjected to five cycles of freeze (-80°C) and thaw (95°C). Amplification of peptide synthetase gene fragments was carried out with the NRPS degenerated primers A2gamF (5'-AAG GCN GGC GSB GCS TAY STG CC-3') and A3gamR (5'-TTG GGB IKB CCG GTS GIN CCS GAG GTG-3') (Marahiel *et al.*, 1997) and PKS degenerated primers KSDPQQF (5'-MGN GAR GCN NWN SMN ATG GAY CCN CAR CAN MG-3') and KSHGTGR (5'-GGR TCN CCN ARN SWN GTN CCN GTN CCR TG -3') domain (Piel, 2002). All primers were manufactured by MWG-Biotech (Ebersberg, Germany). PCR was performed with an ProgeneThermal cycler (Techne, Burkhardtorf, Germany) as follows: 1 µl template DNA, 1 µl of each of the appropriate primers, and 23 µl DNA free water (Fluka, Sigma-Aldrich Chemie GmbH, Germany) were added to puReTaq Ready-To-Go PCR beads (Amersham Biosciences Europe GmbH, Germany). The NRPS-PCR run comprised 40 cycles with denaturing conditions for 1 min at 95°C, annealing for 1 min at 70°C and extension for 2 min at 72°C, respectively. *Pseudomonas* sp. DSM 50117 was used as positive control. The amplification of PKS gene fragments included an initial denaturing step at 94°C for 2 min, followed by 45 cycles at 94°C for 1 min, annealing at 55°C for 1 min and elongation at 72°C for 2 min. *Bacillus subtilis* 168 as utilized for positive control.

### ***PCR amplification and sequencing of 16S rRNA gene fragments.***

PCR amplification of partial 16S rRNA gene of active strains, purification of PCR products and subsequent sequencing analysis were performed according to the method of Thiel and Imhoff (2003). The almost complete 16S rDNA sequences of strains were compared for homology to the NCBI GenBank and EMBL databases using BLAST (Basic Local Alignment Search Tool) and FASTA searches (Altschul *et al.*, 1990, Pearson 1990). Sequences were aligned using the program ClustalX Version 1.83 (Thomson *et al.* 1997). For phylogenetic calculations the PhyML software (Guindon and Gascuel, 2003) as well as the online version of PhyML (Guindon *et al.*, 2005) were used. Trees were calculated by Maximum Likelihood (ML) method (Felsenstein, 1981) using the GTR model and estimated proportion of invariable sites as well as the Gamma distribution parameter.

### ***Cloning and sequencing of (putative) peptide synthetase and polyketide domains***

The amplified PCR-products were purified using the High Pure PCR Product Purification Kit (Roche Diagnostics, Mannheim, Germany) following the manufacturers protocol. The Big Dye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems) was used for subsequent sequencing on an ABI 310 analyzer (Perkin Elmer Applied Biosystems, Foster City, USA).

### ***DNA sequence accession numbers***

The 16S rRNA gene sequences have been entered into the GenBank database under the sequence accession number AY000000-, the putative peptide synthetase and polyketide synthase sequences obtained under AM287210-AM287213.

## RESULTS AND DISCUSSION

### Results

#### *Inhibitory interaction test*

To estimate antimicrobial activity of coral associated-bacteria, inhibitory interaction tests between test strains and coral associated isolates were carried out. As shown in Table 1, a total of 13 bacterial isolates belonging to the genera *Arthrobacter* (1 strain), *Bacillus* (7 strains), *Micrococcus* (1 strain), *Pseudoalteromonas* (1 strain), *Pseudovibrio* (1 strain), and *Vibrio* (2 strain), were successfully for their inhibitory effect against at least 1 test strain. All 13 coral derived strains inhibited the growth of the Gram-positive bacteria *B. subtilis*. All isolates but one *Pseudovibrio* isolate inhibited the second Gram-positive test strain, *S. lentus*. Two *Bacillus* strains showed an antibacterial activity against the Gram-negative test strain *E. coli*, and one *Bacillus* isolate showed an antifungal activity against the yeast *C. glabrata*.

#### *Detection of NRPS and PKS gene fragments by PCR*

PCR-based analysis using specific primers of non-ribosomal peptide synthetase (NRPS) and polyketide synthase (PKS) revealed that the coral-associated bacterial strains *Bacillus* sp BM1.5 and *Micrococcus* sp BJB showed a PCR amplicon for NRPS. In addition, two other bacterial strains (*Bacillus subtilis* BJ.7 and *Vibrio coralliitycus* MJ.5) exhibited a PCR product of PKS..

#### *Putative peptide synthetase and polyketide sequences.*

Two PKS-amplifying bacterial strains (BJ.7 and MJ.5) had homologies with polyketide synthase genes from *Bacillus subtilis* and *Anabaena* sp 90 (Table 2), meanwhile the other 2 NRPS-amplifying strains (BM1.5 and BJB) showed similarity with peptide

synthetase genes of *Bacillus subtilis* and *Actinoplanes teichomyceticus*, respectively (Table 3).

### Discussion

In a study aimed at the search for antimicrobial compounds from the sea, we investigated bacteria associated with corals from North Java Sea, Indonesia. Our attention was focused on the examination of coral associated-bacteria by using growth inhibition tests followed by PCR-based approach for the occurrence of gene fragments of non-ribosomal peptide synthetase (NRPS) and polyketide synthase (PKS) which are believed to be responsible for the biosynthesis of two main family of secondary metabolites, polyketides and peptides.

Microbes including bacteria play crucial ecological roles in many marine ecosystems such as tropical coral reefs. Unfortunately, studies on bacterial associations with corals have been much less extensive (Knowlton and Rohwer, 2003). Considering the intense interaction among coral colonizers, in particular for the competition for space and nutrient (Slattery *et al.* 2001), coral associated-bacteria have been considered as potential sources of biologically active compounds (Moore, 1999).

In our study, we show that the corals *Porites lutea*, *Galaxea fascicularis*, *Acropora* sp. and *Pavona* sp. offered promising potency as the source of coral associates with biological activities against *E. coli*, *S. lentus*, *B. subtilis* and *C. glabrata*. A total of 13 bacterial isolates belonging to Gram-positive genera, like *Bacillus*, *Micrococcus*, and *Arthrobacter*, and to Gram-negative genera, like *Pseudoalteromonas*, *Pseudovibrio* and *Vibrio* were found to inhibit the growth of at least 1 test strain.. It is interesting to note that 7 out of 13 active strains belonged to the genus *Vibrio* that were isolated from corals *Acropora* sp (2), *G. fascicularis* (3) and *Pavona* sp(2) but not

from *P. lutea*. Furthermore, *Acropora* sp was found to dominate the existence of active strains including *Bacillus* (2), *Micrococcus* (1), *Arthrobacter* (1) and *Pseudovibrio* (1).

One isolate obtained from the coral *Pavona* sp. showed closest similarity to *Pseudoalteromonas* sp. The genus of *Pseudoalteromonas* is widely distributed in the marine environment (Alexeeva *et al.* 2003) and considered one of the most abundant group of marine  $\gamma$ -proteobacteria. For the survival in such environments, the members of genus *Pseudoalteromonas* have been known to develop metabolic pathways that produce biologically active metabolites. Extensive studies regarding the capability of *Pseudoalteromonas* to produce diverse secondary metabolites were reported (Austin, 1989; Jensen and Fenical, 1994; Holmstrom and Kjelleberg, 1999). Further work by (Reid *et al.* 1993; Deng *et al.* 1995) showed that *Pseudoalteromonas luteoviolacea*, owns a non-ribosomal peptide synthetase, which produces the siderophore alterobactin. Radjasa *et al.* (2007) reported the antibacterial activity of a bacterium *Pseudoalteromonas luteoviolacea* TAB4.2 isolated from coral *Acropora* sp. collected from the neighboring sampling site in the North Java Sea. A novel antimicrobial protein was also reported from a marine bacterium strain X153 which was closely related to *P. piscicida* (Longeon *et al.* 2004). Furthermore, Sobolevskaya *et al.* (2005) successfully isolated the recently described *Pseudoalteromonas maricaloris* KMM 635<sup>T</sup> collected in the Coral Sea at a depth of 10 m that produces a brominated cyclic depsipeptides.

Hutchinson (2003) mentioned that microorganisms, especially ones that inhabit marine environments are noted for the ability to produce a wide range of chemicals known as secondary metabolites devoted to the property of self-defense, intercellular communication and other aspects of microbial life. Du *et al.* (2001) mentioned that polyketides and non-ribosomal peptides represent the largest group of natural

products many of which are clinically important drugs. For instance, non-ribosomal peptides including cyclosporin, penicillin and vancomycin are synthesized by non-ribosomal peptide synthetases (NRPS) (Kleinkauf and von Doehren, 1990). On the contrary, polyketides such as daunorubicin, erythromycin, lovastatin and rapamycin are derived from sequential condensation of short carboxylic acids.

In the present study, however, only 2 strains amplified NRPS and 2 other strains amplified PKS gene fragments *Bacillus* sp BM1.5 showed closest similarity to the genes for peptide synthetase and protein building protein of *Bacillus subtilis*. Toh *et al.* (2004) reported the occurrence of emetic toxin of a foodborne pathogen, *Bacillus cereus* that is putatively a product of non-ribosomal peptide synthesis. The presence of NRPS gene fragment in coral bacterium *Bacillus* BM1.5 supported the fact that other *Bacillus* species produce peptides using non-ribosomal peptide synthesis (Stachelhaus and Marahiel, 1995). Meanwhile another bacterium associated with coral *Acropora* sp, *Bacillus subtilis* BJ.7 was found to amplify the gene fragment of PKS and showed sequence similarity of 73% to previously reported polyketide gene of *Bacillus subtilis* (acc. no. U11039). Scotti *et al.* (1993) studied the potential ability of *B. subtilis* to synthesize polyketides.

*Micrococcus* sp BJB, had sequence similarity to peptide synthetase of *Actinoplanes teichomyceticus*. The members of *Micrococcus* are heterotrophic bacteria distributed in various environments such as seawater (Tanaka *et al.* 2001), marine sediments (Zhong *et al.* 2002) and prawn-rearing water (Phatarpekar *et al.* 2002). Members of the genus *Micrococcus* have also the potency as producers of antimicrobial compounds. A marine bacterium *Micrococcus* MCCB 104 isolated from hatchery water demonstrated extracellular antagonistic properties against *Vibrio alginolyticus*, *V. parahaemolyticus*, *V. fluvialis*, *V. nereis*, *V. proteolyticus*, *V.*

*mediterranei*, *V. cholerae* and *Aeromonas* sp (Jayaprakash *et al.* 2005). Variacin, a bacteriocin produced by *Micrococcus varians* inhibited other Gram-positive bacteria, but not Gram-negative ones (Pridmore *et al.*, 1996).

Two antibacterial active isolates of vibrios, *V. coralliitycus* MJ.5 and *V. parahaemolyticus* MJ.11 were obtained from coral *Porites lutea*. Isolate *V. coralliitycus* MJ.5 also amplified the gene fragment of PKS and had sequence similarity (62%) to polyketide synthase of *Anabaena* sp90. Polyketide natural products are common metabolites of blue-green algae (cyanobacteria) (Burdja *et al.* 2001). Neilan *et al.* (1999) mentioned that *Cyanobacteria* produced a myriad array of secondary metabolites, including alkaloids, polyketides, and non ribosomal peptides, some of which are potent toxins. Isolate *V. parahaemolyticus* MJ.11 inhibited the growth of *Bacillus subtilis* and *Staphylococcus lentus*. *V. parahaemolyticus* has been reported as the member of *Vibrio*, a dominant genus responsible for the observed mortality in prawn-farmings (Lightner, 1988). As far as known it is uncertain, if *V. parahaemolyticus* is also a pathogen of corals. Cervino *et al.* (2004) considered that further studies should include the examination to determine if strains of *V. parahaemolyticus* and *V. alginolyticus* are capable of inducing disease signs in *Montastraea* spp.

## CONCLUSION

Our study on coral-associated bacteria provides evidence of antimicrobial properties among coral colonizers. Although in minor numbers, the coral-associated bacteria might be capable of producing polyketides and peptides, because of the detection of NRPS and PKS gene fragments as well as their antimicrobial activities. The discovery of the potential for the synthesis

of polyketides and peptides among coral-associated bacteria should be of interest because of promising applications in the development of pharmacological polyketides and peptides. Furthermore, considering the fact that research on coral-bacteria association has been fairly limited, the future study on the search for secondary metabolite producers among coral colonizers should be given prominence.

## ACKNOWLEDGEMENTS

This work was partly supported by the grant from Ministry of Research and Technology, Indonesia under the competitive research grant scheme (RUT XI) No. 011.27.SK.RUT.2004. The work was also part of a research grant provided by German Center of Excellence within *BIOTECmarin*. Ocky Karna Radjasa was research fellow of the Alexander von Humboldt Foundation (AvH) within the Georg Forster Research Fellowship Programme.

## REFERENCES

- Alexeeva YV, Kalinovskaya NI, Kuznetsova TA, Ivanova EP 2003 Impact of cultivation conditions on haemolytic activity of *Pseudoalteromonas issachenkonii* KMM 3549<sup>T</sup>. *Lett. Appl. Microbiol* 38:38-42
- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ 1990 Basic local alignment search tool. *J Mol. Biol* 215:403-410
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman D J 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids.Res* 25:3389-3402

- Armstrong E, Yan L, Boyd KG, Wright PC, Burgess JG 2001. The symbiotic role of marine microbes on living surfaces. *Hydrobiologia* 461: 37-4
- Cane DE 1997 Polyketide and non-ribosomal polypeptide biosynthesis. *Chem.Rev* 97: 2463-2706
- Cervino JM, Hayes RL, Polson SW, Polson SC, Goreau TJ, Martinez SJ, Smith GW. 2004. Relationship of *Vibrio* species infection and elevated temperatures to Yellow Blotch/Band disease in Caribbean corals. *Appl. Environ.Microbiol.* 70: 6855-6864
- Christiansen G, Dittmann E, Ordorika IV, Rippka R, Herdman M, Borner T 2001. Nonribosomal peptide synthetase genes occur in most cyanobacterial genera as evidenced by their distribution in axenic strains of PCC. *Arch. Microbiol* 176:452-458
- Coffroth MA 1990. The function and fate mucous sheets produced by reef coelenterates. *Procc The 6<sup>th</sup> Int Coral Reef Symp* 2:15-20. Australia
- Coll JC, Sammarco PW 1986. Soft corals: chemistry and ecology. *Oceanus* 29:33-37
- Deng J G, Hamada Y, Shioiri T 1995. Total synthesis of alterobactin-A, a super siderophore from an open-ocean bacterium. *J Am Chem Soc* 117 (29): 824-7825
- Du L, Sanchez C, Shen B 2001. Hybrid peptide-polyketide natural products: biosynthesis and prospects toward engineering novel molecules. *Metabol. Engineer* 3:78-95
- Faulkner DJ 2000 Marine pharmacology. *Antonie. Van. Leeuwenhoek.* 77: 135-145
- Felsenstein J 1981 Evolutionary trees from DNA-sequences - a maximum-likelihood approach. *J. Mol. Evol* 17:368-376
- Fenical W 1993 Chemical studies of marine bacteria: developing a new source. *Chem. Rev* 93:1673-1683
- Guindon S, Lethiec F, Duroux P, Gascuel O 2005 PHYML Online - a web server for fast maximum likelihood-based phylogenetic inference. *Nucleic. Acids. Res* 33:W557-W559
- Guindon S, Gascuel O 2003 A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol* 52:696-704
- Jensen P.R, Fenical W 2000 Marine microorganisms and drug discovery: current status and future potential. In: Fusetani N (ed). *Drugs from the sea*, Basel: Karger, pp 6-29
- Kim K. 1994. Antimicrobial activity in gorgonian corals (Coelenterata, Octocorallia). *Coral. Reefs* 13:75-80
- Kiorboe T, Grossart HP, Ploug H, Kam T 2003. Microbial dynamics on particles: colonization, growth, detachment, and grazing mortality of attached bacteria. *Appl. Environ. Microbiol* 69:3036-3047.
- Kleinkauf H, von Doehren H 1996. A nonribosomal system of peptide biosynthesis. *Eur. J. Biochem* 236:335-351
- Knowlton N., Rohwer F 2003. Multispecies microbial mutualisms on coral reefs: the host as a habitat. *Am. Nat* 162:51-62

- Kushmaro A, Loya Y, Fine M, Rossenberg E 1996 Bacterial infection and coral bleaching. *Nature* 380:396
- Kushmaro A, Rossenberg E, Fine M, Loya Y, 1997. Bleaching of the coral *Oculina patagonica* by *Vibrio* AK-1. *Mar. Ecol. Prog. Ser* 147 159-165
- Lightner, DV. 1988. *Vibrio* disease of penaeid shrimp, p. 42–47. In C. J. Sindermann and D. V. Lightner (ed.), Disease diagnosis and control in North American marine aquaculture. Elsevier, Amsterdam. The Netherlands.
- Longeon A, Peduzzi J, Barthelemy, Corre S, Nicolas J-L, Guyot M 2004. Purification and partial identification of novel antimicrobial protein from marine bacterium *Pseudoalteromonas* species strain X153. *Mar. Biotechnol* 6:633-641
- Long R , Azam F 2001 Antagonistic interactions among marine pelagic bacteria. *Appl. Environ. Microbiol* 67:4975-4983
- Ludwig W, Strunk O, Klugbauer S, Klugbauer N, Weizenegger M, Neumaier J, Bachleitner M, Schleifer KH 1998. Bacterial phylogeny based on comparative sequence analysis. *Electrophoresis* 19:554-68
- Madigan MT, Martinko JM, Parker J, Brock TD, 2000. Biology of microorganisms. Prentice-Hall, Inc., Upper Saddle River, New Jersey 07458
- Marahiel MA, Stachelhaus T, Mootz HD 1997. Modular peptide synthetases involved in nonribosomal peptide synthesis. *Chem. Rev.* 97:2651-2673
- Messing J 1983. New M13 vectors for cloning. *Methods. Enzymol* 10:20-78
- Moore BS 1999. Biosynthesis of marine natural products: microorganisms and macroalgae. *Nat. Prod. Rep* 16:653-674
- Munro MHG, Blunt JW, Dumdei EJ, Hickford SJH, Lill RE, Li S, Battershill CN, Duckworth AR 1999. The discovery and development of marine compounds with pharmaceutical potential. *J. Biotechnol* 70:15-25
- Neilan BA, Dittmann, Rouhiainen L, Bass RA, Schaub V, Sivonen K, Borner T 1999. Nonribosomal peptide synthesis and toxigenicity of cyanobacteria. *J. Bacteriol* 181:4089-4097
- Paul JH, DeFlaun ME, Jeffrey WH 1986 Elevated levels of microbial activity in the coral surface microlayer. *Mar. Ecol. Prog. Ser.* 33:29-40
- Pascal H, Vacelet E 1981. Bacterial utilization of mucus on the coral reef of Aqaba (Red Sea). *Procc. 4<sup>th</sup>. Intl Coral Reef Symp.* Manila: pp 669-677
- Pearson WR 1990. Rapid and Sensitive Sequence Comparison with FASTP and FASTA. *Methods. Enzymol.* 183:63-98
- Piel J 2002. A polyketide synthase-peptide synthetase gene cluster from an uncultured bacterial symbiont of *Paederus* beetles. *PNAS* 29:14002-14007
- Piel J, Hui D, Fusetani N, Matsunaga S 2003. Targeting modular polyketide synthetase with iteratively acting acyltransferases from metagenomes of uncultured bacterial consortia. *Environ. Microbiol.* 6: 921-927
- Pietra F, 1997. Secondary metabolites from marine microorganisms- bacteria,

- protozoa, algae and fungi-achievements and prospectives. *J. Nat. Prod.* 14: 453-464
- Proksch P, Edrada RA, Ebel R. 2002. Drugs from the seas-current status and microbiological implications. *Appl Microbiol Biotechnol.* 59: 125-134
- Radjasa, O.K., T. Martens., H-P. Grossart., T. Brinkoff., A. Sabdono., and M. Simon. 2007. Antagonistic activity of a marine bacterium *Pseudoalteromonas luteoviolacea* TAB4.2 associated with coral *Acropora* sp. *J. Biol. Sci.* 7(2):239-246
- Reid RT, Live DH, Faulkner DJ, Butler A 1993. A siderophore from a marine bacterium with an exceptional ferric ion affinity constant. *Nature.* 366:455-458
- Rohwer F, Breitbart M, Jara J, Azam F, Knowlton N 2001. Diversity of bacteria associated with the Caribbean coral *Montastraea franksi*. *Coral. Reefs.* 20:85-95
- Rohwer F, Seguritan V, Azam F, Knowlton N 2002 Diversity and distribution of coral-associated bacteria. *Mar. Ecol. Prog. Ser.* 243:1-10
- Rublee PA, Lasker HR, Gottfried M, Roman MR 1980. Production and bacterial colonization of mucus from the soft coral *Briararium asbestinum*. *Bull. Mar. Sci.* 30 : 888-893
- Sammarco PW, Coll JC 1992. Chemical adaptation in the Octocorallia: Evolutionary considerations. *Mar. Ecol. Prog. Ser* 88: 93-104
- Santavy DL, Peters EC, Kozlowski J, Wilkinson S 1995. Characterization of the bacterium suspected in the incidence of white band disease. *Abstr. Gen. Meet. Am. Soc. Microbiol.* p. 332
- Scotti C, Piatti M, Cuzzoni A, Perani P, Tognoni A, Grandi G, Galizzi A, Albertini AM 1993. A *Bacillus subtilis* large ORF coding for a polypeptide highly similar to polyketide synthases. *Gene* 16:65-71.
- Scudiero DA, Shoemaker RH, Paull KD, Monks A, Tierney S, Nofziger TH, Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic. Acids. Res.* 25:4876-4882
- Shashar, N., Cohen, Y., Ioya, Y., and Sar, N. 1994. Nitrogen fixation (acetylene reduction) in stony corals: evidence for coral-bacteria interactions. *Mar. Ecol. Prog. Ser.* 111: 259-264.
- Silakowski B, Nordsiek G, Kunze B, Blöker H, Müller R 2001. Novel features in a combined polyketide synthase/non-ribosomal peptide synthetase: the myxalamid biosynthetic gene cluster of the myxobacterium *Stigmatella aurantiaca* Sga15. *Chem. Biol.* 8:59-69
- Slattery M, Rajbhandari I, Wesson K 2001. Competition - mediated antibiotic induction in the marine bacterium *Streptomyces tenjimariensis*. *Microb. Ecol.* 41:90-96
- Sobolevskaya MP, Smetanina OF, Speitling M, Shevchenko LS, Dmitrenok PS, Laatsch H, Kuznetsova TA, Ivanova EP, Elyakov GB (2005) Controlling production of brominated cyclic depsipeptides by *Pseudoalteromonas*

- maricaloris* KMM 636<sup>T</sup>. *Lett. Appl. Microbiol.* 40:243-248
- Strunk O, Gross O, Reichel B, May M, Hermann S, Stuckmann N, Nonhoff B, Lenke M, Ginhart A, Vilbig A, Ludwig T, Bode A, Schleifer K-H, Ludwig W 1998 ARB: a software environment for sequence data. <http://www.mikro.biologie.tu-muenchen.de/pub/ARB>. Department of Microbiology, Technische Universität München, Munich, Germany.
- Thiel V, Imhoff JF 2003 Phylogenetic identification of bacteria with antimicrobial activities isolated from Mediterranean sponges. *Biolmol. Eng.* 20:421-423
- Toh M, Moffitt MC, Henrichsen L, Raftery M, Barrow K, Cox JM, Marquis CP, Neilan BA 2004. Cereulide, the emetic toxin of *Bacillus cereus*, is putatively a product of nonribosomal peptide synthesis. *J. Appl. Microbiol.* 97:992-1000
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic. Acids. Res.* 25:4876-4882
- Williams WM, Viner AB, Broughton WJ 1987. Nitrogen fixation (acetylene reduction) associated with the living coral *Acropora variabilis*. *Mar. Biol* 94:531-535