



## International Journal of Science and Engineering(IJSE)

Home page: <http://ejournal.undip.ac.id/index.php/ijse>



# The Synthesis Biodiesel from Palm Oil Through Interesterification Using Immobilized Lipase Enzym as Catalyst

### The Effect of Amount of Biocatalyst, Mole Ratio of Reactan, Temperature to Yield

Renita Manurung<sup>1</sup>, Melina Widyawati<sup>2</sup>, Ricky Afrianto<sup>3</sup>  
Chemical Engineering Department, Engineering Faculty, University of Sumatera Utara<sup>1,2,3</sup>  
Jl. Almamater Kampus USU Medan 20155, Indonesia

renitachem@yahoo.com<sup>1</sup>, winzseasonz@yahoo.co.id<sup>2</sup>, rickyafrianto@yahoo.com<sup>3</sup>

**Abstract** - Biodiesel usually synthesized by transesterification of triglyceride and alcohol by addition of acid or base catalyst so there is could produce a waste of chemical process. Alternative process is by using biocatalyst such as enzyme to synthesize biodiesel without chemical process waste. In this research, synthesis of biodiesel from Crude Palm Oil (CPO) that through the process of degumming and methyl acetate as acyl donor has been investigated with using Lipozyme as biocatalyst. Variables in this research are amount of biocatalyst, mole ratio of reactant, and temperature, and its respond to the yield conversion of biodiesel that presented by using Response Surface Methodology (RSM). Yield raging from 15% - 68% were achieved during 10 hours reaction time. The results showed that the most influential variable is amount of biocatalyst.

**Keywords**—Biodiesel, Methyl acetate, CPO, lipozyme, RSM

Submission: June 12, 2014

Corrected: July 30, 2014

Accepted: October 10, 2014

Doi: [10.12777/ijse.7.2.174-177](https://doi.org/10.12777/ijse.7.2.174-177)

**[How to cite this article:** Aditia, M., Sunarso., C.C. Sevilla, and A.A. Angeles (2014). Growth Performance and Mineral Status on Goats

(*Caprahircuslinn.*) Supplemented with Zinc Protein and Selenium Yeast. *International Journal of Science and Engineering*, 7(2), 174-177; Doi: [10.12777/ijse.7.2.174-177](https://doi.org/10.12777/ijse.7.2.174-177)

## I. INTRODUCTION

Since 1990, research and development in biodiesel field had done extensively to obtain the renewable fuel oil. Indonesia has various species of plants produce oil or fat as biodiesel. [1] One of raw material for biodiesel is crude palm oil (CPO). Indonesia is a bigger producer of Crude Palm Oil (CPO) in the world since 2006 with the area of oil palm is 5 million hectare. [2]

Biodiesel is produced by reaction of vegetable oil and alcohol using base as catalyst in certain composition and temperature. [3] But recently, a biodiesel synthesis had be developed using lipase enzyme as biocatalyst. [4] The advantage of enzymatic process is product separation was easier and without produce the waste of chemical process.

Lipase represent soluble enzyme in water and catalyze the hydrolysis reaction of fat substrate ester bind that did not soluble in water and role as interface layer between water and organic phase. Enzymatic action of lipase on substrate is a product of nucleophilic on atom of carbonyl carbon from ester group. Some lipase also able to catalyze the esterification,

interesterification, transesterification, acidosis, aminolysis processes and indicates enantioselectivity character. [5]

For industrial application, specificity of lipase is an important factor. This enzyme will present specificity of substrate (fat acid or alcohol) include the isomer differentiation. Lipase can be divided into 3 groups based on their specificity, i.e. non specific lipase, 1,3-specific lipase and fatty acid lipase. [6]

The using of enzyme independently for product of biodiesel production has any technical limitation and unreliable practically because it is not recovered and reuse, and will increase the production process cost and increase the contamination of product by remains enzyme. These difficultness can be minimized by using immobilized enzyme that enable reuse of biocatalyst in anytime, minimize the cost and increase the quality of product.

The using of methanol and ethanol in biodiesel synthesis produce the glycerol as by product that could block the active side of lipase enzyme. Therefore, the using of alternative acyl

group donor (non alcohol route) such as methyl acetate, ethyl acetate and propan-2-ol, had be studied. [6-10]

The synthesis of biodiesel through non alcohol route is classified into transesterification reaction in which transesterification can be depict as group change between two ester by the presence of catalyst. [11]




Figure 1. Interesterification Reaction by Methyl Acetate

## II. MATERIAL AND METHOD

The main material used in this research such as CPO is supplied by PT. Perkebunan Nusantara IV Indonesia, methyl acetate and phosphate acid from Merck and Lipozyme RM IM from Sigma Aldrich.

The analysis of fatty acid composition of CPO and the same product as FAME is using Chromatography gas method (Shimadzu GC 148 by FID detector, column type of DB-1HT; 1.5 mm x 0.25 mm ID, film thick is 0.1 μm, carrier gas; helium, flushing gas; nitrogen, oven temperature 50 °C, injector temperature 400 °C, detector temperature is 400 °C).

Degumming procedure of CPO using phosphate acid 0.6% (w/w) on temperature of 60 °C. Determining of FFA content on CPO is using AOCS Official method Ca 5a-40 before and after degumming. Procedure of transesterification reaction is the degumming CPO was reacted to methyl acetate during 10 hours in 150 rpm with molar ratio 1:4 – 1 :9, on temperature 45 – 60 °C by 10-30% (w/w) biocatalyst using erlenmeyer in heater shaker. Analysis of physical characteristic of biodiesel is using OECD 109 method for density and ASTM D 445 method for kinematic viscosity.

## III. RESULT AND DISCUSSION

### A. The Analysis of Crude Palm Oil (CPO)

This research was conducted by using Crude Palm Oil (CPO) as raw material that had been degumming. Degumming is a separation process gum that consists of phospholipids, protein, residue, carbohydrate, water and resin. [12]. The content of Free Fatty Acid (FFA) content in CPO before and after degumming process is shown in Figure 2.




Figure 2. Analysis of FFA Content in CPO Before and After Degumming

Figure 2 indicates that there is decreasing FFA of CPO after degumming for 14.33%. The decreasing of FFA content means the increasing of enzyme performance as a

consequence for the decreasing of content and number of pollutant such as gum that could block the porous and active side of enzyme. Previously, there is an introduction study using CPO as raw material without degumming and the biodiesel yield is 16.05%, in which this yield is smaller than using degumming CPO as raw material. Based on this condition, the degumming process must be conducted as a pretreatment in using CPO as biodiesel raw material in enzymatic process. On vegetable oil and fat, saturated fatty acid is found on external position of sn-1 and sn-3 and unsaturated fatty acid on inner side of sn-2. [13]



Figure 3. Interesterification Reaction of Triacylglycerol using Lipase sn-1,3 Specific (A,B, C, X = fatty acid/acyl group) [14]

Composition of saturated and unsaturated fatty acid showed in table 1

Composition	Percentage (%)
Saturated Fatty Acid	39.2172
Unsaturated fatty acid	60.7827

In this research, it use immobilized lipase enzyme using support of porous ion exchange resin (Lipozyme RM IM). Lipozyme RM IM is a biocatalyst in specificity sn-1,3 that release the fatty acid from position 1 and 3 of glycerida. [6] By using lipase specific sn-1,3 on transesterification reaction, exchange a half of acyl group is focus to sn-1 and sn-3 positions that increase the product by characteristic that did not found from transesterification chemically. [15]

Based on composition of saturated and unsaturated fatty acid in CPO, it is possible that did not less than 39.2172% fatty acid will converted to be ester using Lipozyme. Because the dominant fatty acid in CPO is unsaturated fatty acid for 60.7827% in sn-2 position, the using of non specific enzyme could produces a best yield.

### B. Analysis of Experimental Variable

The influence of used experiment variable is processed statistically and presented in table 2.

Table 2. Estimation of Statistic Equation Model Parameter

Term	Coef	SE Coef	T	P
Constant	22,727	2,102	10,813	0,000
Amount of Biocatalyst (X <sub>1</sub> )	10,679	1,395	7,658	0,000
Mole Ratio of Reactan (X <sub>2</sub> )	6,254	1,395	4,485	0,001
Temperature (X <sub>3</sub> )	1,713	1,395	1,228	0,248
X <sub>1</sub> *X <sub>1</sub>	8,912	1,358	6,565	0,000
X <sub>2</sub> *X <sub>2</sub>	3,148	1,358	2,319	0,043
X <sub>3</sub> *X <sub>3</sub>	6,852	1,358	5,047	0,001
X <sub>1</sub> *X <sub>2</sub>	1,240	1,822	0,681	0,512
X <sub>1</sub> *X <sub>3</sub>	-5,678	1,822	-3,116	0,011
X <sub>2</sub> *X <sub>3</sub>	-1,965	1,822	-1,079	0,306
S = 5.153	R-Sq = 94.0%	R-Sq(adj) = 88.5%		

By using analysis of surface response methodology with coded level, there is a correlation of %yield and the three variables, i.e.

$$Y = 22,727 + 10,679 X_1 + 6,254 X_2 + 1,713 X_3 + 8,912 X_1^2 + 3,148 X_2^2 + 6,852 X_3^2 + 1,240 X_1 X_2 - 5,678 X_1 X_3 - 1,965 X_2 X_3 \quad (1)$$




Figure 4. Contour Plot %Yield of Biodiesel for Amount of Biocatalyst vs Mole ratio of reactant

Figure 4 shows that the increasing of amount of biocatalyst has a significant influence to the conversion of yield than mole ratio of reactant in constant temperature 50 °C. Based on Contour Plot it indicates that the increasing of biocatalyst amount will increase the yield significantly.

The number of enzyme is an important variable of operation to achieve a rapid and efficient reaction. But, the increasing of lipase did not produce a higher conversion. [16] While that in reaction with immobilized enzyme as catalyst in which enzyme cannot be interacted so the increasing of enzyme up to certain amount will influence the velocity of reaction positively.

Figure 4 indicates that the bigger yield conversion (>80%) is obtained by increasing the biocatalyst more than 29% and mole ratio of reactant is more than 8.5. Based on statistical analysis of surface respond method on table II it indicates that interaction between the amount of biocatalyst and mole ratio of reactant will have a positive yield for 1.240. While if the amount of biocatalyst and mole ratio of reactant is smaller, the yield conversion is smaller

The higher of mole ratio of reactant, the higher of number of substrate, while the higher of biocatalyst, the highest of active side of enzyme. Therefore, by the increasing of the both of variable will increase the yield conversion. This is caused by the more of interaction between the active side of enzyme that contact to available substrate directly. Based on this research it concluded that interaction between amount of biocatalyst and mole ratio of reactant on certain limit will influence the yield significantly.

Figure 5 shows the increasing of mole ratio of reactant indicates a significant change than by the increasing of temperature with the fixed biocatalyst amount 20%. Contour plot indicates that if mole ratio of reactant is increased by the constant temperature, it increase the yield significantly.

Stoichiometric ratio for transesterification reaction requires 3 mole methyl acetate and 1 mole triglyceride to produce 3 mole methyl ester and 1 mole triacetilglyserol. So the number of methyl acetate in a big number is required to drive the reaction to produce a product.

As reported that if methyl acetate is over it make the oil is more liquid cause the declining of conversion from methyl ester. [17]




Figure 5. Contour Plot of %Yield of Biodiesel for Mole Ratio of reactant vs. Temperature.

Figure 5 shows that a big yield conversion is obtained by increasing the mole ratio of reactant and maintain the permanent temperature in optimum condition.




Figure 6. Contour Plot of % yield of Biodiesel for temperature vs Amount of Biocatalyst.

Figure 6 shows that the increasing of number of biocatalyst has a significant influence to the %yield with fixed variable of mole ratio of reactant 1 : 6. But it is not same to the temperature without a significant influence to %yield. This is caused by deactivated of lipase enzyme in higher temperature so it decrease %yield of biodiesel. Contour plot indicates that if temperature is lower and the number of biocatalyst is increase, it increase the %yield of biodiesel product.

The higher temperature will increase the reaction rate because it minimize the viscosity of lipid compound and increase the transfer between substrate and product in surface or in enzyme particle. But, the higher temperature is also lower the stability and half time of enzyme. [18]

Temperature has an important role in transesterification reaction enzymatically. A research by using Lipozyme TL IM and vegetable oil as raw material in temperature 35 – 38 °C as a higher yield of conversion for 90%. [19] A research by using sunflower seed oil and Novozyme 435 as catalyst produce yield for 99.6% on temperature 45 °C. [20] While for CPO as raw material based on Figure 6, the optimum temperature for Lipozyme is < 45 °C.

Figure 6 shows that a higher yield conversion is obtained by addition of biocatalyst in the lower temperature. It is caused by a higher temperature of reaction will deactivate the performance of lipase enzyme. Therefore it concluded that temperature is not a dominant variable because it did not have a significant influence when interacted to other variables.

### C. Analysis of Physical Characteristic of Biodiesel

The below is a result of density and viscosity analysis of biodiesel as shown in table 3.

Table 3. The Result of Physical Characteristic of Biodiesel

Amount of Biocatalyst (b/b)	Molar Ratio of Reactant	Temperature (°C)	Density (g/ml)	Kinematic Viscosity (cSt)
30 %	1 : 6	50	0.86524	3.517

The result of density and viscosity analysis is suitable to SNI standard, i.e. for density is in range of 0.84 – 0.89 g/ml in temperature 40 °C while for kinematic viscosity is in range of 2.3 – 6.0 cSt in temperature of 40 °C.

## IV. CONCLUSIONS

The performance of Lipozyme that only specific to break down the chain 1 and 3 on triglyceride cause a few of fatty acid will converted to be ester so the using of non specific enzyme will give a best yield. On interesterification of CPO, a dominant variable are the amount of biocatalyst, mole ratio of reactant, and temperature. Temperature is variable that has not significant influence when interacted to the other both factors.

## REFERENCES

- [1] Riyanti, Fahma., Poedji L.H., and Catur D.L. 2012. Pengaruh Variasi Konsentrasi Katalis KOH pada Pembuatan Metil Ester dari Minyak Biji Ketapang (Terminalia catappa Linn). Jurnal Penelitian Sains, Vol. 15, No. 2, 2012. (1-5)
- [2] Pandiangan, Payaman. 2008. Studi Proses Interesterifikasi Enzimatik (EIE) Campuran Minyak Sawit dan Minyak Kelapa untuk Produksi Bahan Baku Margarin Bebas Asam Lemak Trans. Tesis. Sekolah Pascasarjana, Institut Pertanian Bogor. Bogor.
- [3] Yuniwati, Murni., dan Karim, Amelia Abdul. 2009 Kinetika Reaksi Pembuatan Biodiesel dari Minyak Goreng Bekas (Jelantah) dan Metanol dengan Katalisator KOH. Jurnal. Jurusan Teknik Kimia, Institut Sains & Teknologi AKPRIND.
- [4] Hermansyah, Heri., Marno, Septhian., Arbianti, Rita., Utami, Tania Surya., dan Wijanarko, Anondho. 2008. Interesterifikasi Minyak Kelapa Sawit dengan Metil Asetat untuk Sintesis Biodiesel Menggunakan Candida Rugosa Lipase Terimobilisasi. Jurnal Teknik Kimia Indonesia, Vol. 8, No. 1. Departemen Teknik Kimia. Fakultas Teknik. Universitas Indonesia.
- [5] Stoytcheva, M., Montero, G., Toscano, L., Gochev, V., Valdez, B. 2011. The Immobilized Lipases in Biodiesel Production. InTech Journal, Vol. 19, 50-72.
- [6] Ribeiro, Bernardo Dias., Aline Machado de Castro. 2011. Coelho, Maria Alice Zarur., and Denise Maria Guimarães Freire. Production and Use of Lipases in Bioenergy: A Review from the Feedstocks to Biodiesel Production. Journal of Enzyme Research, Volume: 2011, 16 pages.
- [7] Du, W., Xu, Y., Liu D., and Zeng, J. 2004. Comparative Study on Lipase-Catalyzed Transformation of Soybean Oil for Biodiesel Production with Different Acyl Acceptors. Journal of Molecular Catalysis B: Enzymatic, Vol. 30, No. 3-4, 125-129, ISSN 1381-1177.
- [8] Modi, M. K., Reddy J.R.C., Rao, B.V.S.K., and Prasad, R.B.N. 2007. Lipase-Mediated Conversion of Vegetable Oils Into Biodiesel Using Ethyl Acetate as Acyl Acceptor. Bioresource Technology, Vol. 98, No. 6, 1260-1264, ISSN 0960-852.
- [9] Ognjanović, N., Bezbradica, D. & Knežević-Jugović, Z. 2009. Enzymatic Conversion Of Sunflower Oil To Biodiesel In A Solvent-Free System: Process Optimization And Immobilized System Stability. Journal of Bioresource Technology, Vol. 100, No. 21.
- [10] Hermansyah, Heri., Arbianti, Rita., Rizkiyadi, Muhammad Ekky., Surendro., dan Risan Aji. 2012. Interesterification of Fried Palm Oil with Methyl Acetate using Candida rugosa Lipase To Produce Biodiesel. Artikel, Departemen Teknik Kimia, Universitas Indonesia.
- [11] Maulana, Mirza Akbar. 2012. Pemanfaat Whole Cell Candida Rugosa sebagai Biokatalis untuk Sintesis Biodiesel Melalui Rute Non Alkohol. Skripsi. Universitas Indonesia. Depok.
- [12] Lin L, Rhee KC, Koseoglu SS. 1998. Recent Progress in Membrane Degumming of Crude Vegetable Oils on a Pilot-Plant Scale. Journal of Food Protein R&D Center.
- [13] Sidabutar, G.U. 2011. Aktivitas Lipase dari Getah (Carica papaya, Linn.) Terhadap Minyak Lemak. Skripsi. Universitas Sumatera Utara.
- [14] Mukherjee, Kumar D. 1998. Lipase-catalyzed Synthesis of Designer Lipids with Improved Nutritional Properties. Journal of Food, Nutrition and Well Being, Pusaltee 68, D-48147.
- [15] Schuch, Ricardo and Mukherjee, Kumar D. 1987. Interesterification of Lipids Using an Immobilized sn-1,3-specific Triacylglycerol Lipase. Journal of Agricultural and Food Chemistry, 35, 1005-1008.
- [16] Zhong, Nanjing., Zhongyu Gui., Li Xu., Jianrong Huang., Kun Hu., Yongqing Gao., Xia Zhang., Zhenbo Xu., Jianyu Su., and Bing Li. 2013. Solvent-free Enzymatic Synthesis of 1,3-Diacylglycerols by Direct Esterification of Glycerol with Saturated Fatty Acids. Journal Lipid in Health and Disease, 12:65.
- [17] Yuang, H and Yan, Y. 2008. Lipase-Catalyzed Biodiesel Production with Methyl Acetate as Acyl Acceptor. Z Naturforsch C, Mar-Apr;63 (3-4):297-302.
- [18] Xu, X., Fomuso, L., Akoh, C. 2000. Synthesis of Structured Triacylglycerols by Lipase-Catalyzed Acidolysis in a Packed Bed Bioreactor. Journal Agric. Food Chem, 48, 3–10.
- [19] Ferreira, P.J., Sousa, H.S., Caetano, N.S. 2008. Biodiesel Production From Vegetable Frying Oil and Ethanol Using Enzymatic Catalysis. Journal of Bioenergy Challenges and Opportunities.
- [20] Ognjanović, Nevena D., Šaponjić, Svetlana V., and Bezbradica Dejan I. 2008. Lipase-Catalyzed Biodiesel Synthesis with Different Acyl Acceptors. APTEFF, 39, 1-212 (2008) UDC: 665.3:665.75:665.058.6.