

SCREENING APPROACHES FOR METHANE MITIGATING POTENTIAL OF TANNIN-CONTAINING PLANTS UNDER *IN VITRO* RUMEN ENVIRONMENT

A. Jayanegara, M. Ridla and Nahrowi

Department of Nutrition and Feed Technology, Faculty of Animal Science,
Bogor Agricultural University, Jl. Agatis Kampus IPB Dramaga Bogor 16680 - Indonesia
Corresponding E-mail: anuragaja@ipb.ac.id

Received July 17, 2012; Accepted September 06, 2012

ABSTRAK

Tujuan penelitian ini adalah untuk melakukan pemilahan terhadap hijauan mengandung tanin yang memiliki sifat menurunkan emisi gas metana dengan menggunakan pendekatan univariabel, bivariabel dan multivariabel (*principal component analysis*, PCA). Sampel hijauan berasal dari beberapa negara, yakni Indonesia (n = 27 spesies), Mongolia (n = 14), Swiss (n = 16) dan Jerman (n = 3). Hijauan diinkubasi secara *in vitro* dengan cairan rumen-buffer pada suhu 39°C selama 24 jam. Total produksi gas diamati sebagai indikator kualitas hijauan dan emisi gas metana diukur. Hasil menunjukkan bahwa, berdasarkan pendekatan bivariabel, hijauan yang menghasilkan emisi metana rendah umumnya juga memiliki kualitas yang kurang baik atau rendah total produksi gasnya, kecuali *Rhus typhina* yakni sebesar 43 ml metana/200 mg bahan kering. *Loading plot* dari PCA menunjukkan bahwa semua fraksi fenolik hijauan berlawanan arah dengan gas metana dan total produksi gas. Hijauan yang berlawanan arah dengan produksi metana adalah *Bergenia crassifolia*, *Swietenia mahagoni*, *Clidemia hirta*, *Peltiphyllum peltatum*, *Acacia villosa* dan *R. typhina*. Dapat disimpulkan bahwa, untuk hijauan mengandung tanin, pemilahan berdasarkan pendekatan univariabel, bivariabel dan multivariabel terkait emisi gas metana menunjukkan hasil yang serupa.

Kata kunci : Metana, kualitas hijauan, pemilahan, korelasi, tanin, fenolik

ABSTRACT

The aim of the present study was to conduct univariate, bivariate and multivariate (principal component analysis, PCA) approaches in the screening of tannin-containing plants from various collection sites for their CH₄ mitigating properties. Plant samples were obtained from various collection sites in different countries, i.e. Indonesia (n = 27 species), Mongolia (n = 14), Switzerland (n = 16) and Germany (n = 3). The plants were incubated *in vitro* with buffered-rumen fluid at 39°C for 24 h. Total gas production was recorded as an indicator of feed quality and emission of CH₄ was measured. Results showed that, based on bivariate screening, generally, plants possessed low CH₄ production had low quality or low total gas production except *Rhus typhina*, i.e. 43 ml/200 mg DM. The loading plot of PCA showed that all phenolic fractions were in the opposite direction with CH₄ and total gas production. Plants clustered together in reverse direction to that of CH₄ were *Bergenia crassifolia* root and leaf, *Swietenia mahagoni*, *Clidemia hirta*, *Peltiphyllum peltatum*, *Acacia villosa* and *R. typhina*. It was concluded that, for tannin-containing plants, screenings based on univariate, bivariate and multivariate approaches in relation to ruminal CH₄ emission led to similar results

Keywords : Methane, forage quality, screening, correlation, tannin, phenolic

INTRODUCTION

Forage quality is a determining factor for the productivity of ruminants. Several important properties related to forage quality are energy

content, fiber, protein, anti-nutritional compounds and the digestibility of forage in the digestive tract. On the other hand, the concern on global warming due to accumulation of green-house gases has increased in the last decades, including

methane (CH₄). Ruminants are among major contributors of atmospheric CH₄ which is generated from microbial fermentation in the rumen (Moss *et al.*, 2000), and may be more prevalent in the tropical area. It has been estimated that ruminants produce 80 million tons of CH₄ annually which accounts for 28% of total anthropogenic CH₄ emissions (Beauchemin *et al.*, 2008). Such emission is not only associated with environmental problem but also reflects the loss of energy from animal and, hence, may reduce the amount of energy for production purposes. Up to 12% of the gross energy from feed consumed by ruminants is lost as CH₄ (McCrabb and Hunter, 1999).

The above-mentioned factors have led to the exploration of forages that are not only good in quality but also are able to reduce CH₄ formation in the rumen. Tannin-containing forages or extracts have been reported to be able to reduce CH₄ emission from ruminants, both in *in vitro* and *in vivo* experiments (e.g., Tavendale *et al.*, 2005; Tiemann *et al.*, 2008; Bhatta *et al.*, 2009). However, further research still needs to be conducted to find most promising tannin-containing plants, i.e. effective in mitigating CH₄ emission at simultaneously of high quality. Screenings of plants that possess CH₄ mitigating properties have been done using different approaches, such as using univariate approach (Soliva *et al.*, 2008), bivariate (Bodas *et al.*, 2008; Garcia-Gonzalez *et al.*, 2008) and multivariate (Jayanegara *et al.*, 2011a). However, no studies so far have attempted to address these different approaches simultaneously when screening the plants and to compare the results obtained.

Based on this background, therefore, the aim of the present study is to conduct univariate, bivariate and multivariate approaches in the screening of tannin-containing plants from various collection sites for their CH₄ mitigating properties. A main point of interest is to observe whether these different approaches will lead to a different conclusion when screening the plants. Additionally, as a second objective, this study is aimed at observing the relationships between plant phenolic fractions and CH₄ emission using an extended database of plants collected from various sites, i.e. Indonesia, Mongolia, Switzerland and Germany.

MATERIALS AND METHODS

Plant Samples

Plant samples were obtained from various collection sites in different countries, i.e. Indonesia (n = 27 species), Mongolia (n = 14), Switzerland (n = 16) and Germany (n = 3). The data were based on some previous studies, i.e. Jayanegara *et al.* (2009a) for the plants from Mongolia and Germany, Jayanegara *et al.* (2011a) for the plants from Indonesia, and Jayanegara *et al.* (2011b) for the plants from Switzerland. Plant samples from Indonesia consisted of one grass, four herbs, nine shrubs and 13 tree species; all of them are either commonly used as ruminant feeds in rural areas or as traditional veterinary medicinal plants. The plants collected from Mongolia are used locally in the region of production as medicinal plants. Plants from Switzerland were obtained from the region of the Alps at altitudes of 800-2300 m above sea level, consisted of two grasses, eight non-leguminous herbs, three herbaceous legumes and three tree species. The majority of these plants is used for grazing of ruminants during European summer period, particularly for dairy production. The plants from Germany were collected from the Botanical Garden of the University of Hohenheim in Stuttgart. All plant samples were oven dried at 50-60°C and ground to pass 1 mm sieve. Each plant was encoded by its country of origin, i.e. I, M, S and G for plants obtained from Indonesia, Mongolia, Switzerland and Germany, respectively, and numbered according to alphabetical order within country.

Chemical Composition and *In Vitro* Procedures

The plant samples were analyzed for their chemical composition. These included crude protein (CP), ether extract (EE; AOAC, 1997), neutral detergent fiber (NDF), acid detergent fiber (ADF; Van Soest *et al.*, 1991), total phenols (TP), non-tannin phenols (NTP), total tannins (TT), condensed tannins (CT; Makkar, 2003a) and hydrolysable tannins (HT; Singh *et al.*, 2005). The plants were incubated *in vitro* with buffered-rumen fluid by using the Hohenheim gas test method (Menke and Steingass, 1988), incubated at 39°C for 24 h. In addition, standard hay and concentrate (obtained from the Institute of Animal Nutrition, University of Hohenheim, Stuttgart, Germany) were incubated, serving as control for the successful incubation. After the incubation,

total gas production was recorded as an indicator of feed quality. Emission of CH₄ was measured by using an infrared methane analyzer (Goel *et al.*, 2008) or a gas chromatograph (Soliva and Hess, 2007). These two different methods result in a similar CH₄ value of hay standard, i.e. between 160-170 ml/l gas.

Statistical Analysis

Data of all plants from various collection sites were tabulated into a database. Minimum and maximum values for each variable were recorded, and mean and standard deviation were calculated. Pearson's correlation analysis was performed to the data, i.e. the chemical composition and fermentation variables (total gas production and CH₄ emission). Screening approaches were based on univariate (CH₄ only), bivariate (total gas and CH₄) and multivariate (all variables investigated) using the principal component analysis (PCA). For the univariate approach, the plants considered were those which produced CH₄ ≤ average CH₄ across all plants minus 1 × standard deviation. For the bivariate approach, the screening was based on a two-dimensional plot between total gas and CH₄. For the multivariate approach, a PCA was applied to the data with Kaiser's criterion, i.e. Eigenvalue ≥ 1.0 to extract the principal components (PC) without rotation method. This procedure generated factor loading and factor score. The first two PC were plotted both for the loading and the score. The loading plot is used for describing the relationship among variables and the score plot is used for classifying the plants according to the loading. All statistical analyses were conducted using SPSS version 17.0 (2008) and the figures were generated using SigmaPlot version 11.0 (2008).

RESULTS

Chemical Composition

Summary of chemical composition of the plants investigated is presented in Table 1. Plant samples collected from Indonesia contained the highest CP contents (on average basis) compared to from other collection sites. Very high CP contents were found in the leaves of *Carica papaya* (386 g/kg DM; code I7) and *Manihot esculenta* (377 g/kg DM; I17) across all plants. All plants contained low EE with the average less

than 35 g/kg DM from the four collection sites. Plants from Mongolia contained higher ADF than the others. Quite a diverse TP contents were found across all collection sites, except those from Switzerland. The average of TP in these plants was 31 g/kg DM with the highest was found in *Hedysarum hedysaroides*, i.e. 69 g/kg DM (S8). This pattern was similarly observed for TT, CT and HT. Several plants, i.e. *Rhus typhina* (G2), *Acacia villosa* (I2) and *Clidemia hirta* (I8) contained TT more than 200 g/kg DM. Very high CT contents were found in *Vaccinium vitis idea* (M14) and *Swietenia mahagoni* (I27) with the values of 175 and 86 g/kg DM, respectively.

Relationships between Variables

Variables CP and EE were unrelated to total gas production and CH₄ (Table 2). Both NDF and ADF were positively correlated (both at P<0.001) with CH₄ emission, however, only ADF was negatively correlated with total gas (P<0.05). All phenolic fractions were negatively correlated with CH₄ by following the order of magnitude: TP > TT > HT > NTP > CT. The phenolics were also negatively correlated with total gas production.

Screening of Plants

Across all plants, the average CH₄ emission after 24 h *in vitro* incubation was 137 (±32) ml/l gas (Table 1). Therefore, for the screening based on univariate approach, the promising plants were those which produced CH₄ ≤ 105 ml/l gas. These plants were (ordered from lowest to highest CH₄) *Bergenia crassifolia* root (M4), *Peltiphyllum peltatum* (G1), *S. mahagoni* (I27), *A. villosa* (I2), *Eugenia aquea* (I11), *Myristica fragrans* (I21), *B. crassifolia* leaf (M3), *Pithecellobium jiringa* (I24), *R. typhina* (G2) and *C. hirta* (I8). Screening based on bivariate approach is presented in Figure 1. It appeared that, generally, plants possessed low CH₄ production had low quality, i.e. below the average of total gas production. Among plants that produced low CH₄, only *R. typhina* (G2) resulted in a high total gas production, i.e. 43 ml/200 mg DM. *Salsola laricifolia* (M8), *Hibiscus tiliaceus* (I12) and *Canna indica* (I6) had simultaneously high CH₄ emission and low total gas.

In relation to screening plants based on the multivariate approach, loading and score plots generated from PCA are shown in Figure 2. The PC1 and PC2 explained 42.3% and 23.3% of the

Table 1: Summary of Chemical Composition of Plants from Various Collection Sites and Ruminal Fermentation Variables *in vitro*

Item	CP	EE	NDF	ADF	TP	NTP	TT	CT	HT	Gas	CH ₄
	 (g/kg dry matter)									(ml)	(ml/l)
Indonesia (n = 27 species)											
Minimum	79	11	155	135	14	4	2	0	2	5.3	68
Maximum	386	65	710	476	236	102	220	86	206	42.8	185
Mean	215	30	371	281	90	26	65	21	44	23.4	127
SD	90	14	137	100	71	24	66	23	59	10.7	29
Mongolia (n = 14 species)											
Minimum	33	6	226	169	20	9	4	0	0	11.5	44
Maximum	246	35	626	509	320	144	178	175	164	44.8	220
Mean	114	21	444	352	94	43	51	20	33	27.5	153
SD	52	10	136	107	109	47	64	46	51	9.7	43
Switzerland (n = 16 species)											
Minimum	91	6	218	157	8	8	0	0	0	29.5	129
Maximum	247	35	753	344	69	35	38	21	34	49.9	176
Mean	156	19	379	263	31	20	12	3	9	40.6	148
SD	47	8	128	52	15	7	12	6	8	5.8	12
Germany (n = 3 species)											
Minimum	113	17	191	174	57	13	36	1	21	22.2	57
Maximum	169	56	322	185	222	53	209	16	208	49.3	125
Mean	141	31	244	181	160	29	131	11	120	38.2	95
SD	28	22	69	6	90	21	88	8	94	14.2	35
Overall (n = 60 species)											
Minimum	33	6	155	135	8	4	0	0	0	5.3	44
Maximum	386	65	753	509	320	144	220	175	208	49.9	220
Mean	172	25	384	288	79	28	51	15	36	30.0	137
SD	81	13	136	97	79	29	63	28	55	11.9	32

ADF, acid detergent fiber; CP, crude protein; CT, condensed tannins; EE, ether extract; HT, hydrolysable tannins; NDF, neutral detergent fiber; NTP, non-tannin phenols; SD, standard deviation; TP, total phenols; TT, total tannins

total variation, respectively. It is clear that all phenolic fractions were in the opposite direction with CH₄ and total gas production (Figure 2a). However, less clear relationships were observed between the other chemical constituents (CP, EE, NDF and ADF) and the *in vitro* fermentation variables. Some plants were clustered together in reverse direction to that of CH₄ (Figure 2b). These plants were *B. crassifolia* root (M4), *B. crassifolia* leaf (M3), *S. mahagoni* (I27), *C. hirta* (I8), *P. peltatum* (G1), *A. villosa* (I2) and *R.*

typhina (G2). Variable CH₄ and total gas production appeared to be in close direction in the loading plot of PCA.

DISCUSSION

Plant Phenolics and CH₄ Emission

Negative relationships between all phenolic fractions and CH₄ emission suggest the role of plant phenolics on mitigating the respective green-house gas under anaerobic condition in the

Table 2. Correlations between Chemical Composition of Plants and Total Gas Production and CH₄ Emission (n = 60)

Variable	Gas	CH ₄
CP	0.08	0.05
EE	-0.01	-0.18
NDF	-0.23	0.49***
ADF	-0.27*	0.45***
TP	-0.53***	-0.77***
NTP	-0.35**	-0.52***
TT	-0.51***	-0.73***
CT	-0.46***	-0.33*
HT	-0.36**	-0.67***

ADF, acid detergent fiber; CP, crude protein; CT, condensed tannins; EE, ether extract; HT, hydrolysable tannins; NDF, neutral detergent fiber; NTP, non-tannin phenols; TP, total phenols; TT, total tannins

* P<0.05; ** P<0.01; *** P<0.001

rumen, and this confirms some studies that had been reported previously (Tiemann *et al.*, 2008; Bhatta *et al.*, 2009). Phenolics, particularly tannins reduce CH₄ emission through several ways, i.e. through reduction in fiber digestion which decreases H₂ production, inhibition of the growth of methanogens (Tavendale *et al.*, 2005), and reduction of protozoa population where part of the methanogens are associated symbiotically on the surface or inside the micro-fauna (Vogels *et al.*, 1980). Within the tannin classes, based on the correlation data, both CT and HT contributed to lower CH₄ emission as also shown by Bhatta *et al.* (2009). It is hypothesized that lower CH₄ due to CT is more through reduction in fiber digestion as compared to the HT, confirmed by much stronger negative correlation of the former with the total gas production. This might be related to the fact that CT is resistant from microbial degradation in the rumen due to its aromatic structure, while HT is degradable due to the presence of carbohydrate in its structure (McSweeney *et al.*, 2001; Makkar, 2003b). Interestingly, the NTP plays also a role in lowering CH₄ with less adverse effect on the total gas production. This is in accordance to our




Figure 1: Bivariate approach for screening plants based on total gas produced and CH₄ emission. Vertical and horizontal lines show the average of total gas and CH₄ across all plants, respectively. I, plant samples from Indonesia (□); M, plant samples from Mongolia (●); S, plant samples from Switzerland (▲); G, plant samples from Germany (◇)

screening by univariate approach is not as selective as the others. One may obtain plants with low CH₄ production potential, but without considering the other important aspect, i.e. the forage quality. Among the promising plants obtained by the univariate screening, for instance, *A. villosa* (I2) and *E. aquea* (I11) resulted in poor quality performance as shown by the very low total gas production (≤ 10 ml/200 mg DM incubated) of both plants. The bivariate approach may overcome this problem by considering the forage quality related variable (Bodas *et al.*, 2008). Although almost all plants possessed low CH₄ emission had low quality, still, a good quality plant could be obtained, i.e. *R. typhina* (G2). What is clear from this approach is to avoid the utilization of plants that possessed high CH₄ emission and low total gas production such as *Salsola laricifolia* (M8), *Canna indica* (I6) and *Hibiscus tiliaceus* (I12).

The loading plot of PCA showed the relationships between variables in a two-dimensional plot. Variables that are in the same direction are positively correlated whereas variables that are in the opposite direction are negatively correlated (Esbensen, 2004). This apparently provides the same message as the correlation analysis, but not only restricted to pairwise comparisons. Since variable CH₄ and total gas production were in a close direction in the loading plot of PCA, it may imply that there is little probability in obtaining many tannin-containing plants with both desired characteristics, i.e. low CH₄ production potential and high forage quality. Therefore, the screening results of this approach were largely in agreement with the univariate approach. This is probably connected to the role of tannins in mitigating CH₄ emission as discussed above but reducing the forage quality as well at the same time. The adverse effects of tannins on forage digestibility have been well-documented (Makkar, 2003b; Mueller-Harvey, 2006). However, this does not exclude the possibility of finding plants with both desired characteristics, like *R. typhina* (G2) in the present dataset, since the structure and the activity of tannins in each plants are quite diverse (Mueller-Harvey, 2006).

CONCLUSION

Plant phenolic contents may be used as an

indicator for CH₄ mitigating properties of the plants, although they are also reversely related to the forage quality. Different approaches, i.e. univariate, bivariate and multivariate (PCA) for screening of tannin-containing plants in relation to their ruminal CH₄ emission *in vitro* led to similar results to certain extent. However, screening based on univariate approach has to be conducted cautiously since it does not consider the forage quality, which is also quite an important factor. The bivariate and multivariate approaches may overcome such problem by considering the forage quality related variable. Although almost all tannin-containing plants that possessed low CH₄ emission had low quality in the present study, still, a good quality plant could be obtained, i.e. *R. typhina*. This is possible since great variation have occurred in the structure and activity of each plant phenolics. The searching of plants with both desired characteristics is subjected to further research.

REFERENCES

- AOAC. 1997. Official Methods of Analysis. Association of Official Analytical Chemists, Arlington, USA.
- Beauchemin, K.A., M. Kreuzer, F. O'Mara and T.A. McAllister. 2008. Nutritional management for enteric methane abatement: a review. *Aust. J. Exp. Agric.* 48: 21-27.
- Bhatta, R., Y. Uyeno, K. Tajima, A. Takenaka, Y. Yabumoto, I. Nonaka, O. Enishi and M. Kurihara. 2009. Difference in the nature of tannins on *in vitro* ruminal methane and volatile fatty acid production and on methanogenic archaea and protozoal populations. *J. Dairy Sci.* 92: 5512-5522.
- Bodas, R., S. Lopez, M. Fernandez, R. Garcia-Gonzalez, A.B. Rodriguez, R.J. Wallace and J.S. Gonzalez. 2008. *In vitro* screening of the potential of numerous plant species as antimethanogenic feed additives for ruminants. *Anim. Feed Sci. Technol.* 145: 245-258.
- Esbensen, K.H. 2004. Multivariate Data Analysis - In Practice. 5th Edition. CAMO Software AS, Oslo, Norway.
- Garcia-Gonzalez, R., S. Lopez, M. Fernandez, R. Bodas and J.S. Gonzalez. 2008. Screening the activity of plants and spices for decreasing ruminal methane production *in vitro*. *Anim. Feed Sci. Technol.* 147: 36-52.

- Goel, G., H.P.S. Makkar and K. Becker. 2008. Effect of *Sesbania sesban* and *Carduus pycnocephalus* and Fenugreek (*Trigonella foenum-graecum* L.) seeds and their extracts on partitioning of nutrients from roughage- and concentrate-based feeds to methane. *Anim. Feed Sci. Technol.* 147: 72-89.
- Jayanegara, A., N. Togtokhbayar, H.P.S. Makkar and K. Becker. 2009a. Tannins determined by various methods as predictors of methane production reduction potential of plants by an *in vitro* rumen fermentation system. *Anim. Feed Sci. Technol.* 150: 230-237.
- Jayanegara, A., H.P.S. Makkar and K. Becker. 2009b. Methane reduction effect of simple phenolic acids evaluated by *in vitro* Hohenheim gas production method. *Proc. Soc. Nutr. Physiol.* 18:98.
- Jayanegara, A., E. Wina, C.R. Soliva, S. Marquardt, M. Kreuzer and F. Leiber. 2011a. Dependence of forage quality and methanogenic potential of tropical plants on their phenolic fractions as determined by principal component analysis. *Anim. Feed Sci. Technol.* 163: 231-243.
- Jayanegara, A., S. Marquardt, M. Kreuzer and F. Leiber. 2011b. Nutrient and energy content, *in vitro* ruminal fermentation characteristics and methanogenic potential of alpine forage plant species during early summer. *J. Sci. Food Agric.* 91: 1863-1870.
- Makkar, H.P.S. 2003a. Quantification of Tannins in Tree and Shrub Foliage: A Laboratory Manual. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Makkar, H.P.S. 2003b. Effects and fate of tannins in ruminant animals, adaptation to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Rum. Res.* 49: 241-256.
- McCrabb, G.J. and R.A. Hunter. 1999. Prediction of methane emissions from beef cattle in tropical production systems. *Aust. J. Agric. Res.* 50: 1335-1340.
- McSweeney, C.S., B. Palmer, D.M. McNeill and D.O. Krause. 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Anim. Feed Sci. Technol.* 91:83-93.
- Menke, K.H. and H. Steingass. 1988. Estimation of the energetic feed value obtained from chemical analysis and *in vitro* gas production using rumen fluid. *Anim. Res. Dev.* 28: 7-55.
- Moss, A.R., J.P. Jouany and J. Newbold. 2000. Methane production by ruminants: its contribution to global warming. *Ann. Zootech.* 49: 231-253.
- Mueller-Harvey, I. 2006. Unravelling the conundrum of tannins in animal nutrition and health. *J. Sci. Food Agric.* 86:2010-2037.
- SigmaPlot. 2008. SigmaPlot Software version 11.0. Systat Software Inc., San Jose, USA.
- Singh, B., A. Sahoo, R. Sharma and T.K. Bhat. 2005. Effect of polyethylene glycol on gas production parameters and nitrogen disappearance of some tree forages. *Anim. Feed Sci. Technol.* 123-124: 351-364.
- Soliva, C.R. and H.D. Hess. 2007. Measuring methane emission of ruminants by *in vitro* and *in vivo* techniques. In: Makkar, H.P.S. and P.E. Vercoe (Eds.). *Measuring Methane Production from Ruminants*. Springer, Dordrecht, The Netherlands.
- Soliva, C.R., A.B. Zeleke, C. Clement, H.D. Hess, V. Fievez and M. Kreuzer. 2008. *In vitro* screening of various tropical foliages, seeds, fruits and medicinal plants for low methane and high ammonia generating potentials in the rumen. *Anim. Feed Sci. Technol.* 147: 53-71.
- SPSS Inc. 2008. SPSS Statistics version 17.0. SPSS Inc., Chicago, USA.
- Tavendale, M.H., L.P. Meagher, D. Pacheco, N. Walker, G.T. Attwood and S. Sivakumaran. 2005. Methane production from *in vitro* rumen incubations with *Lotus pedunculatus* and *Medicago sativa*, and effects of extractable condensed tannin fractions on methanogenesis. *Anim. Feed Sci. Technol.* 123-124: 403-419.
- Tiemann, T.T., C.E. Lascano, H.R. Wettstein, A.C. Mayer, M. Kreuzer and H.D. Hess. 2008. Effect of the tropical tannin-rich shrub legumes *Calliandra calothyrsus* and *Flemingia macrophylla* on methane emission and nitrogen and energy balance in growing lambs. *Animal* 2: 790-799.
- Van Soest, P.J., J.B. Robertson and B.A. Lewis. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74: 3583-3597.
- Vogels, G.D., W.F. Hoppe and C.K. Stumm. 1980. Association of methanogenic bacteria with rumen ciliates. *Appl. Environ. Microbiol.* 40: 608-612.