

TWO NICE DETERMINANTAL EXPRESSIONS AND A RECURRENCE RELATION FOR THE APOSTOL–BERNOULLI POLYNOMIALS

FENG QI^{1,2,3} AND BAI-NI GUO⁴

¹Institute of Mathematics, Henan Polytechnic University,
Jiaozuo City, Henan Province, 454010, China

²College of Mathematics, Inner Mongolia University for Nationalities,
Tongliao City, Inner Mongolia Autonomous Region, 028043, China

³Department of Mathematics, College of Science, Tianjin Polytechnic
University, Tianjin City, 300387, China

E-mail: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

⁴School of Mathematics and Informatics, Henan Polytechnic University,
Jiaozuo City, Henan Province, 454010, China

E-mail: bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com

Abstract. In the paper, the authors establish two nice determinantal expressions and a recurrence relation for the Apostol–Bernoulli polynomials.

Key words and Phrases: Apostol–Bernoulli polynomial; determinantal expression; recurrence relation; determinant; derivative of a ratio between two functions.

Abstrak. Pada makalah ini, para penulis menyajikan dua pernyataan berbentuk determinan dan sebuah relasi rekurensi untuk suku banyak Apostol–Bernoulli.

Kata kunci: Suku banyak Apostol–Bernoulli, ekspresi berbentuk determinan, relasi rekurensi, determinan, turunan dari rasio dua fungsi.

1. INTRODUCTION

It is well-known that the Bernoulli numbers B_k , the Bernoulli polynomials $B_k(u)$, and the Apostol–Bernoulli polynomials $B_k(u, z)$ for $k \geq 0$ can be generated respectively by

$$\frac{x}{e^x - 1} = \sum_{k=0}^{\infty} B_k \frac{x^k}{k!} = 1 - \frac{x}{2} + \sum_{k=1}^{\infty} B_{2k} \frac{x^{2k}}{(2k)!}, \quad |x| < 2\pi,$$

2000 Mathematics Subject Classification: Primary 11B68; Secondary 11B83, 11C20, 15A15, 26A06, 26A09, 33B10.

Received: 01-08-2016, revised: 03-03-2017, accepted: 04-03-2017.

$$\frac{xe^{ux}}{e^x - 1} = \sum_{k=0}^{\infty} B_k(u) \frac{x^k}{k!}, \quad |x| < 2\pi,$$

and

$$\frac{xe^{ux}}{ze^x - 1} = \sum_{k=0}^{\infty} B_k(u, z) \frac{x^k}{k!}, \quad |x| < \begin{cases} 2\pi, & z = 1; \\ |\ln z|, & z \neq 1. \end{cases} \quad (1)$$

It is clear that these notions have the relations

$$B_k = B_k(0) \quad \text{and} \quad B_k(u) = B_k(u, 1).$$

In [1, 2], Apostol connected special values of the Lerch zeta functions with the Apostol–Bernoulli polynomials $B_k(u, z)$. In [8], Luo gave a relation between the λ -power sums and the Apostol–Bernoulli polynomials $B_k(u, z)$, which generalize J. Bernoulli's formula on the representation of power sums in terms of the Bernoulli polynomials $B_k(u)$. In [7], Kim and Hu obtained the sums of products identity for the Apostol–Bernoulli numbers $B_k(u, z)$, which is an analogue of the classical sums of products identity for the Bernoulli numbers B_k dating back to Euler.

Let $p = p(x)$ and $q = q(x) \neq 0$ be two differentiable functions. Then

$$\frac{d^k}{dz^k} \left[\frac{p(x)}{q(x)} \right] = \frac{(-1)^k}{q^{k+1}} \begin{vmatrix} p & q & 0 & \cdots & 0 \\ p' & q' & q & \cdots & 0 \\ p'' & q'' & 2q' & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ p^{(k-1)} & q^{(k-1)} & \binom{k-1}{1} q^{(k-2)} & \cdots & q \\ p^{(k)} & q^{(k)} & \binom{k}{1} q^{(k-1)} & \cdots & \binom{k}{k-1} q' \end{vmatrix}, \quad k \geq 0. \quad (2)$$

See [3, p. 40]. We can rewrite the formula (2) as

$$\frac{d^k}{dx^k} \left[\frac{p(x)}{q(x)} \right] = \frac{(-1)^k}{q^{k+1}(x)} |W_{(k+1) \times (k+1)}(x)|, \quad (3)$$

where $|W_{(k+1) \times (k+1)}(x)|$ denotes the determinant of the $(k+1) \times (k+1)$ matrix

$$W_{(k+1) \times (k+1)}(x) = (U_{(k+1) \times 1}(x) \quad V_{(k+1) \times k}(x)),$$

the quantity $U_{(k+1) \times 1}(x)$ is a $(k+1) \times 1$ matrix whose elements $u_{\ell,1}(x) = p^{(\ell-1)}(x)$ for $1 \leq \ell \leq k+1$, and $V_{(k+1) \times k}(x)$ is a $(k+1) \times k$ matrix whose elements

$$v_{i,j}(x) = \begin{cases} \binom{i-1}{j-1} q^{(i-j)}(x), & i-j \geq 0 \\ 0, & i-j < 0 \end{cases}$$

for $1 \leq i \leq k+1$ and $1 \leq j \leq k$. For more information, please refer to related texts in the recently published papers [6, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 25, 26] and the closely related references therein.

The determinant expressions for the classical Bernoulli polynomials $B_k(u)$ have a long history, see [5, p. 53].

Applying the formula (3) to $p(x) = 1$ and $q(x) = \int_0^1 e^{x(s-u)} ds$, Qi and Chapman [13, Theorem 1.2] obtained the determinantal expressions

$$B_k(u) = (-1)^k \left| \frac{1}{\ell+1} \binom{\ell+1}{m} [(1-u)^{\ell-m+1} - (-u)^{\ell-m+1}] \right|_{1 \leq \ell \leq k, 0 \leq m \leq k-1}$$

and

$$B_k = (-1)^k \left| \frac{1}{\ell+1} \binom{\ell+1}{m} \right|_{1 \leq \ell \leq k, 0 \leq m \leq k-1},$$

where $|\cdot|_{1 \leq \ell \leq k, 0 \leq m \leq k-1}$ denotes a $k \times k$ determinant.

Influenced by the paper [13], Hu and Kim [6, Theorem 1.10] established

$$B_{k+1}(u, z) = \frac{(-1)^k(k+1)}{(z-1)^{k+1}} \left| \binom{\ell}{m} [z(z-u)^{\ell-m} - (-u)^{\ell-m}] \right|_{1 \leq \ell \leq k, 0 \leq m \leq k-1} \quad (4)$$

and

$$B_{k+1}(z) = \frac{(-1)^k(k+1)}{(z-1)^{k+1}} \left| \binom{\ell}{m} (z - \delta_{\ell,m}) \right|_{1 \leq \ell \leq k, 0 \leq m \leq k-1} \quad (5)$$

for $k \in \mathbb{N}$ and $z \neq 1$, where

$$\delta_{\ell,m} = \begin{cases} 1, & \ell = m \\ 0, & \ell \neq m \end{cases}$$

denotes the Kronecker delta. When deriving (4) and (5), Hu and Kim used the Leibnitz theorem for differentiation of a product and applied the formula (3) to $p(x) = 1$ and $q(x) = ze^{(1-u)x} - e^{-ux}$.

In this paper, we will apply the formula (3) again to establish two nice determinantal expressions and consequently derive a recurrence relation of the Apostol–Bernoulli polynomials $B_k(u, z)$ for $k \in \mathbb{N}$ and $z \neq 1$.

2. MAIN RESULTS AND THEIR PROOFS

Our main results, two nice determinantal expressions and a recurrence relation of the Apostol–Bernoulli polynomials $B_k(u, z)$ for $k \in \mathbb{N}$ and $z \neq 1$, can be stated as the following theorem.

Theorem 2.1. *The Apostol–Bernoulli polynomials $B_k(u, z)$ for $k \in \mathbb{N}$ and $z \neq 1$ can be determinantly expressed by*

$$B_k(u, z) = \frac{(-1)^{k-1}k}{(z-1)^k} \begin{vmatrix} 1 & z-1 & 0 & \cdots & 0 & 0 & 0 \\ u & z & z-1 & \cdots & 0 & 0 & 0 \\ u^2 & z & \binom{2}{1}z & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ u^{k-3} & z & \binom{k-3}{1}z & \cdots & \binom{k-3}{k-4}z & z-1 & 0 \\ u^{k-2} & z & \binom{k-2}{1}z & \cdots & \binom{k-2}{k-4}z & \binom{k-2}{k-3}z & z-1 \\ u^{k-1} & z & \binom{k-1}{1}z & \cdots & \binom{k-1}{k-4}z & \binom{k-1}{k-3}z & \binom{k-1}{k-2}z \end{vmatrix} \quad (6)$$

and

$$B_k(u, z) = \frac{(-1)^{k+1}}{(z-1)^k} \begin{vmatrix} 1 & z-1 & 0 & \cdots & 0 & 0 \\ 2u & \binom{2}{1}z & z-1 & \cdots & 0 & 0 \\ 3u^2 & \binom{3}{1}z & \binom{3}{2}z & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ (k-2)u^{k-3} & \binom{k-2}{1}z & \binom{k-2}{2}z & \cdots & z-1 & 0 \\ (k-1)u^{k-2} & \binom{k-1}{1}z & \binom{k-1}{2}z & \cdots & \binom{k-1}{k-2}z & z-1 \\ ku^{k-1} & \binom{k}{1}z & \binom{k}{2}z & \cdots & \binom{k}{k-2}z & \binom{k}{k-1}z \end{vmatrix}. \quad (7)$$

Consequently, the Apostol–Bernoulli polynomials $B_k(u, z)$ for $k \in \mathbb{N}$ and $z \neq 1$ satisfy the recurrence relation

$$B_k(u, z) = \frac{z}{1-z} \left[kB_{k-1}(u, z) + \sum_{r=1}^{k-2} \binom{k}{r} B_r(u, z) - \frac{ku^{k-1}}{z} \right], \quad k \geq 2. \quad (8)$$

Proof of Theorem 2.1. Applying the formula (3) to $p(x) = xe^{ux}$ and $q(x) = ze^x - 1$ for $z \neq 1$ gives

$$u_{\ell,1}(x) = (xe^{ux})^{(\ell-1)} \rightarrow (\ell-1)u^{\ell-2}, \quad x \rightarrow 0$$

for $\ell \geq 1$ and

$$v_{i,j}(x) = \begin{cases} \binom{i-1}{j-1} (ze^x - 1)^{(i-j)}, & i-j \geq 0 \\ 0, & i-j < 0 \end{cases} \rightarrow \begin{cases} z-1, & i-j = 0 \\ \binom{i-1}{j-1} z, & i-j > 0 \\ 0, & i-j < 0 \end{cases} \quad (9)$$

as $x \rightarrow 0$ for $i, j \geq 1$. As a result, the Apostol–Bernoulli polynomials $B_k(u, z)$ for $k \geq 1$ can be expressed as

$$B_k(u, z) = \lim_{x \rightarrow 0} \frac{d^k}{dx^k} \left(\frac{xe^{ux}}{ze^x - 1} \right) \\ = \frac{(-1)^k}{(z-1)^{k+1}} \begin{vmatrix} 0 & z-1 & 0 & \cdots & 0 & 0 & 0 \\ 1 & z & z-1 & \cdots & 0 & 0 & 0 \\ 2u & z & \binom{2}{1}z & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ (k-2)u^{k-3} & z & \binom{k-2}{1}z & \cdots & \binom{k-2}{k-3}z & z-1 & 0 \\ (k-1)u^{k-2} & z & \binom{k-1}{1}z & \cdots & \binom{k-1}{k-3}z & \binom{k-1}{k-2}z & z-1 \\ ku^{k-1} & z & \binom{k}{1}z & \cdots & \binom{k}{k-3}z & \binom{k}{k-2}z & \binom{k}{k-1}z \end{vmatrix}$$

$$= \frac{(-1)^{k+1}}{(z-1)^k} \begin{vmatrix} 1 & z-1 & 0 & \cdots & 0 & 0 & 0 \\ 2u & \binom{2}{1}z & z-1 & \cdots & 0 & 0 & 0 \\ 3u^2 & \binom{3}{1}z & \binom{3}{2}z & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ (k-2)u^{k-3} & \binom{k-2}{1}z & \binom{k-2}{2}z & \cdots & \binom{k-2}{k-3}z & z-1 & 0 \\ (k-1)u^{k-2} & \binom{k-1}{1}z & \binom{k-1}{2}z & \cdots & \binom{k-1}{k-3}z & \binom{k-1}{k-2}z & z-1 \\ ku^{k-1} & \binom{k}{1}z & \binom{k}{2}z & \cdots & \binom{k}{k-3}z & \binom{k}{k-2}z & \binom{k}{k-1}z \end{vmatrix}.$$

The determinantal expression (7) is thus proved.

Since $B_0(u, z) = 0$ for $z \neq 1$, the equation (1) can be rewritten as

$$\frac{e^{ux}}{ze^x - 1} = \sum_{k=0}^{\infty} B_{k+1}(u, z) \frac{x^k}{(k+1)!}, \quad |x| < \begin{cases} 2\pi, & z = 1; \\ |\ln z|, & z \neq 1. \end{cases}$$

This implies that

$$\frac{B_{k+1}(u, z)}{k+1} = \lim_{x \rightarrow 0} \frac{d^k}{dx^k} \left(\frac{e^{ux}}{ze^x - 1} \right), \quad k \geq 0.$$

Further applying the formula (3) to $p(x) = e^{ux}$ and $q(x) = ze^x - 1$ for $z \neq 1$ gives

$$u_{\ell,1}(x) = (e^{ux})^{(\ell-1)} \rightarrow u^{\ell-1}, \quad x \rightarrow 0$$

for $\ell \geq 1$ and (9). Therefore, we have

$$\frac{B_{k+1}(u, z)}{k+1} = \frac{(-1)^k}{(z-1)^{k+1}} \begin{vmatrix} 1 & z-1 & 0 & \cdots & 0 & 0 & 0 \\ u & z & z-1 & \cdots & 0 & 0 & 0 \\ u^2 & z & \binom{2}{1}z & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ u^{k-2} & z & \binom{k-2}{1}z & \cdots & \binom{k-2}{k-3}z & z-1 & 0 \\ u^{k-1} & z & \binom{k-1}{1}z & \cdots & \binom{k-1}{k-3}z & \binom{k-1}{k-2}z & z-1 \\ u^k & z & \binom{k}{1}z & \cdots & \binom{k}{k-3}z & \binom{k}{k-2}z & \binom{k}{k-1}z \end{vmatrix}$$

for $k \geq 0$. The determinantal expression (6) is thus proved.

Let $M_0 = 1$ and

$$M_n = \begin{vmatrix} m_{1,1} & m_{1,2} & 0 & \cdots & 0 & 0 \\ m_{2,1} & m_{2,2} & m_{2,3} & \cdots & 0 & 0 \\ m_{3,1} & m_{3,2} & m_{3,3} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ m_{n-2,1} & m_{n-2,2} & m_{n-2,3} & \cdots & m_{n-2,n-1} & 0 \\ m_{n-1,1} & m_{n-1,2} & m_{n-1,3} & \cdots & m_{n-1,n-1} & m_{n-1,n} \\ m_{n,1} & m_{n,2} & m_{n,3} & \cdots & m_{n,n-1} & m_{n,n} \end{vmatrix}$$

for $n \in \mathbb{N}$. In [4, p. 222, Theorem], it was proved that the sequence M_n for $n \geq 0$ satisfies $M_1 = m_{1,1}$ and

$$M_n = m_{n,n}M_{n-1} + \sum_{r=1}^{n-1} (-1)^{n-r} m_{n,r} \left(\prod_{j=r}^{n-1} m_{j,j+1} \right) M_{r-1}, \quad n \geq 2.$$

See also [17, Lemma 2], [18, Lemma 5], [24, Lemma 2], and [25, Remark 3]. Applying this conclusion to determinants in (6) and (7) readily produces the same recurrence relation (8) respectively. The proof of Theorem 2.1 is complete. \square

Remark 2.1. *This paper is a slightly modified version of the preprint [11].*

Acknowledgements. The authors appreciate the anonymous referees for their careful corrections to and valuable comments on the original version of this paper.

REFERENCES

- [1] T. M. Apostol, *Addendum to 'On the Lerch zeta function'*, Pacific J. Math. **2** (1952), 10–10; Available online at <http://projecteuclid.org/euclid.pjm/1103051938>.
- [2] T. M. Apostol, *On the Lerch zeta function* Pacific J. Math. **1** (1951), 161–167; Available online at <http://projecteuclid.org/euclid.pjm/1103052188>.
- [3] N. Bourbaki, *Functions of a Real Variable, Elementary Theory*, Translated from the 1976 French original by Philip Spain. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004; Available online at <http://dx.doi.org/10.1007/978-3-642-59315-4>.
- [4] N. D. Cahill, J. R. D'Errico, D. A. Narayan, and J. Y. Narayan, *Fibonacci determinants*, College Math. J. **3** (2002), 221–225; Available online at <http://dx.doi.org/10.2307/1559033>.
- [5] J. W. L. Glaisher, *Expressions for Laplace's coefficients, Bernoullian and Eulerian numbers, etc., as determinants*, Messenger Math. (2) **6** (1875), 49–63.
- [6] S. Hu and M.-S. Kim, *Two closed forms for the Apostol-Bernoulli polynomials*, arXiv preprint (2015), available online at <http://arxiv.org/abs/1509.04190>.
- [7] M.-S. Kim and S. Hu, *Sums of products of Apostol-Bernoulli numbers*, Ramanujan J. **28** (2012), no. 1, 113–123; Available online at <http://dx.doi.org/10.1007/s11139-011-9340-z>.
- [8] Q.-M. Luo, *The multiplication formulas for the Apostol-Bernoulli and Apostol-Euler polynomials of higher order*, Integral Transforms Spec. Funct. **20** (2009), no. 5-6, 377–391; Available online at <http://dx.doi.org/10.1080/10652460802564324>.
- [9] F. Qi, *A determinantal representation for derangement numbers*, Glob. J. Math. Anal. **4** (2016), no. 3, 17–17; Available online at <http://dx.doi.org/10.14419/gjma.v4i3.6574>.
- [10] F. Qi, *Derivatives of tangent function and tangent numbers*, Appl. Math. Comput. **268** (2015), 844–858; Available online at <http://dx.doi.org/10.1016/j.amc.2015.06.123>.
- [11] F. Qi, *Two nice determinantal expressions for the Apostol-Bernoulli polynomials*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.2.33797.76004>.
- [12] F. Qi, V. Čerňanová, and Y. S. Semenov, *On tridiagonal determinants and the Cauchy product of central Delannoy numbers*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.1.3772.6967>.
- [13] F. Qi and R. J. Chapman, *Two closed forms for the Bernoulli polynomials*, J. Number Theory **159** (2016), 89–100; Available online at <http://dx.doi.org/10.1016/j.jnt.2015.07.021>.
- [14] F. Qi and B.-N. Guo, *A determinantal expression and a recurrence relation for the Euler polynomials*, Preprints **2016**, 2016100034, 8 pages; Available online at <http://dx.doi.org/10.20944/preprints201610.0034.v1>.
- [15] F. Qi and B.-N. Guo, *Explicit and recursive formulas, integral representations, and properties of the large Schröder numbers*, Kragujevac J. Math. **41** (2017), no. 1, 121–141.

- [16] F. Qi and B.-N. Guo, *Expressing the generalized Fibonacci polynomials in terms of a tridiagonal determinant*, *Matematiche (Catania)* **72** (2017), no. 1, in press.
- [17] F. Qi and B.-N. Guo, *Some determinantal expressions and recurrence relations of the Bernoulli polynomials*, *Mathematics* **4** (2016), no. 4, Article 65, 11 pages; Available online at <http://dx.doi.org/10.3390/math4040065>.
- [18] F. Qi and B.-N. Guo, *Some properties of a solution to a family of inhomogeneous linear ordinary differential equations*, *Preprints* **2016**, 2016110146, 11 pages; Available online at <http://dx.doi.org/10.20944/preprints201611.0146.v1>.
- [19] F. Qi, M. Mahmoud, X.-T. Shi, and F.-F. Liu, *Some properties of the Catalan–Qi function related to the Catalan numbers*, *SpringerPlus* (2016), **5**:1126, 20 pages; Available online at <http://dx.doi.org/10.1186/s40064-016-2793-1>.
- [20] F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, *Integers* **16** (2016), Paper No. A23, 15 pages.
- [21] F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin, *Several formulas for special values of the Bell polynomials of the second kind and applications*, *J. Appl. Anal. Comput.* **7** (2017), no. 3, in press.
- [22] F. Qi, J.-L. Wang, and B.-N. Guo, *A recovery of two determinantal representations for derangement numbers*, *Cogent Math.* (2016), **3**: 1232878, 7 pages; Available online at <http://dx.doi.org/10.1080/23311835.2016.1232878>.
- [23] F. Qi, J.-L. Wang, and B.-N. Guo, *A representation for derangement numbers in terms of a tridiagonal determinant*, *Kragujevac J. Math.* **42** (2018), no. 1, 7–14.
- [24] F. Qi and J.-L. Zhao, *Some properties of the Bernoulli numbers of the second kind and their generating function*, *J. Differ. Equ. Appl.* (2017), in press.
- [25] F. Qi, J.-L. Zhao, and B.-N. Guo, *Closed forms for derangement numbers in terms of the Hessenberg determinants*, *Preprints* **2016**, 2016100035, 11 pages; Available online at <http://dx.doi.org/10.20944/preprints201610.0035.v1>.
- [26] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, *J. Inequal. Appl.* **2015**, **2015**:219, 8 pages; Available online at <http://dx.doi.org/10.1186/s13660-015-0738-9>.

